



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2024 – 06:56 PM EDT

PDB ID : 6CUU  
Title : Thermus thermophiles RNA polymerase in complex with promoter DNA and antibiotic Kanglemycin A  
Authors : Molodtsov, V.; Murakami, K.S.  
Deposited on : 2018-03-26  
Resolution : 2.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	
1	B	315	
2	C	1119	
3	D	1524	
4	E	99	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	F	423	
6	G	22	
7	H	27	

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 28403 atoms, of which 62 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	227	Total 1787	C 1141	N 311	O 333	S 2	0	0	0
1	B	223	Total 1758	C 1124	N 305	O 327	S 2	0	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	1112	Total 8717	C 5517	N 1555	O 1622	S 23	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	958	THR	PRO	conflict	UNP Q72HM5

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	1485	Total 11652	C 7391	N 2046	O 2180	S 35	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	274	ARG	GLN	conflict	UNP Q72HM6
D	1041	LEU	MET	conflict	UNP Q72HM6
D	1313	VAL	ALA	conflict	UNP Q72HM6

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	E	94	761	486	132	139	4	0	0	0

- Molecule 5 is a protein called RNA polymerase sigma factor SigA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	F	346	2793	1763	508	518	4	0	0	0


- Molecule 6 is a DNA chain called DNA (5'-D(P\*TP\*GP\*CP\*AP\*TP\*CP\*AP\*GP\*AP\*GP\*CP\*CP\*CP\*AP\*AP\*AP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
6	G	17	348	165	69	97	17	0	0	0

- Molecule 7 is a DNA chain called DNA (5'-D(\*TP\*AP\*TP\*AP\*AP\*TP\*GP\*GP\*GP\*AP\*GP\*CP\*CP\*TP\*CP\*TP\*GP\*AP\*TP\*GP\*CP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
7	H	22	451	216	84	130	21	0	0	0

- Molecule 8 is Kanglemycin A (three-letter code: KNG) (formula: C<sub>50</sub>H<sub>67</sub>NO<sub>19</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
8	C	1	132	50	62	1	19	0	0

- Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
9	D	2	2	2	0	0


- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
10	D	2	2	2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase subunit alpha




- Molecule 1: DNA-directed RNA polymerase subunit alpha




- Molecule 2: DNA-directed RNA polymerase subunit beta






• Molecule 3: DNA-directed RNA polymerase subunit beta'








• Molecule 4: DNA-directed RNA polymerase subunit omega




• Molecule 5: RNA polymerase sigma factor SigA






• Molecule 6: DNA (5'-D(P\*TP\*GP\*CP\*AP\*TP\*CP\*AP\*GP\*AP\*GP\*CP\*CP\*CP\*AP\*AP\*A P\*A)-3')



• Molecule 7: DNA (5'-D(\*TP\*AP\*TP\*AP\*AP\*TP\*GP\*GP\*GP\*AP\*GP\*CP\*CP\*TP\*CP\*TP \*GP\*AP\*TP\*GP\*CP\*A)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	185.22Å 101.31Å 294.28Å 90.00° 98.89° 90.00°	Depositor
Resolution (Å)	46.30 – 2.99 46.31 – 2.99	Depositor EDS
% Data completeness (in resolution range)	98.7 (46.30-2.99) 98.7 (46.31-2.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 3.01Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.201 , 0.248 0.203 , 0.244	Depositor DCC
$R_{free}$ test set	1990 reflections (1.85%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	77.4	Xtrriage
Anisotropy	0.501	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 70.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	28403	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	95.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, KNG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/1819	0.61	0/2473
1	B	0.34	0/1790	0.64	0/2435
2	C	0.33	0/8883	0.60	2/12022 (0.0%)
3	D	0.35	1/11857 (0.0%)	0.61	3/16040 (0.0%)
4	E	0.33	0/775	0.55	0/1045
5	F	0.31	0/2838	0.56	0/3820
6	G	0.96	1/391 (0.3%)	1.25	4/600 (0.7%)
7	H	0.83	0/505	1.30	6/776 (0.8%)
All	All	0.37	2/28858 (0.0%)	0.64	15/39211 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	875	THR	C-N	5.32	1.46	1.34
6	G	12	DG	C3'-O3'	-5.04	1.37	1.44

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	H	23	DG	O4'-C4'-C3'	-8.67	100.80	106.00
7	H	12	DC	O4'-C1'-N1	-8.29	102.20	108.00
6	G	14	DC	O4'-C1'-N1	-8.17	102.28	108.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	6	DA	O4'-C4'-C3'	-7.08	101.67	104.50
7	H	20	DG	O4'-C1'-N9	-7.04	103.07	108.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	422	ARG	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1787	0	1839	54	0
1	B	1758	0	1808	77	0
2	C	8717	0	8787	279	0
3	D	11652	0	11828	360	2
4	E	761	0	778	33	0
5	F	2793	0	2863	89	1
6	G	348	0	190	15	0
7	H	451	0	251	17	0
8	C	70	62	0	6	0
9	D	2	0	0	0	0
10	D	2	0	0	0	0
All	All	28341	62	28344	821	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 821 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:678:PRO:HA	2:C:683:ASN:HD21	1.14	1.12
3:D:806:PHE:HB2	3:D:829:VAL:HG22	1.29	1.10
3:D:1048:PRO:O	3:D:1079:LYS:NZ	1.87	1.06

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:669:ASN:ND2	5:F:420:ASP:OD1	1.88	1.05
6:G:6:DA:H8	6:G:6:DA:H5'	1.19	1.03

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:34:TYR:OH	3:D:327:GLU:OE1[4_1359]	1.95	0.25
3:D:1287:GLU:OE1	5:F:302:LYS:NZ[3_445]	2.03	0.17

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/315 (71%)	222 (99%)	3 (1%)	0	100	100
1	B	221/315 (70%)	217 (98%)	4 (2%)	0	100	100
2	C	1108/1119 (99%)	1080 (98%)	22 (2%)	6 (0%)	29	68
3	D	1481/1524 (97%)	1444 (98%)	34 (2%)	3 (0%)	47	82
4	E	92/99 (93%)	89 (97%)	2 (2%)	1 (1%)	14	50
5	F	344/423 (81%)	337 (98%)	5 (2%)	2 (1%)	25	64
All	All	3471/3795 (92%)	3389 (98%)	70 (2%)	12 (0%)	41	76

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	766	GLU
2	C	156	GLY
5	F	322	GLY
2	C	477	GLY
3	D	276	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/273 (73%)	194 (98%)	5 (2%)	47	79
1	B	196/273 (72%)	194 (99%)	2 (1%)	76	91
2	C	922/941 (98%)	887 (96%)	35 (4%)	33	69
3	D	1233/1279 (96%)	1184 (96%)	49 (4%)	31	68
4	E	83/88 (94%)	83 (100%)	0	100	100
5	F	297/370 (80%)	288 (97%)	9 (3%)	41	75
All	All	2930/3224 (91%)	2830 (97%)	100 (3%)	37	72

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	632	VAL
3	D	972	LEU
5	F	420	ASP
3	D	650	LEU
3	D	810	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
5	F	269	ASN
3	D	1441	GLN
3	D	709	HIS
3	D	1333	HIS
3	D	350	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	KNG	C	2001	-	75,75,75	3.82	28 (37%)	107,114,114	2.38	28 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	KNG	C	2001	-	-	23/76/113/113	0/5/6/6

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	C	2001	KNG	O18-C46	-14.12	1.21	1.44
8	C	2001	KNG	O03-C06	11.12	1.58	1.37
8	C	2001	KNG	O17-C47	-10.88	1.20	1.43
8	C	2001	KNG	C04-C10	8.42	1.57	1.43
8	C	2001	KNG	O16-C37	7.74	1.62	1.42

The worst 5 of 28 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	C	2001	KNG	C33-C24-C23	-8.30	95.10	111.43
8	C	2001	KNG	C25-O07-C35	-7.77	105.64	117.72
8	C	2001	KNG	O03-C06-C07	7.50	133.87	121.16
8	C	2001	KNG	C23-C24-C25	5.65	121.36	110.58
8	C	2001	KNG	C18-C19-C20	5.16	124.92	114.56

There are no chirality outliers.

5 of 23 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	C	2001	KNG	C21-C22-C23-O19
8	C	2001	KNG	C23-C24-C25-O07
8	C	2001	KNG	C33-C24-C25-C26
8	C	2001	KNG	C33-C24-C25-O07
8	C	2001	KNG	O06-C27-C28-C29

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	C	2001	KNG	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	227/315 (72%)	0.05	1 (0%) 92 79	67, 92, 127, 170	0
1	B	223/315 (70%)	0.21	3 (1%) 77 51	64, 101, 158, 189	0
2	C	1112/1119 (99%)	0.30	92 (8%) 11 3	40, 82, 174, 268	0
3	D	1485/1524 (97%)	0.16	83 (5%) 24 8	36, 80, 163, 253	0
4	E	94/99 (94%)	0.19	7 (7%) 14 4	47, 83, 155, 175	0
5	F	346/423 (81%)	0.26	20 (5%) 23 7	50, 102, 172, 240	0
6	G	17/22 (77%)	-0.35	0 100 100	75, 94, 187, 191	0
7	H	22/27 (81%)	-0.28	1 (4%) 33 12	79, 113, 202, 219	0
All	All	3526/3844 (91%)	0.21	207 (5%) 22 7	36, 86, 167, 268	0

The worst 5 of 207 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	64	LEU	9.0
5	F	149	GLU	8.8
2	C	311	PHE	8.3
3	D	1497	GLU	7.9
5	F	148	LYS	7.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	KNG	C	2001	70/70	0.86	0.47	87,141,186,187	0
9	ZN	D	2002	1/1	0.96	0.09	91,91,91,91	0
10	MG	D	2003	1/1	0.96	0.56	120,120,120,120	0
10	MG	D	2004	1/1	0.98	0.53	70,70,70,70	0
9	ZN	D	2001	1/1	0.99	0.20	65,65,65,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.