



# Full wwPDB X-ray Structure Validation Report i

Oct 23, 2021 – 11:51 AM EDT

PDB ID : 1CV7  
Title : Crystal structure of enhanced cyan-emission variant of GFP  
Authors : O'Brien, P.; Elsliger, M.A.; Kallio, K.; Remington, S.J.  
Deposited on : 1999-08-23  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	226	7%	45%	42%	12%

## 2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 1768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GREEN FLUORESCENT PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C 1738	N 1121	O 286	S 325	6	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	ARG	LYS	engineered mutation	UNP P42212
A	64	LEU	PHE	engineered mutation	UNP P42212
A	66	CRF	SER	chromophore	UNP P42212
A	66	CRF	TYR	chromophore	UNP P42212
A	66	CRF	GLY	chromophore	UNP P42212
A	146	ILE	ASN	engineered mutation	UNP P42212
A	153	THR	MET	engineered mutation	UNP P42212
A	163	ALA	VAL	engineered mutation	UNP P42212
A	164	HIS	ASN	engineered mutation	UNP P42212


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
2	A	30	Total 30	O 30	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GREEN FLUORESCENT PROTEIN



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.05 Å   62.78 Å   66.83 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	19.50 – 2.50 19.52 – 2.49	Depositor EDS
% Data completeness (in resolution range)	97.4 (19.50-2.50) 97.2 (19.52-2.49)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.74 (at 2.50 Å)	Xtriage
Refinement program	TNT	Depositor
$R$ , $R_{free}$	0.186 , (Not available) 0.192 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.6	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.20 , 85.0	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.48$ , $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1768	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CRF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.84	8/1755 (0.5%)	1.29	27/2377 (1.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	34	GLU	CD-OE2	6.33	1.32	1.25
1	A	90	GLU	CD-OE2	5.88	1.32	1.25
1	A	115	GLU	CD-OE2	5.75	1.31	1.25
1	A	111	GLU	CD-OE2	5.69	1.31	1.25
1	A	17	GLU	CD-OE2	5.65	1.31	1.25
1	A	32	GLU	CD-OE2	5.62	1.31	1.25
1	A	142	GLU	CD-OE2	5.51	1.31	1.25
1	A	95	GLU	CD-OE2	5.26	1.31	1.25

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	ASP	CB-CG-OD2	-8.96	110.24	118.30
1	A	55	VAL	C-N-CD	-8.72	101.43	120.60
1	A	173	ASP	CB-CG-OD2	-7.26	111.76	118.30
1	A	102	ASP	CB-CG-OD2	-7.10	111.91	118.30
1	A	82	ASP	CB-CG-OD1	6.87	124.48	118.30
1	A	36	ASP	CB-CG-OD2	-6.75	112.23	118.30
1	A	173	ASP	CB-CG-OD1	6.63	124.27	118.30
1	A	21	ASP	CB-CG-OD2	-6.50	112.45	118.30
1	A	97	THR	N-CA-CB	6.46	122.57	110.30
1	A	129	ASP	CB-CG-OD1	6.32	123.99	118.30
1	A	210	ASP	CB-CG-OD1	6.21	123.89	118.30
1	A	102	ASP	CB-CG-OD1	6.09	123.78	118.30
1	A	210	ASP	CB-CG-OD2	-6.05	112.86	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	ASP	CB-CG-OD1	6.00	123.70	118.30
1	A	129	ASP	CB-CG-OD2	-5.98	112.92	118.30
1	A	216	ASP	CB-CG-OD2	-5.82	113.06	118.30
1	A	103	ASP	CB-CG-OD2	-5.79	113.09	118.30
1	A	123	ILE	CB-CA-C	-5.71	100.18	111.60
1	A	180	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	A	19	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	A	76	ASP	CB-CG-OD2	-5.61	113.25	118.30
1	A	117	ASP	CB-CG-OD2	-5.59	113.27	118.30
1	A	155	ASP	CB-CG-OD2	-5.56	113.29	118.30
1	A	21	ASP	CB-CG-OD1	5.48	123.23	118.30
1	A	19	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	36	ASP	CB-CG-OD1	5.26	123.03	118.30
1	A	76	ASP	CB-CG-OD1	5.14	122.92	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1738	0	1615	108	0
2	A	30	0	0	2	0
All	All	1768	0	1615	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:THR:HG23	1:A:224:VAL:HG22	1.25	1.11
1:A:66:CRF:HZ3	1:A:150:VAL:HB	1.27	1.08
1:A:150:VAL:HG13	1:A:201:LEU:HB2	1.41	1.00
1:A:93:VAL:HG22	1:A:111:GLU:HG2	1.55	0.89

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:VAL:CG1	1:A:201:LEU:HB2	2.07	0.84
1:A:196:PRO:HB3	1:A:199:HIS:HE1	1.41	0.84
1:A:203:THR:HG23	1:A:224:VAL:CG2	2.05	0.84
1:A:221:LEU:HD12	1:A:222:GLU:N	1.94	0.83
1:A:171:ILE:HG13	1:A:176:VAL:HA	1.61	0.82
1:A:81:HIS:CE1	1:A:197:ASP:HB2	2.16	0.80
1:A:146:ILE:HB	1:A:148:HIS:CE1	2.18	0.78
1:A:18:LEU:HG	1:A:19:ASP:N	1.99	0.77
1:A:201:LEU:HD23	1:A:226:ALA:HA	1.66	0.76
1:A:221:LEU:HD12	1:A:222:GLU:H	1.52	0.74
1:A:194:LEU:C	1:A:195:LEU:HD23	2.08	0.73
1:A:135:ASN:HA	1:A:140:LYS:HD3	1.70	0.71
1:A:66:CRF:HZ3	1:A:150:VAL:CB	2.15	0.70
1:A:152:ILE:HD11	1:A:201:LEU:CD1	2.22	0.70
1:A:196:PRO:HB3	1:A:199:HIS:CE1	2.28	0.68
1:A:195:LEU:HD23	1:A:195:LEU:N	2.09	0.67
1:A:53:LEU:HD12	1:A:54:PRO:HD2	1.75	0.66
1:A:33:GLY:HA3	1:A:44:LEU:HD13	1.78	0.65
1:A:146:ILE:O	1:A:148:HIS:ND1	2.30	0.64
1:A:155:ASP:HB2	1:A:162:LYS:HG3	1.78	0.63
1:A:152:ILE:HD11	1:A:201:LEU:HD12	1.81	0.62
1:A:43:THR:O	1:A:44:LEU:HD22	1.98	0.62
1:A:190:ASP:HB2	2:A:321:HOH:O	2.00	0.61
1:A:96:ARG:HG3	1:A:183:GLN:HB2	1.83	0.61
1:A:167:ILE:O	1:A:178:LEU:HD22	2.00	0.61
1:A:27:PHE:HA	1:A:50:THR:OG1	2.01	0.60
1:A:166:LYS:O	1:A:167:ILE:HD13	2.04	0.58
1:A:201:LEU:HD23	1:A:226:ALA:CA	2.33	0.58
1:A:163:ALA:HB3	1:A:183:GLN:HG2	1.85	0.57
1:A:220:LEU:HD12	1:A:221:LEU:N	2.18	0.57
1:A:14:ILE:HD11	1:A:71:PHE:CE2	2.40	0.56
1:A:171:ILE:HD11	1:A:177:GLN:N	2.20	0.56
1:A:146:ILE:HB	1:A:148:HIS:HE1	1.69	0.56
1:A:43:THR:C	1:A:44:LEU:HD22	2.26	0.56
1:A:12:VAL:HG13	1:A:13:PRO:HD2	1.88	0.56
1:A:100:PHE:HB2	1:A:104:GLY:O	2.06	0.55
1:A:139:HIS:H	1:A:139:HIS:CD2	2.25	0.54
1:A:136:ILE:HD12	1:A:136:ILE:N	2.22	0.54
1:A:136:ILE:HD12	1:A:136:ILE:H	1.73	0.53
1:A:125:LEU:HD23	1:A:126:LYS:N	2.24	0.53
1:A:151:TYR:CD2	1:A:151:TYR:N	2.76	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:LEU:HD23	1:A:207:LEU:N	2.24	0.53
1:A:125:LEU:HD23	1:A:125:LEU:C	2.30	0.52
1:A:146:ILE:HD11	1:A:170:ASN:HD21	1.74	0.52
1:A:220:LEU:HD21	1:A:222:GLU:OE1	2.10	0.51
1:A:171:ILE:HG13	1:A:176:VAL:CA	2.36	0.51
1:A:1:MET:O	1:A:2:SER:O	2.28	0.51
1:A:152:ILE:HG22	1:A:161:ILE:HD12	1.94	0.50
1:A:47:ILE:HD13	1:A:217:HIS:HB3	1.94	0.49
1:A:150:VAL:O	1:A:200:TYR:HB2	2.12	0.49
1:A:135:ASN:HA	1:A:140:LYS:HB2	1.94	0.49
1:A:66:CRF:CZ3	1:A:150:VAL:HB	2.19	0.49
1:A:98:ILE:O	1:A:105:ASN:HB2	2.13	0.49
1:A:142:GLU:HG3	1:A:170:ASN:O	2.12	0.49
1:A:175:SER:O	1:A:176:VAL:HG23	2.13	0.49
1:A:97:THR:HG21	1:A:99:PHE:CZ	2.48	0.49
1:A:199:HIS:O	1:A:200:TYR:HB3	2.13	0.48
1:A:55:VAL:HG23	1:A:56:PRO:N	2.27	0.48
1:A:139:HIS:CD2	1:A:139:HIS:N	2.82	0.48
1:A:56:PRO:HD2	1:A:136:ILE:HG23	1.95	0.48
1:A:57:TRP:HB2	1:A:58:PRO:HD3	1.96	0.48
1:A:33:GLY:CA	1:A:44:LEU:HD13	2.44	0.47
1:A:210:ASP:O	1:A:213:GLU:N	2.42	0.47
1:A:171:ILE:HD11	1:A:176:VAL:C	2.34	0.47
1:A:147:SER:O	1:A:147:SER:OG	2.30	0.47
1:A:197:ASP:O	1:A:199:HIS:N	2.47	0.47
1:A:70:CYS:HA	1:A:84:PHE:HB2	1.97	0.47
1:A:87:ALA:HB1	1:A:91:GLY:O	2.15	0.46
1:A:18:LEU:HD23	1:A:18:LEU:O	2.16	0.46
1:A:123:ILE:CG2	1:A:124:GLU:N	2.79	0.46
1:A:123:ILE:HG22	1:A:124:GLU:N	2.29	0.46
1:A:195:LEU:HA	1:A:196:PRO:HD3	1.83	0.45
1:A:14:ILE:CD1	1:A:71:PHE:CE2	2.98	0.45
1:A:155:ASP:HB2	1:A:162:LYS:CG	2.46	0.45
1:A:137:LEU:HD23	1:A:137:LEU:HA	1.61	0.45
1:A:7:LEU:CD1	1:A:7:LEU:N	2.79	0.45
1:A:55:VAL:HB	1:A:56:PRO:HD2	1.99	0.44
1:A:97:THR:HG22	1:A:99:PHE:CE1	2.52	0.44
1:A:97:THR:CG2	1:A:99:PHE:CE1	3.00	0.44
1:A:15:LEU:HA	1:A:15:LEU:HD23	1.74	0.44
1:A:217:HIS:N	1:A:217:HIS:ND1	2.65	0.44
1:A:146:ILE:O	1:A:148:HIS:N	2.50	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:SER:OG	1:A:219:VAL:HB	2.17	0.44
1:A:96:ARG:HD2	2:A:306:HOH:O	2.17	0.43
1:A:178:LEU:HD22	1:A:178:LEU:HA	1.65	0.43
1:A:217:HIS:N	1:A:217:HIS:HD1	2.16	0.43
1:A:164:HIS:CE1	1:A:182:TYR:CZ	3.06	0.43
1:A:149:ASN:HA	1:A:201:LEU:O	2.18	0.43
1:A:27:PHE:CD1	1:A:54:PRO:HG2	2.54	0.43
1:A:98:ILE:HG12	1:A:181:HIS:CD2	2.54	0.43
1:A:170:ASN:HD22	1:A:176:VAL:CG2	2.32	0.43
1:A:220:LEU:HD12	1:A:221:LEU:H	1.82	0.42
1:A:146:ILE:CB	1:A:148:HIS:CE1	2.98	0.42
1:A:217:HIS:HD1	1:A:217:HIS:H	1.68	0.42
1:A:165:PHE:CE2	1:A:181:HIS:CG	3.08	0.41
1:A:215:ARG:HE	1:A:215:ARG:HB2	1.58	0.41
1:A:157:GLN:C	1:A:159:ASN:H	2.24	0.41
1:A:213:GLU:OE2	1:A:215:ARG:HB2	2.21	0.41
1:A:42:LEU:HD23	1:A:42:LEU:HA	1.91	0.40
1:A:81:HIS:HB3	1:A:196:PRO:HA	2.02	0.40
1:A:152:ILE:HD11	1:A:201:LEU:HG	2.03	0.40
1:A:201:LEU:CD2	1:A:226:ALA:HB2	2.51	0.40
1:A:26:ARG:O	1:A:27:PHE:HB3	2.22	0.40
1:A:115:GLU:HA	1:A:115:GLU:OE1	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	221/226 (98%)	196 (89%)	19 (9%)	6 (3%)	5   7

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	3	LYS
1	A	139	HIS
1	A	147	SER
1	A	191	GLY
1	A	198	ASN

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	169/196 (86%)	143 (85%)	26 (15%)	2   5

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	7	LEU
1	A	9	THR
1	A	15	LEU
1	A	18	LEU
1	A	78	MET
1	A	109	ARG
1	A	126	LYS
1	A	139	HIS
1	A	140	LYS
1	A	145	TYR
1	A	150	VAL
1	A	152	ILE
1	A	178	LEU
1	A	183	GLN
1	A	195	LEU
1	A	197	ASP
1	A	198	ASN
1	A	200	TYR
1	A	202	SER
1	A	204	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	205	SER
1	A	207	LEU
1	A	209	LYS
1	A	215	ARG
1	A	225	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	HIS
1	A	149	ASN
1	A	164	HIS
1	A	170	ASN
1	A	204	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CRF	A	66	1	25,26,27	1.49	4 (16%)	32,37,39	2.88	9 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRF	A	66	1	-	2/10/31/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRF	CG2-CB2	-4.45	1.37	1.46
1	A	66	CRF	CA3-N3	-3.09	1.41	1.47
1	A	66	CRF	C1-N3	2.55	1.41	1.37
1	A	66	CRF	CZ2-CE2	-2.25	1.38	1.41

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CRF	O2-C2-CA2	-9.18	125.80	130.96
1	A	66	CRF	CA2-C2-N3	7.69	107.00	103.37
1	A	66	CRF	C2-N3-C1	-6.32	104.77	107.97
1	A	66	CRF	CG2-CB2-CA2	-4.82	121.47	130.81
1	A	66	CRF	N3-C1-N2	4.08	114.28	111.45
1	A	66	CRF	CA1-C1-N3	-3.46	120.59	124.75
1	A	66	CRF	C1-CA1-N1	-2.35	106.14	109.96
1	A	66	CRF	C2-CA2-N2	-2.25	107.36	108.93
1	A	66	CRF	CH2-CZ3-CE3	-2.09	117.50	120.44

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66	CRF	N2-CA2-CB2-CG2
1	A	66	CRF	C2-CA2-CB2-CG2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	CRF	3	0

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	225/226 (99%)	0.23	15 (6%) 17 18	26, 46, 86, 95	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	159	ASN	5.1
1	A	152	ILE	4.2
1	A	150	VAL	3.6
1	A	228	GLY	3.3
1	A	158	LYS	2.9
1	A	197	ASP	2.9
1	A	148	HIS	2.9
1	A	211	PRO	2.8
1	A	157	GLN	2.7
1	A	161	ILE	2.7
1	A	1	MET	2.7
1	A	26	ARG	2.3
1	A	60	LEU	2.2
1	A	160	GLY	2.1
1	A	153	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CRF	A	66	24/25	0.93	0.21	13,24,38,44	10

### 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

### 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.