



# wwPDB X-ray Structure Validation Summary Report

Oct 31, 2021 – 06:34 PM EDT

PDB ID : 3DD7  
Title : Structure of DocH66Y in complex with the C-terminal domain of Phd  
Authors : Garcia-Pino, A.; Loris, R.  
Deposited on : 2008-06-05  
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	
1	C	135	
2	B	23	
2	D	23	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BR	A	140	-	-	X	-
3	BR	A	143	-	-	X	-
3	BR	C	136	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2451 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Death on curing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	123	937	598	156	180	3	1	7	0
1	C	122	960	614	166	178	2	1	11	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	TYR	HIS	engineered mutation	UNP Q06259
A	126	ASP	GLU	engineered mutation	UNP Q06259
A	127	PRO	-	expression tag	UNP Q06259
A	128	LEU	-	expression tag	UNP Q06259
A	129	GLU	-	expression tag	UNP Q06259
A	130	HIS	-	expression tag	UNP Q06259
A	131	HIS	-	expression tag	UNP Q06259
A	132	HIS	-	expression tag	UNP Q06259
A	133	HIS	-	expression tag	UNP Q06259
A	134	HIS	-	expression tag	UNP Q06259
A	135	HIS	-	expression tag	UNP Q06259
C	66	TYR	HIS	engineered mutation	UNP Q06259
C	126	ASP	GLU	engineered mutation	UNP Q06259
C	127	PRO	-	expression tag	UNP Q06259
C	128	LEU	-	expression tag	UNP Q06259
C	129	GLU	-	expression tag	UNP Q06259
C	130	HIS	-	expression tag	UNP Q06259
C	131	HIS	-	expression tag	UNP Q06259
C	132	HIS	-	expression tag	UNP Q06259
C	133	HIS	-	expression tag	UNP Q06259
C	134	HIS	-	expression tag	UNP Q06259
C	135	HIS	-	expression tag	UNP Q06259

- Molecule 2 is a protein called Prevent host death protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
2	B	18	142	90	20	30	2	0	2	0
2	D	22	168	103	25	39	1	0	2	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	70	MSE	LEU	engineered mutation	UNP Q06253
D	70	MSE	LEU	engineered mutation	UNP Q06253

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	9	Total 9	Br 9	0	0
3	B	1	Total 1	Br 1	0	0
3	C	6	Total 6	Br 6	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	83	Total 83	O 83	0	0
4	B	17	Total 17	O 17	0	0
4	C	103	Total 103	O 103	0	0
4	D	25	Total 25	O 25	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.94Å 38.20Å 63.73Å 90.00° 99.26° 90.00°	Depositor
Resolution (Å)	11.00 – 1.70 11.91 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.5 (11.00-1.70) 96.5 (11.91-1.70)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.182 , 0.198 0.184 , 0.199	Depositor DCC
$R_{free}$ test set	1444 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.9	Xtrriage
Anisotropy	0.054	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2451	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.52	0/972	0.61	1/1319 (0.1%)
1	C	0.63	2/1009 (0.2%)	0.96	4/1365 (0.3%)
2	B	0.53	0/148	0.66	0/195
2	D	0.57	0/174	0.66	0/232
All	All	0.58	2/2303 (0.1%)	0.79	5/3111 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	112[A]	SER	CB-OG	-7.96	1.31	1.42
1	C	112[B]	SER	CB-OG	-7.96	1.31	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	42[A]	ARG	NE-CZ-NH2	-14.57	113.01	120.30
1	C	42[B]	ARG	NE-CZ-NH2	-14.57	113.01	120.30
1	C	42[A]	ARG	NE-CZ-NH1	13.64	127.12	120.30
1	C	42[B]	ARG	NE-CZ-NH1	13.64	127.12	120.30
1	A	64	ARG	NE-CZ-NH2	-5.64	117.48	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	937	0	943	11	0
1	C	960	0	984	10	0
2	B	142	0	135	1	0
2	D	168	0	147	4	0
3	A	9	0	0	7	0
3	B	1	0	0	1	0
3	C	6	0	0	4	0
4	A	83	0	0	5	2
4	B	17	0	0	0	0
4	C	103	0	0	3	2
4	D	25	0	0	2	1
All	All	2451	0	2209	28	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:203:HOH:O	3:B:9:BR:BR	2.12	1.16
3:A:143:BR:BR	4:A:217:HOH:O	2.24	1.08
3:C:140:BR:BR	4:C:282:HOH:O	2.26	1.08
1:A:26[A]:MET:SD	3:A:140:BR:BR	2.92	0.82
3:A:141:BR:BR	4:A:203:HOH:O	2.53	0.80

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:247:HOH:O	4:C:288:HOH:O[1_565]	1.98	0.22
4:A:219:HOH:O	4:C:273:HOH:O[3_545]	2.03	0.17
4:A:223:HOH:O	4:D:185:HOH:O[3_545]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/135 (95%)	128 (100%)	0	0	100	100
1	C	129/135 (96%)	129 (100%)	0	0	100	100
2	B	18/23 (78%)	18 (100%)	0	0	100	100
2	D	22/23 (96%)	22 (100%)	0	0	100	100
All	All	297/316 (94%)	297 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/108 (89%)	95 (99%)	1 (1%)	76	67
1	C	99/108 (92%)	98 (99%)	1 (1%)	76	67
2	B	17/19 (90%)	17 (100%)	0	100	100
2	D	19/19 (100%)	19 (100%)	0	100	100
All	All	231/254 (91%)	229 (99%)	2 (1%)	78	70

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	GLU
1	C	23	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	ASN
1	C	69	ASN
2	D	67	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	123/135 (91%)	0.18	2 (1%) 72 76	16, 21, 26, 33	0
1	C	122/135 (90%)	0.16	1 (0%) 86 88	18, 21, 27, 30	0
2	B	17/23 (73%)	0.30	2 (11%) 4 5	14, 17, 27, 28	0
2	D	21/23 (91%)	0.85	3 (14%) 2 2	15, 17, 25, 38	0
All	All	283/316 (89%)	0.23	8 (2%) 53 57	14, 21, 27, 38	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	72	ASN	8.5
2	D	51	ALA	5.7
1	A	66	TYR	4.8
2	D	71	VAL	4.1
1	A	28	ASP	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BR	A	141	1/1	0.91	0.10	43,43,43,43	1
3	BR	A	142	1/1	0.95	0.07	20,20,20,20	1
3	BR	A	143	1/1	0.95	0.10	18,18,18,18	1
3	BR	B	9	1/1	0.95	0.08	26,26,26,26	1
3	BR	A	144	1/1	0.97	0.05	23,23,23,23	1
3	BR	A	139	1/1	0.97	0.12	30,30,30,30	1
3	BR	C	138	1/1	0.97	0.10	29,29,29,29	1
3	BR	C	140	1/1	0.97	0.06	22,22,22,22	1
3	BR	A	140	1/1	0.98	0.09	25,25,25,25	1
3	BR	A	137	1/1	0.98	0.15	24,24,24,24	1
3	BR	C	136	1/1	0.99	0.05	18,18,18,18	1
3	BR	C	137	1/1	0.99	0.07	21,21,21,21	1
3	BR	A	138	1/1	0.99	0.12	23,23,23,23	1
3	BR	C	139	1/1	0.99	0.06	17,17,17,17	1
3	BR	A	136	1/1	0.99	0.10	18,18,18,18	1
3	BR	C	141	1/1	0.99	0.14	26,26,26,26	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.