



# Full wwPDB NMR Structure Validation Report ⓘ

Dec 5, 2022 – 08:02 AM EST

PDB ID : 8DGH  
BMRB ID : 51447  
Title : NMR Structure of calmodulin bound to C-terminal site in the beta-subunit of cyclic nucleotide-gated channel  
Authors : Bej, A.; Ames, J.B.  
Deposited on : 2022-06-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.31.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 68%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	149	
2	B	12	

## 2 Ensemble composition and analysis i

This entry contains 10 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:84-A:149, B:1128-B:1136 (75)	0.30	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 7, 9
2	4, 8
Single-model clusters	2; 3; 5; 6; 10

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1271 atoms, of which 625 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Calmodulin-1.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	68	1051	333	506	90	118	4	0

- Molecule 2 is a protein called Cyclic nucleotide-gated cation channel beta-1.


Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
2	B	12	220	65	119	22	14	0

## 4 Residue-property plots


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Calmodulin-1



- Molecule 2: Cyclic nucleotide-gated cation channel beta-1




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Calmodulin-1




- Molecule 2: Cyclic nucleotide-gated cation channel beta-1




#### 4.2.2 Score per residue for model 2

- Molecule 1: Calmodulin-1




- Molecule 2: Cyclic nucleotide-gated cation channel beta-1




#### 4.2.3 Score per residue for model 3

- Molecule 1: Calmodulin-1



- Molecule 2: Cyclic nucleotide-gated cation channel beta-1



#### 4.2.4 Score per residue for model 4

- Molecule 1: Calmodulin-1

Chain A:  42% .. 54%

MET ALA ASP GLN LEU THR GLU GLN GLN ILE ALA LEU THR MET PHE LYS GLU ALA LYS PHE SER LYS LEU PHE ASP LYS ASP GLY ASP GLY THR ILE THR THR LYS GLU LEU THR VAL MET ARG SER LEU GLN ASN PRO THR GLU ALA GLU LEU GLN ASP MET ILE ASN VAL ASP ALA ASP GLY

ASN GLY THR ILE ASP PHE PRO GLU PHE LEU THR MET MET ALA ARG LYS MET LYS ASP THR ASP E82 E83 D119 M138 T147 A148 K149

- Molecule 2: Cyclic nucleotide-gated cation channel beta-1

Chain B:  58% 17% 25%

K1128 L1132 R1133 K1137 E1138 L1139

#### 4.2.5 Score per residue for model 5

- Molecule 1: Calmodulin-1

Chain A:  42% .. 54%

MET ALA ASP GLN LEU THR GLU GLN GLN ILE ALA LEU THR MET PHE LYS GLU ALA LYS PHE SER LYS LEU PHE ASP LYS ASP GLY ASP GLY THR ILE THR THR LYS GLU LEU THR VAL MET ARG SER LEU GLN ASN PRO THR GLU ALA GLU LEU GLN ASP MET ILE ASN VAL ASP ALA ASP GLY

ASN GLY THR ILE ASP PHE PRO GLU PHE LEU THR MET MET ALA ARG LYS MET LYS ASP THR ASP E82 E83 L106 M110 M138 K149

- Molecule 2: Cyclic nucleotide-gated cation channel beta-1

Chain B:  67% 8% 25%

K1128 R1133 K1137 E1138 L1139

#### 4.2.6 Score per residue for model 6


- Molecule 1: Calmodulin-1

Chain A:  40% .. 54%

MET ALA ASP GLN LEU THR GLU GLN GLN ILE ALA LEU THR MET PHE LYS GLU ALA LYS PHE SER LYS LEU PHE ASP LYS ASP GLY ASP GLY THR ILE THR THR LYS GLU LEU THR VAL MET ARG SER LEU GLN ASN PRO THR GLU ALA GLU LEU GLN ASP MET ILE ASN VAL ASP ALA ASP GLY


ASN GLY THR ILE ASP PHE PRO GLU PHE LEU THR MET MET ALA ARG LYS MET LYS ASP THR ASP E82 E83 L106 M110 E121 M125 V137 M138 K149

- Molecule 2: Cyclic nucleotide-gated cation channel beta-1



#### 4.2.7 Score per residue for model 7

- Molecule 1: Calmodulin-1




- Molecule 2: Cyclic nucleotide-gated cation channel beta-1




#### 4.2.8 Score per residue for model 8

- Molecule 1: Calmodulin-1





- Molecule 2: Cyclic nucleotide-gated cation channel beta-1




#### 4.2.9 Score per residue for model 9

- Molecule 1: Calmodulin-1






- Molecule 2: Cyclic nucleotide-gated cation channel beta-1




### 4.2.10 Score per residue for model 10

- Molecule 1: Calmodulin-1



- Molecule 2: Cyclic nucleotide-gated cation channel beta-1



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 10 were deposited, based on the following criterion: *structures with acceptable covalent geometry*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
HADDOCK	refinement	
HADDOCK	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1450
Number of shifts mapped to atoms	1450
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	68%

## 6 Model quality i

### 6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	530	495	495	1±1
2	B	75	89	88	0±0
All	All	6050	5840	5830	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:94:ASP:OD1	1:A:97:GLY:HA2	0.47	2.09	7	1
1:A:142:PHE:O	1:A:145:MET:HG2	0.47	2.10	9	4
1:A:123:ASP:O	1:A:127:ARG:HG3	0.45	2.11	1	1
1:A:106:LEU:O	1:A:110:MET:HG2	0.45	2.12	6	2
1:A:107:ARG:HD3	1:A:122:VAL:HG21	0.42	1.91	10	1
1:A:146:MET:HA	2:B:1133:ARG:NH2	0.42	2.29	10	1
1:A:85:GLU:HB3	2:B:1133:ARG:NH1	0.42	2.29	1	1
1:A:121:GLU:O	1:A:125:MET:HG3	0.41	2.16	6	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	65/149 (44%)	64±1 (98±1%)	1±1 (2±1%)	0±0 (0±0%)	100	100
2	B	8/12 (67%)	8±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
All	All	730/1610 (45%)	716 (98%)	14 (2%)	0 (0%)	100	100

There are no Ramachandran outliers.

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	56/127 (44%)	54±1 (96±2%)	2±1 (4±2%)	33	81
2	B	7/10 (70%)	6±1 (84±15%)	1±1 (16±15%)	5	42
All	All	630/1370 (46%)	595 (94%)	35 (6%)	25	74

All 16 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	138	ASN	7
2	B	1132	LEU	4
1	A	123	ASP	4
2	B	1133	ARG	4
1	A	119	ASP	3
1	A	132	ASP	2
1	A	147	THR	2
2	B	1129	LEU	1
1	A	102	SER	1
1	A	84	GLU	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	137	VAL	1
1	A	96	ASP	1
1	A	130	ASP	1
1	A	144	GLN	1
2	B	1131	HIS	1
2	B	1136	LEU	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 68% for the well-defined parts and 65% for the entire structure.

### 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: *assigned\_chemical\_shifts\_1*

#### 7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1450
Number of shifts mapped to atoms	1450
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	143	$-0.41 \pm 0.09$	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	132	$0.12 \pm 0.16$	None needed (< 0.5 ppm)
$^{13}\text{C}'$	123	$-0.45 \pm 0.13$	None needed (< 0.5 ppm)
$^{15}\text{N}$	138	$0.03 \pm 0.56$	None needed (< 0.5 ppm)

#### 7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 68%, i.e. 640 atoms were assigned a chemical shift out of a possible 940. 8 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	325/375 (87%)	132/150 (88%)	127/150 (85%)	66/75 (88%)
Sidechain	297/508 (58%)	181/294 (62%)	116/187 (62%)	0/27 (0%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	18/57 (32%)	18/31 (58%)	0/24 (0%)	0/2 (0%)
Overall	640/940 (68%)	331/475 (70%)	243/361 (67%)	66/104 (63%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 65%, i.e. 652 atoms were assigned a chemical shift out of a possible 1004. 8 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	333/400 (83%)	135/160 (84%)	130/160 (81%)	68/80 (85%)
Sidechain	301/547 (55%)	183/317 (58%)	118/202 (58%)	0/28 (0%)
Aromatic	18/57 (32%)	18/31 (58%)	0/24 (0%)	0/2 (0%)
Overall	652/1004 (65%)	336/508 (66%)	248/386 (64%)	68/110 (62%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

