



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 7, 2023 – 01:01 PM EST

PDB ID : 8DGQ
Title : Crystal structure of p120RasGAP SH2-SH3-SH2 in complex with p190RhoGAP doubly phosphorylated peptide
Authors : Stiegler, A.L.; Vish, K.J.; Boggon, T.J.
Deposited on : 2022-06-24
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	
1	B	273	
2	U	29	
2	V	29	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5079 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ras GTPase-activating protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	272	Total	C	N	O	S	0	1	0
			2223	1413	384	417	9			
1	B	271	Total	C	N	O	S	0	2	0
			2230	1417	387	417	9			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	172	GLY	-	expression tag	UNP P20936
A	173	SER	-	expression tag	UNP P20936
A	236	SER	CYS	conflict	UNP P20936
A	261	SER	CYS	conflict	UNP P20936
A	372	SER	CYS	conflict	UNP P20936
A	402	SER	CYS	conflict	UNP P20936
B	172	GLY	-	expression tag	UNP P20936
B	173	SER	-	expression tag	UNP P20936
B	236	SER	CYS	conflict	UNP P20936
B	261	SER	CYS	conflict	UNP P20936
B	372	SER	CYS	conflict	UNP P20936
B	402	SER	CYS	conflict	UNP P20936

- Molecule 2 is a protein called Rho GTPase-activating protein 35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
2	U	25	Total	C	N	O	P	S	0	0	0
			210	125	33	49	2	1			
2	V	20	Total	C	N	O	P	S	0	0	0
			162	96	23	40	2	1			

- Molecule 3 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 3 4	0	0
3	A	1	Total C O 7 3 4	0	0
3	B	1	Total C O 7 3 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	101	Total O 101 101	0	0
4	B	110	Total O 110 110	0	0
4	U	11	Total O 11 11	0	0
4	V	11	Total O 11 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ras GTPase-activating protein 1




- Molecule 1: Ras GTPase-activating protein 1



- Molecule 2: Rho GTPase-activating protein 35



- Molecule 2: Rho GTPase-activating protein 35



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.21Å 113.50Å 119.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.65 – 1.95 49.65 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.65-1.95) 100.0 (49.65-1.95)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 1.95Å)	Xtrriage
Refinement program	PHENIX 1.19.2	Depositor
R, R_{free}	0.213 , 0.256 0.212 , 0.256	Depositor DCC
R_{free} test set	2010 reflections (3.63%)	wwPDB-VP
Wilson B-factor (Å ²)	38.3	Xtrriage
Anisotropy	0.336	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.000 for -h,l,k	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5079	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.59 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4941e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/2280	0.59	0/3085
1	B	0.34	0/2287	0.59	0/3094
2	U	0.25	0/179	0.49	0/240
2	V	0.32	0/130	0.48	0/173
All	All	0.33	0/4876	0.58	0/6592

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2223	0	2163	9	0
1	B	2230	0	2172	28	0
2	U	210	0	178	1	0
2	V	162	0	128	2	0
3	A	14	0	4	0	0
3	B	7	0	2	0	0
4	A	101	0	0	0	0
4	B	110	0	0	0	0
4	U	11	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	V	11	0	0	0	0
All	All	5079	0	4647	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:259:VAL:HG23	1:B:267:LYS:HE3	1.65	0.77
1:B:197:GLN:HG2	1:B:405:PRO:HB2	1.81	0.62
1:B:260:SER:HA	1:B:263:LEU:O	2.00	0.62
1:B:398:ARG:NH2	2:V:1087:PTR:O1P	2.34	0.61
1:B:364:LEU:HG	1:B:441:VAL:HG21	1.81	0.61
1:B:258:HIS:HA	1:B:267:LYS:HE2	1.86	0.58
1:A:197:GLN:HG2	1:A:405:PRO:HB2	1.85	0.58
1:B:353:HIS:CE1	1:B:443:MET:HG2	2.39	0.57
1:A:315:ASP:HB2	1:A:317:TRP:HD1	1.69	0.57
1:B:315:ASP:HB2	1:B:317:TRP:HD1	1.69	0.56
1:B:260:SER:HB2	1:B:265:GLY:H	1.71	0.56
1:B:277:PRO:HB2	1:B:345:PRO:HG2	1.91	0.52
1:B:256:TYR:CD1	1:B:261:SER:HB3	2.44	0.52
1:B:259:VAL:CG2	1:B:267:LYS:HE3	2.37	0.51
1:B:315:ASP:N	1:B:315:ASP:OD1	2.43	0.51
1:A:389:TYR:CZ	1:A:398:ARG:HD2	2.47	0.50
1:B:256:TYR:HD1	1:B:261:SER:HB3	1.77	0.48
1:B:315:ASP:HB2	1:B:317:TRP:CD1	2.50	0.46
1:B:218:LEU:HD21	1:B:262:LEU:HD11	1.98	0.46
1:A:235:MET:HB3	1:A:240:TYR:CE1	2.51	0.45
1:A:211:ARG:HD2	2:U:1100:GLU:OE1	2.17	0.44
1:B:188:ARG:CZ	2:V:1103:ASN:HB3	2.48	0.44
1:B:372:SER:HA	1:B:390:PHE:CE1	2.52	0.44
1:B:287:ILE:HG12	1:B:335:LEU:O	2.17	0.44
1:B:428:LYS:HG3	1:B:429:GLU:HG2	2.00	0.42
1:B:353:HIS:HE1	1:B:443:MET:HG2	1.82	0.42
1:B:416:TYR:CD1	1:B:422:ILE:HG12	2.55	0.42
1:A:318:MET:HE1	1:A:336:VAL:HG21	2.02	0.42
1:A:197:GLN:HG2	1:A:405:PRO:CB	2.50	0.42
1:A:240:TYR:HA	1:A:244:ARG:O	2.19	0.42
1:B:389:TYR:CZ	1:B:398:ARG:HD2	2.54	0.42
1:B:318:MET:SD	1:B:333:GLU:HG3	2.60	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:ARG:HG2	1:B:308:ILE:HD13	2.03	0.41
1:B:240:TYR:HA	1:B:244:ARG:O	2.21	0.40
1:A:315:ASP:HB2	1:A:317:TRP:CD1	2.55	0.40
1:B:374:PHE:HA	1:B:389:TYR:O	2.21	0.40
1:B:362:TYR:OH	1:B:398:ARG:HD3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/273 (99%)	266 (98%)	5 (2%)	0	100	100
1	B	271/273 (99%)	266 (98%)	5 (2%)	0	100	100
2	U	21/29 (72%)	21 (100%)	0	0	100	100
2	V	14/29 (48%)	14 (100%)	0	0	100	100
All	All	577/604 (96%)	567 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/244 (100%)	241 (99%)	3 (1%)	71	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	245/244 (100%)	243 (99%)	2 (1%)	81	80
2	U	21/25 (84%)	21 (100%)	0	100	100
2	V	16/25 (64%)	16 (100%)	0	100	100
All	All	526/538 (98%)	521 (99%)	5 (1%)	76	74

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	235	MET
1	A	236	SER
1	A	315	ASP
1	B	315	ASP
1	B	334	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PTR	U	1105	2	15,16,17	1.31	1 (6%)	19,22,24	0.47	0
2	PTR	V	1105	2	15,16,17	1.27	1 (6%)	19,22,24	0.64	0
2	PTR	U	1087	2	15,16,17	1.40	1 (6%)	19,22,24	0.59	0
2	PTR	V	1087	2	15,16,17	1.40	1 (6%)	19,22,24	0.78	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	U	1105	2	-	1/10/11/13	0/1/1/1
2	PTR	V	1105	2	-	0/10/11/13	0/1/1/1
2	PTR	U	1087	2	-	0/10/11/13	0/1/1/1
2	PTR	V	1087	2	-	1/10/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	V	1087	PTR	OH-CZ	-4.86	1.29	1.40
2	U	1087	PTR	OH-CZ	-4.78	1.29	1.40
2	V	1105	PTR	OH-CZ	-4.55	1.30	1.40
2	U	1105	PTR	OH-CZ	-4.42	1.30	1.40

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	V	1087	PTR	CB-CA-C	-2.27	107.21	111.47

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	V	1087	PTR	O-C-CA-CB
2	U	1105	PTR	CZ-OH-P-O2P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	V	1087	PTR	1	0

5.5 Carbohydrates

There are no monosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MLI	A	502	-	6,6,6	1.50	0	7,7,7	1.05	0
3	MLI	B	501	-	6,6,6	1.53	0	7,7,7	1.27	0
3	MLI	A	501	-	6,6,6	1.33	0	7,7,7	1.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	A	502	-	-	2/4/4/4	-
3	MLI	B	501	-	-	0/4/4/4	-
3	MLI	A	501	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	MLI	C3-C1-C2-O6
3	A	502	MLI	C2-C1-C3-O9
3	A	501	MLI	C2-C1-C3-O8
3	A	501	MLI	C3-C1-C2-O7
3	A	501	MLI	C2-C1-C3-O9
3	A	502	MLI	C2-C1-C3-O8

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/273 (99%)	0.99	38 (13%) 2 4	30, 45, 76, 103	0
1	B	271/273 (99%)	1.14	41 (15%) 2 3	29, 46, 85, 108	0
2	U	23/29 (79%)	1.67	7 (30%) 0 0	40, 58, 117, 140	0
2	V	18/29 (62%)	1.42	5 (27%) 0 0	42, 52, 80, 81	0
All	All	584/604 (96%)	1.10	91 (15%) 2 3	29, 47, 84, 140	0

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	U	1098	ARG	7.1
1	B	260	SER	6.8
1	B	174	THR	6.3
1	B	265	GLY	5.7
1	B	173	SER	5.4
1	A	341	ARG	5.1
1	B	224	MET	5.0
1	B	277	PRO	5.0
1	A	224	MET	4.5
2	V	1111	SER	4.5
2	U	1100	GLU	4.3
1	B	259	VAL	4.2
1	B	270	TYR	4.2
1	B	276	GLU	4.0
1	A	276	GLU	3.8
2	U	1101	GLU	3.8
1	B	269	LEU	3.8
1	A	172	GLY	3.7
1	B	443	MET	3.6
1	A	295	ASP	3.5
1	B	175	ALA	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	V	1110	ASP	3.4
1	B	341	ARG	3.4
1	B	410[A]	MET	3.4
1	A	443	MET	3.4
2	U	1099	ASN	3.3
1	B	222	SER	3.2
1	A	211	ARG	3.1
1	A	265	GLY	3.1
1	A	277	PRO	3.0
2	V	1094	VAL	3.0
1	A	422	ILE	2.9
1	A	420	GLY	2.9
1	B	442	PRO	2.9
1	B	177	PRO	2.8
1	B	211	ARG	2.8
1	A	410[A]	MET	2.8
1	B	262	LEU	2.8
1	A	405	PRO	2.8
1	A	416	TYR	2.8
1	B	267	LYS	2.7
1	A	196	ARG	2.7
1	A	243	GLY	2.7
1	B	405	PRO	2.7
1	B	225	ASN	2.7
1	A	258	HIS	2.7
1	A	247	SER	2.7
1	B	315	ASP	2.7
1	B	263	LEU	2.7
1	A	280	ASP	2.6
1	B	223	GLN	2.6
1	A	394	GLU	2.6
1	A	323	LEU	2.5
1	B	419	ILE	2.5
1	B	198	ALA	2.5
1	B	401	ILE	2.4
1	B	394	GLU	2.4
1	A	242	GLY	2.4
1	B	264	LYS	2.4
1	B	278	VAL	2.4
2	U	1109	HIS	2.3
1	A	278	VAL	2.3
1	A	173	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	225	ASN	2.3
1	A	269	LEU	2.3
1	A	230	PHE	2.3
1	B	257	SER	2.3
1	A	406	ASN	2.3
1	A	442	PRO	2.3
1	A	200	LYS	2.3
1	B	178	THR	2.3
1	B	255	TYR	2.2
1	A	245	ARG	2.2
1	B	418	SER	2.2
1	A	401	ILE	2.2
1	A	419	ILE	2.2
2	V	1109	HIS	2.2
1	A	340	GLY	2.2
1	A	176	PRO	2.2
1	B	438	LYS	2.1
1	B	342	GLU	2.1
1	A	270	TYR	2.1
1	A	201	SER	2.1
2	U	1086	ASP	2.1
2	U	1102	GLU	2.1
2	V	1085	SER	2.0
1	A	226	VAL	2.0
1	B	252	LEU	2.0
1	B	179	ASN	2.0
1	B	441	VAL	2.0
1	B	280	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PTR	U	1087	16/17	0.94	0.14	46,52,62,65	0
2	PTR	U	1105	16/17	0.94	0.14	45,53,60,64	0
2	PTR	V	1087	16/17	0.96	0.15	45,47,54,57	0
2	PTR	V	1105	16/17	0.97	0.12	36,42,49,56	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MLI	A	502	7/7	0.71	0.18	55,57,68,72	0
3	MLI	A	501	7/7	0.86	0.18	39,48,57,58	0
3	MLI	B	501	7/7	0.86	0.15	46,51,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.