



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2021 – 09:54 PM EST

PDB ID : 1DII
Title : CRYSTAL STRUCTURE OF P-CRESOL METHYLHYDROXYLASE AT 2.5
A RESOLUTION
Authors : Cunane, L.M.; Chen, Z.W.; Shamala, N.; Mathews, F.S.; Cronin, C.N.; McIn-
tire, W.S.
Deposited on : 1999-11-29
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.17.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.17.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	 2% 88% 10% ..
1	B	521	 2% 88% 11% ..
2	C	80	 85% 5% • 9%
2	D	80	 2% 84% 8% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FAD	A	599	X	-	-	-
4	FAD	B	599	X	-	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called P-CRESOL METHYLHYDROXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	515	Total 4025	C 2562	N 688	O 748	S 27	0	0	0
1	B	515	Total 4083	C 2599	N 701	O 755	S 28	0	6	0

- Molecule 2 is a protein called P-CRESOL METHYLHYDROXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	73	Total 559	C 355	N 95	O 106	S 3	0	0	0
2	D	73	Total 559	C 355	N 95	O 106	S 3	0	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	B	1	Total Cl 1 1	0	0

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
4	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	155	Total 155	O 155	0	0
6	C	21	Total 21	O 21	0	0
6	B	188	Total 188	O 188	0	0
6	D	21	Total 21	O 21	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: P-CRESOL METHYLHYDROXYLASE




- Molecule 1: P-CRESOL METHYLHYDROXYLASE



- Molecule 2: P-CRESOL METHYLHYDROXYLASE



- Molecule 2: P-CRESOL METHYLHYDROXYLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	140.30Å 130.60Å 74.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.50 6.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	88.4 (30.00-2.50) 96.2 (6.00-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 2.49Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.172 , 0.230 0.167 , 0.221	Depositor DCC
R_{free} test set	4300 reflections (10.12%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtrriage
Anisotropy	0.263	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 72.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9805	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/4123	0.80	2/5593 (0.0%)
1	B	0.70	0/4184	0.79	1/5675 (0.0%)
2	C	0.80	0/574	0.86	1/779 (0.1%)
2	D	0.76	0/574	0.85	1/779 (0.1%)
All	All	0.72	0/9455	0.80	5/12826 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	323	LEU	CA-CB-CG	8.88	135.72	115.30
1	B	323	LEU	CA-CB-CG	8.72	135.36	115.30
2	D	619	HIS	ND1-CG-CD2	7.93	119.90	108.80
2	C	619	HIS	ND1-CG-CD2	7.79	119.70	108.80
1	A	274	ILE	N-CA-C	-5.01	97.47	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	434	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4025	0	3959	19	0
1	B	4083	0	4011	25	0
2	C	559	0	538	2	0
2	D	559	0	538	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	53	0	29	2	0
4	B	53	0	30	3	0
5	C	43	0	30	0	0
5	D	43	0	30	0	0
6	A	155	0	0	4	0
6	B	188	0	0	5	0
6	C	21	0	0	0	0
6	D	21	0	0	0	0
All	All	9805	0	9165	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:473[B]:TYR:HE2	6:B:802:HOH:O	1.15	1.26
1:B:473[B]:TYR:CE2	6:B:802:HOH:O	1.89	1.16
1:A:336:VAL:HG21	1:B:336:VAL:HG21	1.68	0.75
1:A:492:ARG:O	1:A:496:HIS:HD2	1.84	0.61
1:B:492:ARG:O	1:B:496:HIS:HD2	1.86	0.59
1:B:114:ASN:H	1:B:114:ASN:HD22	1.50	0.58
1:A:114:ASN:HD22	1:A:114:ASN:H	1.51	0.57
1:A:396:ALA:HB3	6:A:837:HOH:O	2.05	0.57
1:B:92:ASN:HA	4:B:599:FAD:H5'2	1.89	0.54
1:A:86:THR:HG21	1:A:230:ILE:HG23	1.91	0.51
1:B:496:HIS:CE1	1:B:516:ASP:H	2.31	0.48
1:A:491:LYS:HE3	6:B:816:HOH:O	2.13	0.48
1:A:155:PRO:HG2	4:A:599:FAD:H2'	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:496:HIS:CE1	1:A:516:ASP:H	2.32	0.47
1:A:428:PHE:HA	1:A:436:HIS:O	2.15	0.47
1:B:86:THR:HG21	1:B:230:ILE:HG23	1.97	0.47
1:B:487:TYR:HB3	1:B:491:LYS:HD3	1.95	0.47
1:A:92:ASN:HA	4:A:599:FAD:H5'2	1.97	0.46
1:A:487:TYR:HB3	1:A:491:LYS:HD3	1.96	0.46
1:A:368:ARG:NH2	6:A:703:HOH:O	2.49	0.46
1:B:308:VAL:O	1:B:312:MET:HG3	2.16	0.45
1:A:417:LEU:HD23	1:A:457:CYS:SG	2.56	0.45
1:B:155:PRO:HG2	4:B:599:FAD:H2'	1.99	0.44
1:B:368:ARG:NH2	6:B:710:HOH:O	2.48	0.44
1:A:18:PHE:O	1:A:22:VAL:HG23	2.18	0.44
1:B:67:VAL:HG23	6:B:868:HOH:O	2.18	0.44
1:A:308:VAL:O	1:A:312:MET:HG3	2.17	0.43
1:B:434:ASP:OD2	1:B:436[A]:HIS:NE2	2.50	0.43
1:B:18:PHE:O	1:B:22:VAL:HG23	2.19	0.43
1:B:186:GLU:HB3	1:B:233:LYS:HB2	2.01	0.43
1:B:261:ILE:O	1:B:265:LEU:HD23	2.19	0.43
1:B:428:PHE:HA	1:B:436[A]:HIS:O	2.18	0.42
1:A:261:ILE:O	1:A:265:LEU:HD23	2.19	0.42
1:B:496:HIS:HE1	1:B:516:ASP:H	1.66	0.42
1:B:401:ALA:HA	1:B:435[A]:MET:HB2	2.01	0.42
2:D:628:VAL:O	2:D:632:ARG:HD3	2.19	0.42
1:A:278:VAL:HB	1:A:430:VAL:HG22	2.02	0.41
6:A:746:HOH:O	1:B:473[B]:TYR:HD1	2.04	0.41
1:A:174:PRO:HD2	1:A:239:MET:SD	2.61	0.41
4:B:599:FAD:O5B	4:B:599:FAD:H8A	2.21	0.41
2:C:663:LEU:HD12	2:C:663:LEU:HA	1.91	0.41
1:A:453:ARG:NH2	6:A:781:HOH:O	2.55	0.40
1:B:10:PRO:HG3	1:B:104:GLY:O	2.20	0.40
1:B:293:THR:HG22	1:B:385:ASN:O	2.21	0.40
1:B:174:PRO:HD2	1:B:239:MET:SD	2.62	0.40
2:C:617:HIS:O	2:C:623:VAL:HG11	2.21	0.40
1:B:278:VAL:HB	1:B:430:VAL:HG22	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	513/521 (98%)	499 (97%)	12 (2%)	2 (0%)	34	54
1	B	519/521 (100%)	504 (97%)	13 (2%)	2 (0%)	34	54
2	C	71/80 (89%)	68 (96%)	3 (4%)	0	100	100
2	D	71/80 (89%)	68 (96%)	3 (4%)	0	100	100
All	All	1174/1202 (98%)	1139 (97%)	31 (3%)	4 (0%)	41	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	156	SER
1	B	156	SER
1	A	360	GLY
1	B	360	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/433 (99%)	401 (94%)	26 (6%)	18	36
1	B	433/433 (100%)	409 (94%)	24 (6%)	21	41
2	C	60/64 (94%)	58 (97%)	2 (3%)	38	64
2	D	60/64 (94%)	57 (95%)	3 (5%)	24	46
All	All	980/994 (99%)	925 (94%)	55 (6%)	21	40

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	19	ASN
1	A	36	VAL
1	A	42	VAL
1	A	45	ASN
1	A	65	THR
1	A	84	ILE
1	A	114	ASN
1	A	152	PHE
1	A	167	ASP
1	A	180	MET
1	A	272	ASN
1	A	277	SER
1	A	308	VAL
1	A	323	LEU
1	A	357	GLU
1	A	361	ASP
1	A	362	THR
1	A	367	TYR
1	A	371	LEU
1	A	375	VAL
1	A	378	LEU
1	A	415	ARG
1	A	422	LEU
1	A	508	LEU
1	A	520	ASP
2	C	663	LEU
2	C	670	LEU
1	B	9	LEU
1	B	36	VAL
1	B	42	VAL
1	B	45	ASN
1	B	65	THR
1	B	84	ILE
1	B	114	ASN
1	B	152	PHE
1	B	180	MET
1	B	272	ASN
1	B	277	SER
1	B	308	VAL
1	B	323	LEU
1	B	357	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	361	ASP
1	B	362	THR
1	B	367	TYR
1	B	371	LEU
1	B	375	VAL
1	B	378	LEU
1	B	415	ARG
1	B	422	LEU
1	B	508	LEU
1	B	520	ASP
2	D	657	TYR
2	D	663	LEU
2	D	670	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	19	ASN
1	A	114	ASN
1	A	496	HIS
1	B	15	GLN
1	B	19	ASN
1	B	114	ASN
1	B	496	HIS
1	B	518	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FAD	A	599	1	51,58,58	2.05	13 (25%)	60,89,89	3.46	18 (30%)
4	FAD	B	599	1	51,58,58	2.14	14 (27%)	60,89,89	3.44	19 (31%)
5	HEC	D	699	2	26,50,50	1.93	4 (15%)	18,82,82	1.72	5 (27%)
5	HEC	C	699	2	26,50,50	1.82	5 (19%)	18,82,82	1.73	5 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	A	599	1	3/3/9/9	9/30/50/50	0/6/6/6
4	FAD	B	599	1	3/3/9/9	10/30/50/50	0/6/6/6
5	HEC	D	699	2	-	0/6/54/54	-
5	HEC	C	699	2	-	0/6/54/54	-

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	599	FAD	O4B-C1B	6.27	1.49	1.41
5	D	699	HEC	C3C-C2C	-5.90	1.34	1.40
5	C	699	HEC	C3C-C2C	-5.77	1.34	1.40
4	A	599	FAD	C9A-N10	5.75	1.46	1.38
4	B	599	FAD	C8A-N7A	-5.69	1.24	1.34
4	A	599	FAD	C8A-N7A	-5.13	1.25	1.34
4	B	599	FAD	C9A-N10	4.68	1.44	1.38
4	B	599	FAD	C4X-N5	4.63	1.39	1.33
5	D	699	HEC	C3B-C2B	-4.44	1.36	1.40
4	A	599	FAD	O4B-C1B	4.38	1.47	1.41
4	A	599	FAD	C4-N3	4.19	1.40	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	599	FAD	C4-N3	3.94	1.39	1.33
4	A	599	FAD	C4X-N5	3.63	1.38	1.33
4	B	599	FAD	C4X-C10	3.59	1.42	1.38
4	A	599	FAD	C10-N1	3.52	1.37	1.33
4	A	599	FAD	C8M-C8	-3.45	1.44	1.51
5	C	699	HEC	C3B-C2B	-3.35	1.37	1.40
4	A	599	FAD	C1'-N10	3.25	1.51	1.48
4	A	599	FAD	C2A-N3A	2.93	1.36	1.32
4	B	599	FAD	C5'-C4'	-2.88	1.47	1.51
4	B	599	FAD	C8M-C8	-2.88	1.45	1.51
4	B	599	FAD	C10-N1	2.80	1.36	1.33
4	B	599	FAD	C1'-N10	2.71	1.51	1.48
4	B	599	FAD	O5'-C5'	-2.62	1.34	1.44
4	B	599	FAD	C2A-N3A	2.44	1.36	1.32
4	A	599	FAD	C4X-C10	2.38	1.41	1.38
5	C	699	HEC	C1A-C2A	2.26	1.47	1.42
4	A	599	FAD	C4A-N3A	2.19	1.38	1.35
5	D	699	HEC	C1A-C2A	2.14	1.47	1.42
4	B	599	FAD	O3B-C3B	-2.11	1.38	1.43
5	C	699	HEC	C1D-ND	2.11	1.40	1.36
4	A	599	FAD	O3B-C3B	-2.09	1.38	1.43
4	A	599	FAD	O5'-C5'	-2.09	1.36	1.44
5	D	699	HEC	C4D-ND	2.05	1.40	1.36
4	B	599	FAD	PA-O2A	-2.03	1.45	1.55
5	C	699	HEC	CMD-C2D	2.00	1.55	1.51

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	599	FAD	C4-N3-C2	15.62	128.33	115.14
4	B	599	FAD	C4-N3-C2	15.41	128.15	115.14
4	A	599	FAD	C10-C4X-N5	11.84	129.44	121.26
4	B	599	FAD	C10-C4X-N5	11.44	129.17	121.26
4	B	599	FAD	C4-C4X-C10	-6.98	115.33	119.95
4	A	599	FAD	C4-C4X-C10	-6.83	115.43	119.95
4	A	599	FAD	C4X-C4-N3	-5.93	115.32	123.43
4	B	599	FAD	C5'-C4'-C3'	-5.89	100.83	112.20
4	A	599	FAD	C5'-C4'-C3'	-5.66	101.26	112.20
4	B	599	FAD	C4X-C4-N3	-5.59	115.78	123.43
4	B	599	FAD	O4'-C4'-C3'	5.24	121.83	109.10
4	A	599	FAD	O4'-C4'-C3'	5.22	121.79	109.10
4	B	599	FAD	C4X-N5-C5X	-5.17	111.60	116.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	599	FAD	C1'-N10-C9A	5.10	122.31	118.29
4	A	599	FAD	C4X-N5-C5X	-4.54	112.23	116.77
4	B	599	FAD	O3'-C3'-C4'	-4.38	98.24	108.81
4	A	599	FAD	O4'-C4'-C5'	4.38	119.75	109.92
4	A	599	FAD	O3'-C3'-C4'	-4.30	98.42	108.81
4	B	599	FAD	C1'-N10-C9A	4.05	121.48	118.29
4	B	599	FAD	O4'-C4'-C5'	3.92	118.73	109.92
4	A	599	FAD	C4X-C10-N10	-3.65	116.55	120.30
4	A	599	FAD	C6-C5X-N5	-3.61	115.07	119.05
4	B	599	FAD	C4X-C10-N10	-3.55	116.66	120.30
4	B	599	FAD	C6-C5X-N5	-3.42	115.28	119.05
5	D	699	HEC	CMC-C2C-C3C	3.30	129.70	125.82
4	A	599	FAD	C4'-C3'-C2'	3.24	120.11	113.36
4	B	599	FAD	C9A-N10-C10	-3.23	117.68	121.91
5	C	699	HEC	CBD-CAD-C3D	-3.16	106.66	112.49
4	A	599	FAD	C9A-N10-C10	-3.08	117.88	121.91
4	A	599	FAD	C4-C4X-N5	-3.04	115.12	118.60
4	B	599	FAD	C4'-C3'-C2'	3.02	119.65	113.36
5	C	699	HEC	CMC-C2C-C3C	2.93	129.27	125.82
4	B	599	FAD	P-O3P-PA	-2.84	123.09	132.83
4	A	599	FAD	P-O3P-PA	-2.83	123.11	132.83
5	D	699	HEC	CMC-C2C-C1C	-2.78	124.19	128.46
4	B	599	FAD	C9A-C5X-N5	2.78	126.71	122.36
4	B	599	FAD	O3'-C3'-C2'	2.75	115.45	108.81
4	B	599	FAD	C4-C4X-N5	-2.71	115.49	118.60
4	B	599	FAD	C1'-N10-C10	2.71	120.83	118.41
5	D	699	HEC	CMD-C2D-C1D	-2.67	124.36	128.46
4	A	599	FAD	C9A-C5X-N5	2.56	126.36	122.36
5	C	699	HEC	CMB-C2B-C1B	-2.50	124.62	128.46
5	D	699	HEC	CBD-CAD-C3D	-2.46	107.94	112.49
5	C	699	HEC	CMC-C2C-C1C	-2.40	124.77	128.46
4	A	599	FAD	O3'-C3'-C2'	2.31	114.38	108.81
5	C	699	HEC	CMB-C2B-C3B	2.29	128.52	125.82
5	D	699	HEC	CMB-C2B-C1B	-2.16	125.14	128.46

All (6) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	599	FAD	C4'
4	A	599	FAD	C2'
4	A	599	FAD	C3'
4	B	599	FAD	C4'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
4	B	599	FAD	C2'
4	B	599	FAD	C3'

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	599	FAD	N10-C1'-C2'-O2'
4	A	599	FAD	N10-C1'-C2'-C3'
4	A	599	FAD	C2'-C3'-C4'-C5'
4	A	599	FAD	O3'-C3'-C4'-C5'
4	A	599	FAD	C3'-C4'-C5'-O5'
4	A	599	FAD	O4'-C4'-C5'-O5'
4	B	599	FAD	N10-C1'-C2'-O2'
4	B	599	FAD	N10-C1'-C2'-C3'
4	B	599	FAD	C2'-C3'-C4'-C5'
4	B	599	FAD	O3'-C3'-C4'-C5'
4	B	599	FAD	C3'-C4'-C5'-O5'
4	B	599	FAD	O4'-C4'-C5'-O5'
4	A	599	FAD	O3'-C3'-C4'-O4'
4	B	599	FAD	O3'-C3'-C4'-O4'
4	A	599	FAD	C2'-C3'-C4'-O4'
4	B	599	FAD	C2'-C3'-C4'-O4'
4	A	599	FAD	C4'-C5'-O5'-P
4	B	599	FAD	C4'-C5'-O5'-P
4	B	599	FAD	C5'-O5'-P-O3P


There are no ring outliers.


2 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	599	FAD	2	0
4	B	599	FAD	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	515/521 (98%)	-0.67	4 (0%) 86 87	13, 36, 72, 91	0
1	B	515/521 (98%)	-0.65	9 (1%) 70 72	13, 36, 73, 91	0
2	C	73/80 (91%)	-0.50	0 100 100	24, 40, 67, 73	0
2	D	73/80 (91%)	-0.45	2 (2%) 54 58	27, 41, 66, 76	0
All	All	1176/1202 (97%)	-0.64	15 (1%) 77 79	13, 37, 72, 91	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	357	GLU	3.2
1	B	39	ASP	3.1
2	D	602	SER	3.1
1	A	349	GLY	2.8
1	B	7	ALA	2.6
1	B	361	ASP	2.6
2	D	633	GLY	2.4
1	B	360	GLY	2.3
1	B	145	ASN	2.3
1	A	32	ASP	2.1
1	A	7	ALA	2.1
1	A	145	ASN	2.0
1	B	295	ALA	2.0
1	B	19	ASN	2.0
1	B	23	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	HEC	C	699	43/43	0.97	0.10	23,33,44,48	0
5	HEC	D	699	43/43	0.97	0.11	16,45,55,57	0
4	FAD	A	599	53/53	0.98	0.08	6,21,37,44	0
4	FAD	B	599	53/53	0.98	0.08	8,22,39,52	0
3	CL	A	701	1/1	0.99	0.05	29,29,29,29	0
3	CL	B	702	1/1	0.99	0.04	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEC D 699:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 599:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 599:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.