

# wwPDB EM Validation Summary Report (i)

#### Nov 29, 2022 – 06:29 PM EST

PDB ID : 8DLY

EMDB ID : EMD-27520

Title: Cryo-EM structure of SARS-CoV-2 Epsilon (B.1.429) spike protein in complex

with VH ab6 (focused refinement of NTD and VH ab6)

Authors : Zhu, X.; Mannar, D.; Saville, J.W.; Srivastava, S.S.; Berezuk, A.M.; Zhou, S.;

Tuttle, K.S.; Subramaniam, S.

Deposited on : 2022-07-08

Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	С	1288	14% • 85%				
2	Н	119	92%	8%			



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	С	195	Total	С	N	О	S	0	0
1		190	1546	989	260	289	8	0	U

There are 93 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	13	ILE	SER	conflict	UNP P0DTC2
С	152	CYS	TRP	conflict	UNP P0DTC2
С	452	ARG	LEU	conflict	UNP P0DTC2
С	614	GLY	ASP	conflict	UNP P0DTC2
С	682	GLY	ARG	conflict	UNP P0DTC2
С	683	SER	ARG	conflict	UNP P0DTC2
С	685	SER	ARG	conflict	UNP P0DTC2
С	817	PRO	PHE	conflict	UNP P0DTC2
С	892	PRO	ALA	conflict	UNP P0DTC2
С	899	PRO	ALA	conflict	UNP P0DTC2
С	942	PRO	ALA	conflict	UNP P0DTC2
С	986	PRO	LYS	conflict	UNP P0DTC2
С	987	PRO	VAL	conflict	UNP P0DTC2
С	1209	GLY	-	expression tag	UNP P0DTC2
С	1210	SER	-	expression tag	UNP P0DTC2
С	1211	GLY	-	expression tag	UNP P0DTC2
С	1212	TYR	-	expression tag	UNP P0DTC2
С	1213	ILE	-	expression tag	UNP P0DTC2
С	1214	PRO	-	expression tag	UNP P0DTC2
С	1215	GLU	-	expression tag	UNP P0DTC2
С	1216	ALA	-	expression tag	UNP P0DTC2
С	1217	PRO	-	expression tag	UNP P0DTC2
С	1218	ARG	-	expression tag	UNP P0DTC2
С	1219	ASP	-	expression tag	UNP P0DTC2
С	1220	GLY	-	expression tag	UNP P0DTC2
С	1221	GLN	-	expression tag	UNP P0DTC2
С	1222	ALA	-	expression tag	UNP P0DTC2
С	1223	TYR	-	expression tag	UNP P0DTC2

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	1224	VAL	-	expression tag	UNP P0DTC2
C	1225	ARG	-	expression tag	UNP P0DTC2
С	1226	LYS	_	expression tag	UNP P0DTC2
$\Gamma$	1227	ASP	_	expression tag	UNP P0DTC2
С	1228	GLY	_	expression tag	UNP P0DTC2
С	1229	GLU	_	expression tag	UNP P0DTC2
С	1230	TRP	_	expression tag	UNP P0DTC2
С	1231	VAL	_	expression tag	UNP P0DTC2
С	1232	LEU	_	expression tag	UNP P0DTC2
С	1233	LEU	_	expression tag	UNP P0DTC2
С	1234	SER	_	expression tag	UNP P0DTC2
С	1235	THR	-	expression tag	UNP P0DTC2
С	1236	PHE	-	expression tag	UNP P0DTC2
С	1237	LEU	_	expression tag	UNP P0DTC2
С	1238	GLY	-	expression tag	UNP P0DTC2
С	1239	ARG	-	expression tag	UNP P0DTC2
С	1240	SER	-	expression tag	UNP P0DTC2
С	1241	LEU	-	expression tag	UNP P0DTC2
С	1242	GLU	-	expression tag	UNP P0DTC2
С	1243	VAL	-	expression tag	UNP P0DTC2
С	1244	LEU	-	expression tag	UNP P0DTC2
С	1245	PHE	-	expression tag	UNP P0DTC2
С	1246	GLN	-	expression tag	UNP P0DTC2
С	1247	GLY	-	expression tag	UNP P0DTC2
С	1248	PRO	-	expression tag	UNP P0DTC2
С	1249	GLY	-	expression tag	UNP P0DTC2
С	1250	HIS	-	expression tag	UNP P0DTC2
С	1251	HIS	-	expression tag	UNP P0DTC2
С	1252	HIS	-	expression tag	UNP P0DTC2
С	1253	HIS	_	expression tag	UNP P0DTC2
С	1254	HIS	-	expression tag	UNP P0DTC2
С	1255	HIS	-	expression tag	UNP P0DTC2
С	1256	HIS	-	expression tag	UNP P0DTC2
С	1257	HIS	-	expression tag	UNP P0DTC2
С	1258	SER	-	expression tag	UNP P0DTC2
С	1259	ALA	-	expression tag	UNP P0DTC2
С	1260	TRP	-	expression tag	UNP P0DTC2
С	1261	SER	-	expression tag	UNP P0DTC2
С	1262	HIS	-	expression tag	UNP P0DTC2
С	1263	PRO	-	expression tag	UNP P0DTC2
С	1264	GLN	-	expression tag	UNP P0DTC2
С	1265	PHE	-	expression tag	UNP PODTC2

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	1266	GLU	-	expression tag	UNP P0DTC2
С	1267	LYS	-	expression tag	UNP P0DTC2
С	1268	GLY	-	expression tag	UNP P0DTC2
С	1269	GLY	-	expression tag	UNP P0DTC2
С	1270	GLY	-	expression tag	UNP P0DTC2
С	1271	SER	-	expression tag	UNP P0DTC2
С	1272	GLY	-	expression tag	UNP P0DTC2
С	1273	GLY	-	expression tag	UNP P0DTC2
С	1274	GLY	-	expression tag	UNP P0DTC2
С	1275	GLY	-	expression tag	UNP P0DTC2
С	1276	SER	-	expression tag	UNP P0DTC2
С	1277	GLY	-	expression tag	UNP P0DTC2
С	1278	GLY	-	expression tag	UNP P0DTC2
С	1279	SER	-	expression tag	UNP P0DTC2
С	1280	ALA	-	expression tag	UNP P0DTC2
С	1281	TRP	-	expression tag	UNP P0DTC2
С	1282	SER	-	expression tag	UNP P0DTC2
С	1283	HIS	-	expression tag	UNP P0DTC2
С	1284	PRO	-	expression tag	UNP P0DTC2
С	1285	GLN	-	expression tag	UNP P0DTC2
С	1286	PHE	-	expression tag	UNP P0DTC2
С	1287	GLU	-	expression tag	UNP P0DTC2
С	1288	LYS	-	expression tag	UNP P0DTC2

 $\bullet$  Molecule 2 is a protein called VH ab 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	Н	119	Total 923	C 585	- 1	O 177	S 5	0	0

 $\bullet$  Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6).$ 






Mol	Chain	Residues	Atoms				AltConf
3	С	1	Total	С	N	О	0
3		1	14	8	1	5	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Spike glycoprotein









# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	122910	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Michagona	TFS KRIOS	Danagitan
Microscope		Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	40	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	2.038	Depositor
Minimum map value	-0.886	Depositor
Average map value	-0.002	Depositor
Map value standard deviation	0.032	Depositor
Recommended contour level	0.322	Depositor
Map size (Å)	400.0, 400.0, 400.0	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0, 1.0, 1.0	Depositor



## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	С	0.33	0/1590	0.59	0/2164	
2	Н	0.33	0/946	0.68	0/1284	
All	All	0.33	0/2536	0.62	0/3448	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	1546	0	1461	4	0
2	Н	923	0	888	5	0
3	С	14	0	13	0	0
All	All	2483	0	2362	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 8 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:35:HIS:HB2	2:H:96:ALA:HB3	1.79	0.64
2:H:93:TYR:O	2:H:112:GLY:HA2	2.09	0.52
1:C:418:ILE:HA	1:C:422:ASN:HB2	1.93	0.50
1:C:403:ARG:HG3	1:C:406:GLU:HG3	1.97	0.46
2:H:11:VAL:HG22	2:H:116:THR:HB	1.98	0.46

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	$\mathbf{C}$	193/1288 (15%)	184 (95%)	9 (5%)	0	100	100
2	Н	117/119 (98%)	115 (98%)	2 (2%)	0	100	100
All	All	310/1407 (22%)	299 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	$\mathbf{C}$	168/1115 (15%)	166 (99%)	2 (1%)	71 90
2	Н	99/99 (100%)	99 (100%)	0	100 100
All	All	267/1214 (22%)	265 (99%)	2 (1%)	84 94



All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	452	ARG
1	С	492	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	${ m ths}$	$\mathbf{B}$	ond ang	les
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	С	1301	1	14,14,15	0.51	0	17,19,21	0.80	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	С	1301	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	С	1301	NAG	C1-O5-C5	3.08	116.37	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	1301	NAG	O5-C5-C6-O6
3	С	1301	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-27520. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



#### 6.1.2 Raw map



The images above show the map projected in three orthogonal directions.




### 6.2 Central slices (i)

#### 6.2.1 Primary map







X Index: 200


Y Index: 200

Z Index: 200

#### 6.2.2 Raw map







X Index: 200

Y Index: 200

Z Index: 200

The images above show central slices of the map in three orthogonal directions.



### 6.3 Largest variance slices (i)


#### 6.3.1 Primary map







Y Index: 190



Z Index: 252


#### 6.3.2 Raw map



X Index: 187



Y Index: 190



Z Index: 254

The images above show the largest variance slices of the map in three orthogonal directions.



#### 6.4 Orthogonal surface views (i)

#### 6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.322. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

### 6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis (i)

This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution (i)




The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



EMD-27520, 8DLY

#### 7.2 Volume estimate (i)




The volume at the recommended contour level is  $102~\mathrm{nm}^3$ ; this corresponds to an approximate mass of  $92~\mathrm{kDa}$ .

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum (i)




<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.333  $\rm \mathring{A}^{-1}$ 



### 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

#### 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.333  $\rm \mathring{A}^{-1}$ 



## 8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
rtesolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	3.00	-	-	
Author-provided FSC curve	3.02	4.00	3.08	
Unmasked-calculated*	4.10	8.80	4.20	


<sup>\*</sup>Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.10 differs from the reported value 3.0 by more than 10 %



## 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-27520 and PDB model 8DLY. Per-residue inclusion information can be found in section 3 on page 7.


### 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.322 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


#### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.322).



## 9.4 Atom inclusion (i)



At the recommended contour level, 96% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.



### 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.322) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8526	0.4950
С	0.8720	0.5050
Н	0.8198	0.4780



