



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2023 – 12:39 AM EDT

PDB ID : 5DO8  
Title : 1.8 Angstrom crystal structure of *Listeria monocytogenes* Lmo0184 alpha-1,6-glucosidase  
Authors : Light, S.H.; Halavaty, A.S.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)  
Deposited on : 2015-09-10  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	 96%
1	B	555	 96%
1	C	555	 96%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lmo0184 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	553	4629	2936	764	907	22	0	8	0
1	A	552	4655	2946	771	917	21	0	9	0
1	C	551	4614	2922	763	908	21	0	6	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	554	GLY	-	expression tag	UNP Q8YAE6
B	555	ALA	-	expression tag	UNP Q8YAE6
A	554	GLY	-	expression tag	UNP Q8YAE6
A	555	ALA	-	expression tag	UNP Q8YAE6
C	554	GLY	-	expression tag	UNP Q8YAE6
C	555	ALA	-	expression tag	UNP Q8YAE6

- Molecule 2 is beta-D-glucopyranose (three-letter code: BGC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 12 6 6	0	0
2	A	1	Total C O 12 6 6	0	0
2	C	1	Total C O 12 6 6	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0
3	A	1	Total Cl 1 1	0	0
3	C	2	Total Cl 2 2	0	0

- Molecule 4 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C<sub>8</sub>H<sub>19</sub>NO<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	632	Total	O	0	4
			632	632		
5	A	627	Total	O	0	3
			627	627		
5	C	502	Total	O	0	0
			502	502		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lmo0184 protein



- Molecule 1: Lmo0184 protein



- Molecule 1: Lmo0184 protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.12Å 103.94Å 193.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80 29.81 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.1 (30.00-1.80) 99.2 (29.81-1.80)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.74 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, $R_{free}$	0.154 , 0.188 0.163 , 0.196	Depositor DCC
$R_{free}$ test set	7982 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.2	Xtrriage
Anisotropy	0.176	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	15741	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, BGC, BTB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/4774	0.71	2/6474 (0.0%)
1	B	0.47	0/4770	0.70	2/6469 (0.0%)
1	C	0.45	0/4739	0.68	0/6427
All	All	0.46	0/14283	0.70	4/19370 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	341	ARG	NE-CZ-NH1	6.53	123.56	120.30
1	B	341	ARG	NE-CZ-NH2	-5.73	117.44	120.30
1	A	176	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	394	ARG	NE-CZ-NH2	5.24	122.92	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4655	0	4380	6	0
1	B	4629	0	4381	9	0
1	C	4614	0	4345	5	0
2	A	12	0	12	0	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	12	0	12	0	0
2	C	12	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	2	0	0	0	0
4	A	14	0	19	0	0
4	B	14	0	19	0	0
4	C	14	0	19	0	0
5	A	627	0	0	1	0
5	B	632	0	0	1	0
5	C	502	0	0	0	0
All	All	15741	0	13199	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:14:GLN:HE22	1:B:326:ASN:HB2	1.68	0.59
1:A:390:MET:SD	1:A:394:ARG:HD3	2.43	0.58
1:A:14:GLN:HE22	1:A:326:ASN:HB2	1.68	0.57
1:C:14:GLN:HE22	1:C:326:ASN:HB2	1.70	0.56
1:C:281:GLU:HB3	1:C:306:ASN:HD21	1.75	0.51
1:B:338:ASN:O	1:B:343:ARG:HA	2.11	0.50
1:B:305:GLU:HG2	5:B:704:HOH:O	2.16	0.46
1:C:338:ASN:O	1:C:343:ARG:HA	2.16	0.45
1:B:218:ILE:HG23	1:B:219:TYR:CD2	2.52	0.44
1:B:375[A]:HIS:CE1	1:B:409:TYR:CE2	3.06	0.44
1:A:419:PRO:HB3	1:A:430:PHE:CG	2.52	0.44
1:A:101:VAL:HG22	1:A:197:ASP:OD2	2.17	0.43
1:A:338:ASN:O	1:A:343:ARG:HA	2.17	0.43
1:B:233:HIS:CE1	1:B:266:TYR:CE1	3.06	0.43
1:A:305[B]:GLU:HG3	5:A:704:HOH:O	2.17	0.43
1:B:101:VAL:HG22	1:B:197:ASP:OD2	2.20	0.42
1:B:66:ILE:HD12	1:B:101:VAL:HB	2.00	0.42
1:C:66:ILE:HD12	1:C:101:VAL:HB	2.01	0.41
1:B:341:ARG:HA	1:B:341:ARG:NE	2.35	0.41
1:C:101:VAL:HG22	1:C:197:ASP:OD2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	559/555 (101%)	544 (97%)	15 (3%)	0	100	100
1	B	559/555 (101%)	542 (97%)	17 (3%)	0	100	100
1	C	555/555 (100%)	539 (97%)	16 (3%)	0	100	100
All	All	1673/1665 (100%)	1625 (97%)	48 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	513/506 (101%)	507 (99%)	6 (1%)	71	65
1	B	512/506 (101%)	506 (99%)	6 (1%)	71	65
1	C	509/506 (101%)	502 (99%)	7 (1%)	67	59
All	All	1534/1518 (101%)	1515 (99%)	19 (1%)	71	65

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	167	ASP
1	B	288	ASP
1	B	325	ASN
1	B	469	ARG
1	B	481	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	491	PHE
1	A	167	ASP
1	A	325	ASN
1	A	450[A]	GLU
1	A	450[B]	GLU
1	A	469	ARG
1	A	491	PHE
1	C	133	ASN
1	C	167	ASP
1	C	325	ASN
1	C	340	ASN
1	C	341	ARG
1	C	469	ARG
1	C	491	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	14	GLN
1	A	14	GLN
1	A	224	ASN
1	A	522	ASN
1	C	14	GLN
1	C	133	ASN
1	C	427	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	BTB	A	603	-	13,13,13	1.06	2 (15%)	7,16,16	0.40	0
2	BGC	A	601	-	12,12,12	0.48	0	17,17,17	0.74	0
4	BTB	C	604	-	13,13,13	0.94	1 (7%)	7,16,16	0.60	0
4	BTB	B	603	-	13,13,13	0.98	1 (7%)	7,16,16	1.07	0
2	BGC	C	601	-	12,12,12	0.47	0	17,17,17	0.46	0
2	BGC	B	601	-	12,12,12	0.42	0	17,17,17	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BTB	A	603	-	-	3/21/21/21	-
2	BGC	A	601	-	-	0/2/22/22	0/1/1/1
4	BTB	C	604	-	-	1/21/21/21	-
4	BTB	B	603	-	-	12/21/21/21	-
2	BGC	C	601	-	-	0/2/22/22	0/1/1/1
2	BGC	B	601	-	-	0/2/22/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	603	BTB	C5-N	2.33	1.51	1.48
4	B	603	BTB	C2-N	2.32	1.53	1.48
4	C	604	BTB	C7-N	2.18	1.51	1.48
4	A	603	BTB	C7-N	2.01	1.50	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	603	BTB	C1-C2-C4-O4
4	B	603	BTB	C3-C2-C4-O4
4	B	603	BTB	N-C2-C4-O4
4	B	603	BTB	C1-C2-N-C5
4	B	603	BTB	C1-C2-N-C7
4	B	603	BTB	C3-C2-N-C5
4	B	603	BTB	C4-C2-N-C5
4	B	603	BTB	C4-C2-N-C7
4	B	603	BTB	N-C5-C6-O6
4	A	603	BTB	C1-C2-C3-O3
4	A	603	BTB	C4-C2-C3-O3
4	A	603	BTB	N-C2-C3-O3
4	C	604	BTB	O1-C1-C2-C4
4	B	603	BTB	N-C7-C8-O8
4	B	603	BTB	C6-C5-N-C2
4	B	603	BTB	C3-C2-N-C7

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	552/555 (99%)	-0.39	8 (1%) 75 72	12, 19, 33, 59	0
1	B	553/555 (99%)	-0.35	13 (2%) 59 54	12, 19, 36, 59	0
1	C	551/555 (99%)	-0.18	17 (3%) 49 43	15, 23, 40, 62	0
All	All	1656/1665 (99%)	-0.31	38 (2%) 60 56	12, 20, 38, 62	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	555	ALA	6.2
1	B	215	ASP	5.2
1	A	554	GLY	4.7
1	C	215	ASP	4.3
1	B	216	GLY	4.2
1	C	216	GLY	4.1
1	A	216	GLY	4.0
1	A	217	GLN	3.8
1	A	3	GLU	3.8
1	B	554	GLY	3.7
1	C	217	GLN	3.4
1	B	3	GLU	3.2
1	C	218	ILE	3.2
1	A	215	ASP	3.1
1	C	554	GLY	2.9
1	C	4	LYS	2.9
1	C	498	GLU	2.8
1	B	217	GLN	2.8
1	C	288	ASP	2.7
1	C	214	PRO	2.7
1	B	4	LYS	2.7
1	C	499	ASN	2.6
1	C	518	GLU	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	290	ASP	2.4
1	B	290	ASP	2.4
1	C	341	ARG	2.3
1	C	100	LEU	2.2
1	A	133	ASN	2.2
1	C	316	GLU	2.2
1	C	133	ASN	2.1
1	B	288	ASP	2.1
1	A	241	GLN	2.1
1	B	289	SER	2.1
1	B	214	PRO	2.1
1	C	553	LYS	2.1
1	A	214	PRO	2.1
1	B	133	ASN	2.0
1	B	218	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	BTB	C	604	14/14	0.67	0.23	42,48,51,51	0
4	BTB	B	603	14/14	0.76	0.24	38,41,43,43	0
4	BTB	A	603	14/14	0.79	0.19	36,38,40,40	0
2	BGC	C	601	12/12	0.96	0.11	16,18,18,21	0
3	CL	C	603	1/1	0.97	0.06	29,29,29,29	0
2	BGC	A	601	12/12	0.98	0.10	13,14,15,17	0
2	BGC	B	601	12/12	0.98	0.11	14,15,16,17	0
3	CL	B	602	1/1	0.99	0.04	26,26,26,26	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	602	1/1	1.00	0.03	24,24,24,24	0
3	CL	C	602	1/1	1.00	0.04	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.