

wwPDB X-ray Structure Validation Summary Report (i)

Oct 15, 2023 – 03:09 AM EDT

PDB ID : 8DPC

Title : Crystal structure of carbonic anhydrase from Neisseria gonorrhoeae

Authors: Marapaka, A.K.; Das, C.; Flaherty, D.P.; Yadav, R.

Deposited on : 2022-07-15

Resolution : 2.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	243	79%	11% • 9%
1	С	243	77%	11% • 9%
1	Е	243	81%	9% 9%
1	G	243	7%	12% • 9%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7145 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Carbonic anhydrase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	222	Total	С	N	О	S	0	0	0
1	A	222	1732	1098	305	323	6	0	U	U
1	С	222	Total	С	N	О	S	0	0	0
1		222	1732	1097	305	324	6	U	U	0
1	E	222	Total	С	N	О	S	0	0	0
1	15	222	1747	1105	308	328	6	0	U	U
1	G	222	Total	С	N	О	S	0	0	0
1	G	222	1737	1099	304	328	6	0		U

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	HIS	-	expression tag	UNP Q50940
A	-15	HIS	-	expression tag	UNP Q50940
A	-14	HIS	-	expression tag	UNP Q50940
A	-13	HIS	-	expression tag	UNP Q50940
A	-12	HIS	-	expression tag	UNP Q50940
A	-11	HIS	-	expression tag	UNP Q50940
A	-10	ASP	-	expression tag	UNP Q50940
A	-9	SER	-	expression tag	UNP Q50940
A	-8	GLY	-	expression tag	UNP Q50940
A	-7	LEU	-	expression tag	UNP Q50940
A	-6	VAL	-	expression tag	UNP Q50940
A	-5	PRO	ı	expression tag	UNP Q50940
A	-4	ARG	-	expression tag	UNP Q50940
A	-3	GLY	ı	expression tag	UNP Q50940
A	-2	SER	-	expression tag	UNP Q50940
A	-1	HIS	-	expression tag	UNP Q50940
A	0	MET	=	expression tag	UNP Q50940
С	-16	HIS	-	expression tag	UNP Q50940
С	-15	HIS	=	expression tag	UNP Q50940
С	-14	HIS	-	expression tag	UNP Q50940
С	-13	HIS	-	expression tag	UNP Q50940



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
С	-12	HIS	-	expression tag	UNP Q50940
С	-11	HIS	-	expression tag	UNP Q50940
С	-10	ASP	-	expression tag	UNP Q50940
С	-9	SER	-	expression tag	UNP Q50940
С	-8	GLY	-	expression tag	UNP Q50940
С	-7	LEU	-	expression tag	UNP Q50940
С	-6	VAL	-	expression tag	UNP Q50940
С	-5	PRO	-	expression tag	UNP Q50940
С	-4	ARG	-	expression tag	UNP Q50940
С	-3	GLY	-	expression tag	UNP Q50940
С	-2	SER	-	expression tag	UNP Q50940
С	-1	HIS	-	expression tag	UNP Q50940
С	0	MET	-	expression tag	UNP Q50940
E	-16	HIS	-	expression tag	UNP Q50940
Е	-15	HIS	-	expression tag	UNP Q50940
Е	-14	HIS	-	expression tag	UNP Q50940
Е	-13	HIS	-	expression tag	UNP Q50940
Е	-12	HIS	-	expression tag	UNP Q50940
E	-11	HIS	-	expression tag	UNP Q50940
Е	-10	ASP	-	expression tag	UNP Q50940
Е	-9	SER	-	expression tag	UNP Q50940
E	-8	GLY	-	expression tag	UNP Q50940
Е	-7	LEU	-	expression tag	UNP Q50940
Е	-6	VAL	-	expression tag	UNP Q50940
Е	-5	PRO	-	expression tag	UNP Q50940
Е	-4	ARG	-	expression tag	UNP Q50940
Е	-3	GLY	-	expression tag	UNP Q50940
Е	-2	SER	-	expression tag	UNP Q50940
Е	-1	HIS	-	expression tag	UNP Q50940
Е	0	MET	-	expression tag	UNP Q50940
G	-16	HIS	-	expression tag	UNP Q50940
G	-15	HIS	-	expression tag	UNP Q50940
G	-14	HIS	-	expression tag	UNP Q50940
G	-13	HIS	-	expression tag	UNP Q50940
G	-12	HIS	-	expression tag	UNP Q50940
G	-11	HIS	-	expression tag	UNP Q50940
G	-10	ASP	_	expression tag	UNP Q50940
G	-9	SER	-	expression tag	UNP Q50940
G	-8	GLY	-	expression tag	UNP Q50940
G	-7	LEU	-	expression tag	UNP Q50940
G	-6	VAL	-	expression tag	UNP Q50940
G	-5	PRO	-	expression tag	UNP Q50940




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-4	ARG	-	expression tag	UNP Q50940
G	-3	GLY	-	expression tag	UNP Q50940
G	-2	SER	-	expression tag	UNP Q50940
G	-1	HIS	-	expression tag	UNP Q50940
G	0	MET	-	expression tag	UNP Q50940

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	С	1	Total Zn 1 1	0	0
2	Е	1	Total Zn 1 1	0	0
2	G	1	$\begin{array}{cc} \text{Total} & \text{Zn} \\ 1 & 1 \end{array}$	0	0

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S	0	0
3	A	1	5 4 1 Total O S	0	0
- J	А	1	5 4 1	0	U
3	С	1	Total O S 5 4 1	0	0
3	С	1	Total O S 5 4 1	0	0
3	С	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	Е	1	Total O S 5 4 1	0	0
3	G	1	Total O S 5 4 1	0	0
3	G	1	Total O S 5 4 1	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	35	Total O 35 35	0	0
4	С	32	Total O 32 32	0	0
4	E	34	Total O 34 34	0	0
4	G	37	Total O 37 37	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbonic anhydrase







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	71.16Å 122.76Å 78.67Å	Depositor
a, b, c, α , β , γ	90.00° 116.87° 90.00°	Depositor
Resolution (Å)	44.12 - 2.41	Depositor
Resolution (A)	44.12 - 2.41	EDS
% Data completeness	99.4 (44.12-2.41)	Depositor
(in resolution range)	99.4 (44.12-2.41)	EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.45 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.190 , 0.246	Depositor
it, it free	0.195 , 0.250	DCC
R_{free} test set	2281 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 43.6	EDS
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	0.028 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7145	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.41	0/1778	0.78	1/2422 (0.0%)
1	С	0.40	0/1778	0.80	0/2423
1	Е	0.40	0/1793	0.74	0/2441
1	G	0.38	0/1783	0.72	0/2430
All	All	0.40	0/7132	0.76	1/9716 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	С	0	4
1	Е	0	1
1	G	0	6
All	All	0	13

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
1	A	136	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

5 of 13 planarity outliers are listed below:

Mo	ol	Chain	Res	Type	Group
1		A	166	ARG	Sidechain



Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	171	ARG	Sidechain
1	С	154	ASN	Peptide
1	С	26	ARG	Sidechain
1	С	84	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1732	0	1686	21	0
1	С	1732	0	1681	23	0
1	Е	1747	0	1706	9	0
1	G	1737	0	1684	21	0
2	A	1	0	0	0	0
2	С	1	0	0	0	0
2	Е	1	0	0	0	0
2	G	1	0	0	0	0
3	A	20	0	0	1	0
3	С	15	0	0	1	0
3	Е	10	0	0	0	0
3	G	10	0	0	0	0
4	A	35	0	0	4	0
4	С	32	0	0	1	0
4	Е	34	0	0	1	0
4	G	37	0	0	12	0
All	All	7145	0	6757	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:C:152:LYS:O	1:C:154:ASN:N	1.82	1.11
1:C:152:LYS:C	1:C:154:ASN:H	1.75	0.90
1:G:28:CYS:SG	4:G:423:HOH:O	2.30	0.87



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:153:LEU:HG	1:A:155:GLN:CB	2.06	0.84
1:A:73:PRO:HA	1:C:154:ASN:HA	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	220/243 (90%)	212 (96%)	7 (3%)	1 (0%)	29	40
1	C	$220/243 \ (90\%)$	210 (96%)	7 (3%)	3 (1%)	11	14
1	E	$220/243 \ (90\%)$	214 (97%)	6 (3%)	0	100	100
1	G	$220/243 \ (90\%)$	203 (92%)	16 (7%)	1 (0%)	29	40
All	All	880/972 (90%)	839 (95%)	36 (4%)	5 (1%)	25	35

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	148	ALA
1	С	153	LEU
1	G	148	ALA
1	A	148	ALA
1	С	155	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed, and the total nu	imber of residues.
----------------------------	--------------------

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	189/213 (89%)	179 (95%)	10 (5%)	22 36
1	С	189/213 (89%)	176 (93%)	13 (7%)	15 24
1	E	193/213 (91%)	186 (96%)	7 (4%)	35 52
1	G	191/213 (90%)	185 (97%)	6 (3%)	40 58
All	All	762/852 (89%)	726 (95%)	36 (5%)	26 41

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	152	LYS
1	G	166	ARG
1	Е	165	LYS
1	G	29	SER
1	С	22	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	69	GLN
1	G	33	ASN
1	С	69	GLN
1	Ε	142	ASN
1	С	66	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 15 ligands modelled in this entry, 4 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	Chain	Chain	Chain	Chain	Dag	Link	В	Bond lengths			Bond angles		
MIOI	Mol Type Chain	n Res	Res	tes Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2				
3	SO4	С	304	-	4,4,4	0.46	0	6,6,6	0.17	0				
3	SO4	Е	302	-	4,4,4	0.48	0	6,6,6	0.17	0				
3	SO4	A	302	-	4,4,4	0.45	0	6,6,6	0.18	0				
3	SO4	Е	303	-	4,4,4	0.30	0	6,6,6	0.16	0				
3	SO4	A	304	-	4,4,4	0.30	0	6,6,6	0.08	0				
3	SO4	С	303	-	4,4,4	0.34	0	6,6,6	0.16	0				
3	SO4	A	305	-	4,4,4	0.34	0	6,6,6	0.14	0				
3	SO4	A	303	-	4,4,4	0.31	0	6,6,6	0.15	0				
3	SO4	G	302	-	4,4,4	0.41	0	6,6,6	0.28	0				
3	SO4	С	302	-	4,4,4	0.45	0	6,6,6	0.20	0				
3	SO4	G	303	-	4,4,4	0.30	0	6,6,6	0.09	0				

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	SO4	1	0
3	С	302	SO4	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be




highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		$OWAB(A^2)$	Q<0.9	
1	A	222/243 (91%)	-0.35	3 (1%)	75	73	13, 25, 48, 101	0
1	С	222/243 (91%)	-0.52	1 (0%)	91	89	15, 25, 45, 68	0
1	E	222/243 (91%)	-0.49	2 (0%)	84	82	15, 28, 50, 78	0
1	G	222/243 (91%)	0.04	16 (7%)	15	13	20, 38, 74, 101	0
All	All	888/972 (91%)	-0.33	22 (2%)	57	54	13, 29, 56, 101	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	154	ASN	11.9
1	G	6	HIS	6.0
1	G	23	GLU	5.8
1	G	12	HIS	5.6
1	G	5	THR	5.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	ZN	G	301	1/1	0.84	0.16	76,76,76,76	0
3	SO4	G	303	5/5	0.87	0.19	66,66,75,79	0
3	SO4	A	303	5/5	0.92	0.12	53,54,63,66	0
3	SO4	Е	303	5/5	0.94	0.14	59,74,76,78	0
2	ZN	Е	301	1/1	0.96	0.28	68,68,68,68	0
3	SO4	A	305	5/5	0.96	0.20	46,51,53,57	0
3	SO4	С	303	5/5	0.97	0.13	48,52,58,64	0
3	SO4	A	304	5/5	0.97	0.15	49,50,52,59	0
3	SO4	G	302	5/5	0.97	0.12	42,42,43,46	0
2	ZN	С	301	1/1	0.97	0.15	59,59,59,59	0
3	SO4	С	304	5/5	0.98	0.09	44,47,48,52	0
3	SO4	Е	302	5/5	0.99	0.10	36,37,40,43	0
3	SO4	С	302	5/5	0.99	0.11	34,35,40,43	0
3	SO4	A	302	5/5	0.99	0.10	27,28,30,34	0
2	ZN	A	301	1/1	0.99	0.07	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.










Electron density around SO4 G 303:

 $2mF_o$ -DF_c (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)













Electron density around SO4 A 303:





















Electron density around SO4 A 305:













Electron density around SO4 C 303:















Electron density around SO4 G 302:














Electron density around SO4 C 304:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















Electron density around SO4 C 302:













Electron density around SO4 A 302:











6.5 Other polymers (i)

There are no such residues in this entry.

