



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 1, 2022 – 10:10 AM EDT

PDB ID : 8DTC
Title : Crystal Structure of Glucokinase with bound glucose from *Acanthamoeba castellanii*
Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2022-07-25
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.29
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	 3% 89% 5% 6%
1	B	398	 5% 89% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	374	2923	1863	488	562	10	0	9	0
1	B	374	2912	1854	487	559	12	0	7	0

There are 126 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	expression tag	UNP L8GT25
A	-9	ARG	-	expression tag	UNP L8GT25
A	-8	GLY	-	expression tag	UNP L8GT25
A	-7	SER	-	expression tag	UNP L8GT25
A	-6	HIS	-	expression tag	UNP L8GT25
A	-5	HIS	-	expression tag	UNP L8GT25
A	-4	HIS	-	expression tag	UNP L8GT25
A	-3	HIS	-	expression tag	UNP L8GT25
A	-2	HIS	-	expression tag	UNP L8GT25
A	-1	HIS	-	expression tag	UNP L8GT25
A	0	GLY	-	expression tag	UNP L8GT25
A	1	SER	-	expression tag	UNP L8GT25
A	16	GLY	VAL	conflict	UNP L8GT25
A	25	HIS	GLN	conflict	UNP L8GT25
A	59	PRO	SER	conflict	UNP L8GT25
A	105	GLY	SER	conflict	UNP L8GT25
A	142	TYR	PHE	conflict	UNP L8GT25
A	160	ASN	GLN	conflict	UNP L8GT25
A	163	ALA	SER	conflict	UNP L8GT25
A	177	THR	-	insertion	UNP L8GT25
A	178	MET	-	insertion	UNP L8GT25
A	179	LYS	-	insertion	UNP L8GT25
A	180	ARG	-	insertion	UNP L8GT25
A	181	ALA	-	insertion	UNP L8GT25
A	182	ASN	-	insertion	UNP L8GT25

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	183	TYR	-	insertion	UNP L8GT25
A	184	LEU	-	insertion	UNP L8GT25
A	185	VAL	-	insertion	UNP L8GT25
A	186	LEU	-	insertion	UNP L8GT25
A	187	ALA	-	insertion	UNP L8GT25
A	188	VAL	-	insertion	UNP L8GT25
A	189	GLY	-	insertion	UNP L8GT25
A	190	THR	-	insertion	UNP L8GT25
A	191	GLY	-	insertion	UNP L8GT25
A	192	LEU	-	insertion	UNP L8GT25
A	193	GLY	-	insertion	UNP L8GT25
A	194	ILE	-	insertion	UNP L8GT25
A	195	ALA	-	insertion	UNP L8GT25
A	196	LEU	-	insertion	UNP L8GT25
A	197	LEU	-	insertion	UNP L8GT25
A	198	THR	-	insertion	UNP L8GT25
A	199	SER	-	insertion	UNP L8GT25
A	200	LEU	-	insertion	UNP L8GT25
A	201	GLY	-	insertion	UNP L8GT25
A	202	ARG	-	insertion	UNP L8GT25
A	203	GLY	-	insertion	UNP L8GT25
A	204	SER	-	insertion	UNP L8GT25
A	205	ARG	-	insertion	UNP L8GT25
A	206	ASN	-	insertion	UNP L8GT25
A	207	ILE	-	insertion	UNP L8GT25
A	208	PRO	-	insertion	UNP L8GT25
A	209	LEU	-	insertion	UNP L8GT25
A	210	GLN	-	insertion	UNP L8GT25
A	211	VAL	-	insertion	UNP L8GT25
A	224	VAL	ALA	conflict	UNP L8GT25
A	228	ALA	SER	conflict	UNP L8GT25
A	249	ASN	HIS	conflict	UNP L8GT25
A	285	ILE	VAL	conflict	UNP L8GT25
A	287	ALA	THR	conflict	UNP L8GT25
A	342	TYR	HIS	conflict	UNP L8GT25
A	344	ALA	PRO	conflict	UNP L8GT25
A	361	LYS	MET	conflict	UNP L8GT25
A	369	GLN	GLU	conflict	UNP L8GT25
B	-10	MET	-	expression tag	UNP L8GT25
B	-9	ARG	-	expression tag	UNP L8GT25
B	-8	GLY	-	expression tag	UNP L8GT25
B	-7	SER	-	expression tag	UNP L8GT25

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	HIS	-	expression tag	UNP L8GT25
B	-5	HIS	-	expression tag	UNP L8GT25
B	-4	HIS	-	expression tag	UNP L8GT25
B	-3	HIS	-	expression tag	UNP L8GT25
B	-2	HIS	-	expression tag	UNP L8GT25
B	-1	HIS	-	expression tag	UNP L8GT25
B	0	GLY	-	expression tag	UNP L8GT25
B	1	SER	-	expression tag	UNP L8GT25
B	16	GLY	VAL	conflict	UNP L8GT25
B	25	HIS	GLN	conflict	UNP L8GT25
B	59	PRO	SER	conflict	UNP L8GT25
B	105	GLY	SER	conflict	UNP L8GT25
B	142	TYR	PHE	conflict	UNP L8GT25
B	160	ASN	GLN	conflict	UNP L8GT25
B	163	ALA	SER	conflict	UNP L8GT25
B	177	THR	-	insertion	UNP L8GT25
B	178	MET	-	insertion	UNP L8GT25
B	179	LYS	-	insertion	UNP L8GT25
B	180	ARG	-	insertion	UNP L8GT25
B	181	ALA	-	insertion	UNP L8GT25
B	182	ASN	-	insertion	UNP L8GT25
B	183	TYR	-	insertion	UNP L8GT25
B	184	LEU	-	insertion	UNP L8GT25
B	185	VAL	-	insertion	UNP L8GT25
B	186	LEU	-	insertion	UNP L8GT25
B	187	ALA	-	insertion	UNP L8GT25
B	188	VAL	-	insertion	UNP L8GT25
B	189	GLY	-	insertion	UNP L8GT25
B	190	THR	-	insertion	UNP L8GT25
B	191	GLY	-	insertion	UNP L8GT25
B	192	LEU	-	insertion	UNP L8GT25
B	193	GLY	-	insertion	UNP L8GT25
B	194	ILE	-	insertion	UNP L8GT25
B	195	ALA	-	insertion	UNP L8GT25
B	196	LEU	-	insertion	UNP L8GT25
B	197	LEU	-	insertion	UNP L8GT25
B	198	THR	-	insertion	UNP L8GT25
B	199	SER	-	insertion	UNP L8GT25
B	200	LEU	-	insertion	UNP L8GT25
B	201	GLY	-	insertion	UNP L8GT25
B	202	ARG	-	insertion	UNP L8GT25
B	203	GLY	-	insertion	UNP L8GT25

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	204	SER	-	insertion	UNP L8GT25
B	205	ARG	-	insertion	UNP L8GT25
B	206	ASN	-	insertion	UNP L8GT25
B	207	ILE	-	insertion	UNP L8GT25
B	208	PRO	-	insertion	UNP L8GT25
B	209	LEU	-	insertion	UNP L8GT25
B	210	GLN	-	insertion	UNP L8GT25
B	211	VAL	-	insertion	UNP L8GT25
B	224	VAL	ALA	conflict	UNP L8GT25
B	228	ALA	SER	conflict	UNP L8GT25
B	249	ASN	HIS	conflict	UNP L8GT25
B	285	ILE	VAL	conflict	UNP L8GT25
B	287	ALA	THR	conflict	UNP L8GT25
B	342	TYR	HIS	conflict	UNP L8GT25
B	344	ALA	PRO	conflict	UNP L8GT25
B	361	LYS	MET	conflict	UNP L8GT25
B	369	GLN	GLU	conflict	UNP L8GT25

- Molecule 2 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	250	Total O 255 255	0	5
5	B	299	Total O 303 303	0	4

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	114.99Å 52.79Å 118.35Å 90.00° 118.62° 90.00°	Depositor
Resolution (Å)	39.38 – 2.25 46.71 – 2.25	Depositor EDS
% Data completeness (in resolution range)	98.9 (39.38-2.25) 98.9 (46.71-2.25)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.24Å)	Xtriage
Refinement program	PHENIX 1.20.1-4487	Depositor
R, R_{free}	0.169 , 0.201 0.169 , 0.202	Depositor DCC
R_{free} test set	2068 reflections (3.49%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.018 for -h-l,k,h 0.018 for l,k,-h-l 0.025 for h,-k,-h-l 0.068 for -h-l,-k,l 0.031 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6459	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/3015	0.56	0/4100
1	B	0.38	0/2995	0.55	0/4073
All	All	0.37	0/6010	0.56	0/8173

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2923	0	2860	11	0
1	B	2912	0	2846	13	0
2	A	24	0	24	0	0
2	B	24	0	24	2	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	4	0	6	0	0
4	B	4	0	6	0	0
5	A	255	0	0	0	0
5	B	303	0	0	4	0
All	All	6459	0	5766	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:129[B]:GLN:OE1	5:B:501:HOH:O	2.09	0.70
1:A:12:LEU:HD12	1:A:13:VAL:HG23	1.76	0.67
1:B:249:ASN:OD1	5:B:502:HOH:O	2.13	0.67
1:B:45:THR:HB	2:B:402:BGC:H4	1.79	0.64
1:B:17:GLU:HA	1:B:20:ARG:HG3	1.82	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/398 (96%)	376 (98%)	6 (2%)	0	100	100
1	B	379/398 (95%)	369 (97%)	10 (3%)	0	100	100
All	All	761/796 (96%)	745 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/328 (94%)	307 (99%)	2 (1%)	86	91
1	B	307/328 (94%)	304 (99%)	3 (1%)	76	84
All	All	616/656 (94%)	611 (99%)	5 (1%)	81	88

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	SER
1	A	249	ASN
1	B	125	LEU
1	B	249	ASN
1	B	314	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	A	402	-	12,12,12	0.61	0	17,17,17	0.99	0
3	SO4	A	403	-	4,4,4	0.14	0	6,6,6	0.14	0
2	BGC	B	402	-	12,12,12	0.50	0	17,17,17	0.99	1 (5%)
4	EDO	B	404	-	3,3,3	0.58	0	2,2,2	0.28	0
2	BGC	B	401	-	12,12,12	0.48	0	17,17,17	0.85	1 (5%)
3	SO4	B	403	-	4,4,4	0.17	0	6,6,6	0.19	0
2	BGC	A	401	-	12,12,12	0.58	0	17,17,17	0.89	1 (5%)
4	EDO	A	404	-	3,3,3	0.58	0	2,2,2	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	402	-	-	0/2/22/22	0/1/1/1
2	BGC	B	402	-	-	2/2/22/22	0/1/1/1
4	EDO	B	404	-	-	0/1/1/1	-
2	BGC	B	401	-	-	0/2/22/22	0/1/1/1
2	BGC	A	401	-	-	0/2/22/22	0/1/1/1
4	EDO	A	404	-	-	0/1/1/1	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	402	BGC	C6-C5-C4	-2.95	106.08	113.00
2	B	401	BGC	C1-C2-C3	-2.23	105.70	110.31
2	A	401	BGC	C1-C2-C3	-2.14	105.88	110.31

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	402	BGC	C4-C5-C6-O6
2	B	402	BGC	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	402	BGC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	374/398 (93%)	-0.21	10 (2%) 54 57	24, 39, 66, 87	0
1	B	374/398 (93%)	-0.27	21 (5%) 24 26	19, 34, 72, 95	0
All	All	748/796 (93%)	-0.24	31 (4%) 37 40	19, 37, 69, 95	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	294	PRO	4.9
1	B	276	ALA	4.3
1	A	295	CYS	4.0
1	B	203	GLY	3.9
1	B	277	ALA	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BGC	B	402	12/12	0.69	0.28	46,63,76,78	0
2	BGC	A	402	12/12	0.83	0.20	45,62,71,80	0
4	EDO	B	404	4/4	0.94	0.11	40,42,45,50	0
4	EDO	A	404	4/4	0.96	0.13	40,44,46,48	0
2	BGC	A	401	12/12	0.97	0.16	22,26,30,34	0
3	SO4	B	403	5/5	0.97	0.11	55,57,62,63	0
2	BGC	B	401	12/12	0.98	0.14	23,24,27,28	0
3	SO4	A	403	5/5	0.99	0.07	54,55,59,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around BGC A 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BGC A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.