



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2021 – 08:55 PM EST

PDB ID : 1DUW
Title : STRUCTURE OF NONAHEME CYTOCHROME C
Authors : Umhau, S.; Fritz, G.; Diederichs, K.; Breed, J.; Kroneck, P.M.; Welte, W.
Deposited on : 2000-01-19
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.17.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.17.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NONAHEME CYTOCHROME C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	289	2188	1334	415	411	28	0	1	0

- Molecule 2 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0
2	A	1	43	34	1	4	4	0	0
2	A	1	43	34	1	4	4	0	0
2	A	1	43	34	1	4	4	0	0
2	A	1	43	34	1	4	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	423	Total	O	0	0
			423	423		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NONAHEME CYTOCHROME C

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	55.40Å 55.40Å 236.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.89 29.60 – 1.88	Depositor EDS
% Data completeness (in resolution range)	7.5 (30.00-1.89) 97.9 (29.60-1.88)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 1.88Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.183 , 0.235 0.182 , 0.229	Depositor DCC
R_{free} test set	2143 reflections (6.96%)	wwPDB-VP
Wilson B-factor (Å ²)	25.1	Xtrriage
Anisotropy	0.343	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 100.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3004	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.84	1/2246 (0.0%)	1.13	6/3044 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	292	PRO	C-OXT	85.46	2.85	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	207	ARG	NE-CZ-NH1	-9.93	115.33	120.30
1	A	207	ARG	NE-CZ-NH2	7.95	124.28	120.30
1	A	254	PRO	CA-N-CD	-6.84	101.92	111.50
1	A	115	HIS	CA-CB-CG	-5.71	103.90	113.60
1	A	207	ARG	CD-NE-CZ	5.56	131.39	123.60
1	A	250	ASP	CA-CB-CG	5.24	124.94	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2188	0	2124	54	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	387	0	270	20	0
3	A	6	0	8	0	0
4	A	423	0	0	13	0
All	All	3004	0	2402	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:254:PRO:HD2	1:A:256:ARG:H	1.39	0.87
1:A:173:THR:H	1:A:176:GLN:HE21	1.25	0.83
1:A:138:THR:H	1:A:141:GLN:HE21	1.29	0.77
1:A:56:PRO:HG3	4:A:1279:HOH:O	1.85	0.76
1:A:169:VAL:HG21	1:A:219:ASN:O	1.86	0.75
1:A:173:THR:H	1:A:176:GLN:NE2	1.86	0.74
1:A:254:PRO:HD3	4:A:1407:HOH:O	1.87	0.73
1:A:277:GLY:O	1:A:280:LYS:HE3	1.89	0.73
1:A:255:ASN:HA	4:A:1336:HOH:O	1.91	0.69
1:A:291:ARG:HD2	4:A:1045:HOH:O	1.93	0.68
1:A:22:LYS:HG3	1:A:66:GLU:OE2	1.93	0.67
1:A:68:LYS:HG3	1:A:70:GLU:OE1	1.99	0.62
1:A:89:ALA:HB2	4:A:1232:HOH:O	2.01	0.61
2:A:297:HEC:HMC1	2:A:297:HEC:HBC3	1.82	0.60
2:A:301:HEC:HMC1	2:A:301:HEC:HBC3	1.87	0.56
2:A:298:HEC:HMC1	2:A:298:HEC:HBC3	1.85	0.56
1:A:140:GLU:HB2	4:A:1355:HOH:O	2.05	0.56
2:A:300:HEC:HMC1	2:A:300:HEC:HBC3	1.88	0.56
2:A:299:HEC:HMC1	2:A:299:HEC:HBC3	1.88	0.55
1:A:246:THR:O	1:A:258:ASN:HB2	2.10	0.52
1:A:209:LYS:NZ	1:A:209:LYS:HB3	2.25	0.52
1:A:171:PRO:HA	1:A:221:PRO:HA	1.92	0.51
1:A:188:ASP:O	1:A:291:ARG:HD3	2.11	0.51
2:A:297:HEC:HMB1	2:A:297:HEC:HBB3	1.93	0.51
1:A:140:GLU:HA	1:A:140:GLU:OE1	2.10	0.51
2:A:295:HEC:HMC1	2:A:295:HEC:HBC3	1.92	0.51
1:A:8:SER:HB2	2:A:296:HEC:C2B	2.41	0.50
1:A:138:THR:H	1:A:141:GLN:NE2	2.05	0.50
1:A:291:ARG:O	1:A:292:PRO:O	2.30	0.49
1:A:254:PRO:HD2	1:A:256:ARG:HB2	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:LEU:HD13	2:A:297:HEC:CHA	2.43	0.49
1:A:207:ARG:HG3	4:A:1322:HOH:O	2.12	0.48
1:A:241:CYS:HA	2:A:298:HEC:HHC	1.95	0.48
1:A:35:PHE:CZ	2:A:296:HEC:HHD	2.49	0.48
1:A:190:TYR:CD2	1:A:290:ALA:HA	2.49	0.47
1:A:30:MET:HG3	2:A:294:HEC:C4A	2.45	0.47
1:A:278:ARG:HG2	1:A:279:PRO:N	2.26	0.47
1:A:187:ALA:HB3	4:A:1176:HOH:O	2.15	0.47
1:A:63:HIS:CE1	1:A:74:VAL:HB	2.50	0.46
1:A:207:ARG:NH2	4:A:1322:HOH:O	2.48	0.46
2:A:301:HEC:HMB1	2:A:301:HEC:HBB3	1.97	0.46
1:A:122:ARG:HD3	1:A:126:TRP:CH2	2.51	0.46
1:A:254:PRO:CD	1:A:256:ARG:HB2	2.46	0.45
1:A:5:PRO:HA	4:A:1301:HOH:O	2.16	0.45
1:A:114:CYS:O	1:A:118:VAL:HG23	2.17	0.45
1:A:138:THR:OG1	1:A:141:GLN:HG3	2.16	0.45
1:A:84:ASN:O	1:A:84:ASN:ND2	2.50	0.45
2:A:299:HEC:HMB1	2:A:299:HEC:HBB3	1.97	0.45
1:A:188:ASP:O	1:A:291:ARG:NH1	2.49	0.44
1:A:214:ALA:HB1	1:A:218:HIS:CE1	2.53	0.44
1:A:212:LYS:HD3	4:A:1058:HOH:O	2.17	0.44
1:A:241:CYS:HA	2:A:298:HEC:CHC	2.47	0.44
2:A:296:HEC:HMC1	2:A:296:HEC:HBC3	1.99	0.44
1:A:100:CYS:O	1:A:103:GLN:HB2	2.18	0.43
2:A:293:HEC:HBB3	2:A:293:HEC:HMB1	2.00	0.43
1:A:156:GLU:HA	4:A:1152:HOH:O	2.18	0.43
1:A:252:ALA:O	1:A:254:PRO:HA	2.18	0.43
1:A:248:GLU:HA	1:A:248:GLU:OE1	2.19	0.43
1:A:208:ILE:HD13	2:A:295:HEC:HMC2	2.01	0.43
1:A:55:ASP:HA	1:A:56:PRO:HD3	1.90	0.42
1:A:137:MET:HA	1:A:141:GLN:NE2	2.35	0.41
1:A:222:GLU:HB2	1:A:234:LEU:HD13	2.03	0.41
1:A:97:CYS:HA	2:A:297:HEC:CHC	2.51	0.41
1:A:280:LYS:HA	1:A:280:LYS:HD3	1.91	0.40
2:A:294:HEC:HBC3	2:A:294:HEC:HMC1	2.03	0.40
1:A:248:GLU:HB2	4:A:1142:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	288/292 (99%)	279 (97%)	8 (3%)	1 (0%)	41 31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	254	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	242/242 (100%)	229 (95%)	13 (5%)	22 13

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	SER
1	A	55	ASP
1	A	72	ASN
1	A	84	ASN
1	A	121	LYS
1	A	149	LYS
1	A	167	LYS
1	A	170	GLN
1	A	194	ASN
1	A	219	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	253	ASN
1	A	278	ARG
1	A	291	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	103	GLN
1	A	141	GLN
1	A	154	GLN
1	A	176	GLN
1	A	194	ASN
1	A	219	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEC	A	301	1	26,50,50	1.30	2 (7%)	18,82,82	2.72	7 (38%)
2	HEC	A	297	1	26,50,50	1.21	3 (11%)	18,82,82	2.10	7 (38%)
2	HEC	A	300	1	26,50,50	1.25	2 (7%)	18,82,82	1.88	7 (38%)
3	GOL	A	302	-	5,5,5	0.77	0	5,5,5	0.67	0
2	HEC	A	298	1	26,50,50	1.17	2 (7%)	18,82,82	2.03	7 (38%)
2	HEC	A	294	1	26,50,50	1.25	3 (11%)	18,82,82	2.36	5 (27%)
2	HEC	A	293	1	26,50,50	1.16	2 (7%)	18,82,82	1.77	5 (27%)
2	HEC	A	296	1	26,50,50	1.10	2 (7%)	18,82,82	1.91	6 (33%)
2	HEC	A	299	1	26,50,50	1.19	2 (7%)	18,82,82	1.70	4 (22%)
2	HEC	A	295	1	26,50,50	1.23	2 (7%)	18,82,82	2.18	4 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	301	1	-	1/6/54/54	-
2	HEC	A	297	1	-	1/6/54/54	-
2	HEC	A	300	1	-	0/6/54/54	-
3	GOL	A	302	-	-	0/4/4/4	-
2	HEC	A	298	1	-	0/6/54/54	-
2	HEC	A	294	1	-	0/6/54/54	-
2	HEC	A	293	1	-	0/6/54/54	-
2	HEC	A	296	1	-	0/6/54/54	-
2	HEC	A	299	1	-	0/6/54/54	-
2	HEC	A	295	1	-	0/6/54/54	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	HEC	C3C-C2C	-4.19	1.36	1.40
2	A	301	HEC	C3B-C2B	-3.95	1.36	1.40
2	A	295	HEC	C3B-C2B	-3.66	1.36	1.40
2	A	293	HEC	C3C-C2C	-3.48	1.37	1.40
2	A	294	HEC	C3B-C2B	-3.29	1.37	1.40
2	A	297	HEC	C3C-C2C	-3.27	1.37	1.40
2	A	298	HEC	C3C-C2C	-3.25	1.37	1.40
2	A	301	HEC	C3C-C2C	-3.24	1.37	1.40
2	A	294	HEC	C3C-C2C	-3.21	1.37	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	295	HEC	C3C-C2C	-3.16	1.37	1.40
2	A	298	HEC	C3B-C2B	-3.14	1.37	1.40
2	A	297	HEC	C3B-C2B	-3.04	1.37	1.40
2	A	293	HEC	C3B-C2B	-3.01	1.37	1.40
2	A	299	HEC	C3B-C2B	-2.90	1.37	1.40
2	A	299	HEC	C3C-C2C	-2.88	1.37	1.40
2	A	300	HEC	C3B-C2B	-2.75	1.37	1.40
2	A	296	HEC	C3B-C2B	-2.68	1.38	1.40
2	A	294	HEC	C1D-ND	2.08	1.40	1.36
2	A	296	HEC	C3C-C2C	-2.01	1.38	1.40
2	A	297	HEC	C3C-C4C	2.00	1.46	1.43

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	HEC	CMC-C2C-C1C	-6.01	119.22	128.46
2	A	294	HEC	CMC-C2C-C1C	-5.88	119.43	128.46
2	A	301	HEC	CMC-C2C-C3C	5.47	132.26	125.82
2	A	301	HEC	CMD-C2D-C1D	-5.30	120.31	128.46
2	A	295	HEC	CMC-C2C-C1C	-4.85	121.01	128.46
2	A	297	HEC	CMC-C2C-C1C	-4.61	121.38	128.46
2	A	294	HEC	CMB-C2B-C1B	-4.42	121.67	128.46
2	A	296	HEC	CMD-C2D-C1D	-4.35	121.78	128.46
2	A	298	HEC	CMB-C2B-C1B	-4.25	121.92	128.46
2	A	295	HEC	CMB-C2B-C1B	-4.13	122.11	128.46
2	A	295	HEC	CMC-C2C-C3C	3.92	130.43	125.82
2	A	294	HEC	CMC-C2C-C3C	3.85	130.35	125.82
2	A	300	HEC	CMC-C2C-C1C	-3.78	122.65	128.46
2	A	293	HEC	CMD-C2D-C1D	-3.67	122.82	128.46
2	A	298	HEC	CMB-C2B-C3B	3.49	129.93	125.82
2	A	299	HEC	CMB-C2B-C1B	-3.48	123.11	128.46
2	A	297	HEC	CMB-C2B-C1B	-3.42	123.21	128.46
2	A	300	HEC	CMB-C2B-C1B	-3.39	123.26	128.46
2	A	301	HEC	CMB-C2B-C1B	-3.37	123.28	128.46
2	A	298	HEC	CMD-C2D-C1D	-3.35	123.31	128.46
2	A	294	HEC	CMB-C2B-C3B	3.33	129.74	125.82
2	A	299	HEC	CMC-C2C-C1C	-3.33	123.35	128.46
2	A	295	HEC	CMB-C2B-C3B	3.32	129.72	125.82
2	A	298	HEC	CMC-C2C-C1C	-3.26	123.46	128.46
2	A	293	HEC	CMC-C2C-C1C	-3.13	123.66	128.46
2	A	297	HEC	CMD-C2D-C1D	-3.13	123.66	128.46
2	A	296	HEC	CMD-C2D-C3D	3.10	130.78	124.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	293	HEC	CMB-C2B-C1B	-3.04	123.80	128.46
2	A	297	HEC	CMC-C2C-C3C	2.92	129.25	125.82
2	A	296	HEC	CMB-C2B-C1B	-2.87	124.06	128.46
2	A	297	HEC	CMB-C2B-C3B	2.85	129.17	125.82
2	A	301	HEC	CMB-C2B-C3B	2.79	129.10	125.82
2	A	299	HEC	CBA-CAA-C2A	2.78	117.60	112.48
2	A	293	HEC	CMD-C2D-C3D	2.61	129.87	124.94
2	A	301	HEC	CMD-C2D-C3D	2.58	129.80	124.94
2	A	294	HEC	CMD-C2D-C1D	-2.41	124.75	128.46
2	A	300	HEC	CMC-C2C-C3C	2.39	128.63	125.82
2	A	300	HEC	CMD-C2D-C1D	-2.38	124.81	128.46
2	A	297	HEC	CMD-C2D-C3D	2.35	129.38	124.94
2	A	300	HEC	CMB-C2B-C3B	2.33	128.56	125.82
2	A	299	HEC	CMC-C2C-C3C	2.33	128.56	125.82
2	A	296	HEC	C3C-C4C-NC	-2.32	106.57	110.94
2	A	301	HEC	CAA-CBA-CGA	2.30	116.53	112.67
2	A	293	HEC	CMA-C3A-C2A	2.29	129.26	124.94
2	A	297	HEC	CAA-CBA-CGA	2.27	116.48	112.67
2	A	300	HEC	C3C-C4C-NC	-2.22	106.75	110.94
2	A	296	HEC	CBA-CAA-C2A	-2.22	108.39	112.48
2	A	300	HEC	CMA-C3A-C2A	2.21	129.11	124.94
2	A	298	HEC	CMC-C2C-C3C	2.16	128.36	125.82
2	A	298	HEC	CMD-C2D-C3D	2.09	128.89	124.94
2	A	298	HEC	C3C-C4C-NC	-2.02	107.13	110.94
2	A	296	HEC	CMA-C3A-C2A	2.01	128.73	124.94

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	HEC	C2A-CAA-CBA-CGA
2	A	297	HEC	C2A-CAA-CBA-CGA

There are no ring outliers.

9 monomers are involved in 20 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	HEC	2	0
2	A	297	HEC	4	0
2	A	300	HEC	1	0
2	A	298	HEC	3	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	294	HEC	2	0
2	A	293	HEC	1	0
2	A	296	HEC	3	0
2	A	299	HEC	2	0
2	A	295	HEC	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/292 (98%)	0.22	19 (6%) 18 20	18, 27, 53, 93	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	292	PRO	6.1
1	A	253	ASN	4.9
1	A	254	PRO	4.3
1	A	84	ASN	4.0
1	A	291	ARG	4.0
1	A	290	ALA	3.7
1	A	169	VAL	3.5
1	A	170	GLN	3.3
1	A	168	PRO	3.2
1	A	251	PRO	2.8
1	A	250	ASP	2.7
1	A	174	ALA	2.4
1	A	171	PRO	2.4
1	A	72	ASN	2.3
1	A	255	ASN	2.2
1	A	175	MET	2.1
1	A	252	ALA	2.1
1	A	73	PHE	2.1
1	A	152	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	302	6/6	0.93	0.14	28,38,42,46	0
2	HEC	A	296	43/43	0.96	0.12	17,27,33,36	0
2	HEC	A	297	43/43	0.96	0.14	18,22,26,31	0
2	HEC	A	298	43/43	0.96	0.11	17,22,32,51	0
2	HEC	A	293	43/43	0.96	0.11	18,23,37,46	0
2	HEC	A	294	43/43	0.98	0.11	16,22,24,26	0
2	HEC	A	299	43/43	0.98	0.09	14,25,29,30	0
2	HEC	A	300	43/43	0.98	0.10	15,20,29,36	0
2	HEC	A	301	43/43	0.98	0.11	16,21,28,33	0
2	HEC	A	295	43/43	0.98	0.09	16,21,36,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEC A 297:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 298:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC A 293:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around HEC A 299:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 300:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.