

Full wwPDB X-ray Structure Validation Report (i)

Sep 6, 2023 – 10:51 AM EDT

PDB ID : 4DUC

Title : cytochrome P450 BM3h-2G9 MRI sensor, no ligand

Authors: Brustad, E.M.; Lelyveld, V.S.; Snow, C.D.; Crook, N.; Martinez, F.M.; Scholl,

T.J.; Jasanoff, A.; Arnold, F.H.

Deposited on : 2012-02-21

Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	472	86%	11% •
1	В	472	83%	12% • •



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called cytochrome P450 BM3 variant 2G9.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total 3637	C 2326	N 614	O 679	S 18	0	1	0
1	В	456	Total 3633	C 2327	N 614	O 674	S 18	0	3	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	CYS	ARG	engineered mutation	UNP P14779
A	87	LEU	PHE	engineered mutation	UNP P14779
A	268	SER	THR engineered mutation		UNP P14779
A	438	LEU	THR	engineered mutation	UNP P14779
A	464	LEU	-	expression tag	UNP P14779
A	465	GLU	-	expression tag	UNP P14779
A	466	HIS	-	expression tag	UNP P14779
A	467	HIS	-	expression tag	UNP P14779
A	468	HIS	-	expression tag	UNP P14779
A	469	HIS	-	expression tag	UNP P14779
A	470	HIS	-	expression tag	UNP P14779
A	471	HIS	-	expression tag	UNP P14779
В	50	CYS	ARG	engineered mutation	UNP P14779
В	87	LEU	PHE	engineered mutation	UNP P14779
В	268	SER	THR	engineered mutation	UNP P14779
В	438	LEU	THR	engineered mutation	UNP P14779
В	464	LEU	-	expression tag	UNP P14779
В	465	GLU	-	expression tag	UNP P14779
В	466	HIS	-	expression tag	UNP P14779
В	467	HIS	-	expression tag	UNP P14779
В	468	HIS	-	expression tag	UNP P14779
В	469	HIS	- expression tag		UNP P14779
В	470	HIS	-	expression tag	UNP P14779
В	471	HIS	-	expression tag	UNP P14779



 \bullet Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $\rm C_{34}H_{32}FeN_4O_4).$



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	Λ	1	Total	С	Fe	N	О	0	0	
	$\begin{array}{ c c c c } Z & A & A \\ \end{array}$	1	43	34	1	4	4	0		
2	D	1	Total	С	Fe	N	О	0	0	
	1	43	34	1	4	4	0	U		

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	284	Total O 284 284	0	0
3	В	284	Total O 284 284	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: cytochrome P450 BM3 variant 2G9



• Molecule 1: cytochrome P450 BM3 variant 2G9





4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	58.66Å 153.75Å 60.98Å	Depositor	
a, b, c, α , β , γ	90.00° 94.82° 90.00°	Depositor	
Resolution (Å)	29.22 - 1.92	Depositor	
rtesolution (A)	29.22 - 1.92	EDS	
% Data completeness	95.2 (29.22-1.92)	Depositor	
(in resolution range)	95.3 (29.22-1.92)	EDS	
R_{merge}	(Not available)	Depositor	
R_{sym}	0.09	Depositor	
$< I/\sigma(I) > 1$	1.31 (at 1.92Å)	Xtriage	
Refinement program	REFMAC 5.5.0109	Depositor	
Ρ. Р.	0.185 , 0.226	Depositor	
R, R_{free}	0.189 , 0.231	DCC	
R_{free} test set	3864 reflections (4.99%)	wwPDB-VP	
Wilson B-factor (Å ²)	22.2	Xtriage	
Anisotropy	0.129	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36 , 49.5	EDS	
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtriage	
Estimated twinning fraction	0.026 for l,-k,h	Xtriage	
F_o, F_c correlation	0.95	EDS	
Total number of atoms	7924	wwPDB-VP	
Average B, all atoms (Å ²)	27.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	1.03	1/3724 (0.0%)	0.93	9/5041 (0.2%)	
1	В	1.01	$2/3726 \ (0.1\%)$	0.95	14/5045 (0.3%)	
All	All	1.02	3/7450 (0.0%)	0.94	23/10086 (0.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}({ ext{\AA}})$
1	В	107	PHE	CE2-CZ	6.31	1.49	1.37
1	В	393	PHE	CE1-CZ	5.36	1.47	1.37
1	A	310	GLN	CG-CD	5.09	1.62	1.51

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	161	ARG	NE-CZ-NH2	-6.98	116.81	120.30
1	В	323	ARG	NE-CZ-NH2	-6.71	116.95	120.30
1	A	323	ARG	NE-CZ-NH2	-6.61	116.99	120.30
1	A	318	LEU	CB-CG-CD1	-6.57	99.83	111.00
1	В	232	ASP	CB-CG-OD1	6.41	124.07	118.30
1	В	296[A]	ARG	NE-CZ-NH1	6.39	123.50	120.30
1	В	296[B]	ARG	NE-CZ-NH1	6.39	123.50	120.30
1	В	161	ARG	NE-CZ-NH1	6.12	123.36	120.30



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	323	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	A	136	ASP	CB-CG-OD1	5.87	123.58	118.30
1	В	56	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	В	126	LEU	CA-CB-CG	-5.54	102.56	115.30
1	A	36	LEU	CB-CG-CD1	-5.46	101.71	111.00
1	В	296[A]	ARG	NE-CZ-NH2	-5.43	117.59	120.30
1	В	296[B]	ARG	NE-CZ-NH2	-5.43	117.59	120.30
1	A	68	ASP	CB-CG-OD1	5.35	123.11	118.30
1	В	455	LEU	CB-CG-CD2	-5.30	101.98	111.00
1	A	369	ASP	CB-CG-OD1	5.30	123.07	118.30
1	A	437	LEU	CA-CB-CG	5.23	127.34	115.30
1	В	121	ASP	CB-CG-OD1	5.19	122.97	118.30
1	В	436	THR	N-CA-C	-5.02	97.45	111.00
1	A	161	ARG	NE-CZ-NH2	-5.01	117.79	120.30
1	В	432	ASP	CB-CG-OD1	5.00	122.80	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	455	LEU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3637	0	3571	34	0
1	В	3633	0	3566	30	0
2	A	43	0	30	0	0
2	В	43	0	30	1	0
3	A	284	0	0	3	1
3	В	284	0	0	5	1
All	All	7924	0	7197	64	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

A	A. 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)
1:B:204:GLN:HE21	1:B:204:GLN:HA	1.30	0.95
1:A:185:MET:HE1	1:A:437:LEU:HD23	1.58	0.84
1:B:204:GLN:HA	1:B:204:GLN:NE2	1.98	0.77
1:A:267:GLU:HB3	1:A:438:LEU:HD13	1.71	0.73
1:A:370:ASP:OD2	1:A:375:ARG:NH1	2.22	0.72
1:A:229:GLN:HE22	1:A:239:ASN:HD21	1.35	0.71
1:A:135:ALA:O	1:A:136:ASP:CB	2.35	0.71
1:A:185:MET:CE	1:A:437:LEU:HD23	2.22	0.69
1:A:38:GLU:HB2	1:A:54:SER:HB3	1.76	0.68
1:A:185:MET:HE1	1:A:437:LEU:CD2	2.26	0.63
1:A:135:ALA:O	1:A:136:ASP:HB3	1.98	0.63
1:B:187:LYS:NZ	3:B:867:HOH:O	2.25	0.62
1:A:38:GLU:CB	1:A:54:SER:HB3	2.29	0.61
1:A:185:MET:SD	1:A:437:LEU:HD23	2.39	0.61
1:B:104:LEU:N	1:B:105:PRO:HD2	2.16	0.60
1:B:43:GLU:HG2	1:B:48:VAL:HG22	1.83	0.60
1:B:267:GLU:OE2	1:B:440:LYS:NZ	2.32	0.59
1:A:185:MET:SD	1:A:437:LEU:CD2	2.92	0.58
1:A:109:GLN:HE22	1:A:309:LYS:NZ	2.04	0.56
1:A:135:ALA:O	1:A:136:ASP:OD1	2.24	0.56
1:B:187:LYS:CE	3:B:867:HOH:O	2.53	0.56
1:B:409[A]:GLU:OE2	3:B:835:HOH:O	2.19	0.53
1:B:8:PRO:HG3	1:B:32:ILE:HG21	1.89	0.53
1:B:370:ASP:OD2	1:B:375:ARG:NH2	2.41	0.53
1:A:200:GLU:HA	1:A:200:GLU:OE1	2.07	0.53
1:B:301:PRO:HB2	1:B:456:GLY:HA3	1.91	0.53
1:B:267:GLU:HB3	1:B:438:LEU:HD12	1.90	0.52
1:A:387:GLN:NE2	3:A:867:HOH:O	2.18	0.51
1:A:301:PRO:HB2	1:A:456:GLY:HA3	1.92	0.50
1:A:38:GLU:HB2	1:A:54:SER:CB	2.41	0.50
1:A:177:MET:SD	1:A:263:ILE:HD13	2.52	0.49
1:A:253:ASN:O	1:A:257:GLN:HG2	2.12	0.49
1:B:207:GLU:O	1:B:211:VAL:HG23	2.14	0.47
1:B:135:ALA:O	1:B:136:ASP:HB2	2.15	0.47
1:B:97:LYS:HB2	3:B:866:HOH:O	2.14	0.47
1:B:81:PHE:HB3	1:B:209:ILE:HG12	1.96	0.46
1:B:60:GLU:OE2	1:B:342:GLY:HA2	2.15	0.46
1:B:124:VAL:O	1:B:128:GLN:HG3	2.16	0.46
1:B:33:ALA:CB	1:B:359:GLN:HG2	2.46	0.46
1:B:377:GLU:O	1:B:380:GLU:HB2	2.17	0.45
1:B:357:ILE:HG22	1:B:361:HIS:CE1	2.52	0.45



Continued from previous page...

A	A	Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)	
1:B:267:GLU:HB3	1:B:438:LEU:CD1	2.47	0.44	
1:B:109:GLN:HE22	1:B:309:LYS:HZ2	1.64	0.44	
1:A:287:LEU:O	1:A:287:LEU:HD23	2.16	0.44	
1:A:81:PHE:HB3	1:A:209:ILE:HG12	1.99	0.44	
1:B:109:GLN:HE22	1:B:309:LYS:NZ	2.16	0.44	
1:B:215:LEU:HG	1:B:219:ILE:HD12	1.99	0.43	
1:A:453:ILE:HA	1:A:454:PRO:HD2	1.67	0.43	
1:A:17:LEU:N	1:A:18:PRO:CD	2.82	0.43	
1:A:185:MET:CE	1:A:437:LEU:CD2	2.94	0.43	
1:B:367:TRP:HB2	1:B:371:VAL:HG12	2.00	0.42	
1:A:135:ALA:O	1:A:136:ASP:CG	2.57	0.42	
1:A:336:LYS:HG2	3:A:794:HOH:O	2.19	0.42	
1:A:296[A]:ARG:NH2	3:A:626:HOH:O	2.52	0.42	
1:B:296[A]:ARG:HD3	3:B:836:HOH:O	2.19	0.42	
1:A:173:PHE:C	1:A:173:PHE:CD2	2.93	0.42	
1:A:38:GLU:HB3	1:A:54:SER:HB3	2.02	0.42	
1:B:39:ILE:HG13	1:B:52:LEU:HD23	2.01	0.42	
1:A:60:GLU:OE2	1:A:342:GLY:HA2	2.19	0.41	
1:B:400:CYS:HA	2:B:500:HEM:CHA	2.50	0.41	
1:A:24:LYS:HE2	1:A:433:ILE:O	2.20	0.41	
1:A:104:LEU:HB3	1:A:105:PRO:HD3	2.02	0.41	
1:B:298:LEU:HD22	1:B:303:PRO:HB3	2.02	0.41	
1:A:46:GLY:O	1:A:47:ARG:HB3	2.21	0.41	

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1 Atom-2		$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)	
3:A:861:HOH:O	3:B:846:HOH:O[2_656]	1.88	0.32	

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	455/472~(96%)	441 (97%)	14 (3%)	0	100	100
1	В	457/472 (97%)	446 (98%)	10 (2%)	1 (0%)	47	38
All	All	912/944 (97%)	887 (97%)	24 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type	
1	В	191	ALA	

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	388/414 (94%)	381 (98%)	7 (2%)	59 53
1	В	385/414 (93%)	377 (98%)	8 (2%)	53 46
All	All	773/828 (93%)	758 (98%)	15 (2%)	55 51

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	50	CYS
1	A	136	ASP
1	A	148	LEU
1	A	230	SER
1	A	409	GLU
1	A	437	LEU
1	В	23	ASP
1	В	35	GLU
1	В	126	LEU
1	В	148	LEU
1	В	200	GLU
1	В	204	GLN
1	В	430	GLU
1	В	440	LYS



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	109	GLN
1	A	229	GLN
1	A	404	GLN
1	В	109	GLN
1	В	229	GLN
1	В	239	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type C		Chain Res		Link	Bond lengths			Bond angles		
Mol Type Chai	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	HEM	A	500	1,3	41,50,50	2.03	12 (29%)	45,82,82	1.91	13 (28%)
2	HEM	В	500	1,3	41,50,50	1.99	11 (26%)	45,82,82	1.89	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1,3	-	5/12/54/54	-
2	HEM	В	500	1,3	-	2/12/54/54	-

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\text{\AA})$
2	В	500	HEM	C3D-C2D	6.59	1.50	1.36
2	A	500	HEM	C3D-C2D	6.30	1.50	1.36
2	В	500	HEM	CMA-C3A	4.94	1.61	1.51
2	A	500	HEM	CAA-C2A	4.49	1.58	1.52
2	A	500	HEM	C3C-CAC	3.72	1.55	1.47
2	В	500	HEM	FE-ND	3.38	2.13	1.96
2	A	500	HEM	CMB-C2B	3.37	1.57	1.50
2	A	500	HEM	C3C-C2C	-3.22	1.35	1.40
2	A	500	HEM	FE-ND	3.16	2.12	1.96
2	В	500	HEM	C3C-C2C	-3.11	1.36	1.40
2	В	500	HEM	CMB-C2B	2.64	1.56	1.50
2	A	500	HEM	CAB-C3B	2.61	1.54	1.47
2	A	500	HEM	CMA-C3A	2.59	1.57	1.51
2	В	500	HEM	CBD-CGD	2.49	1.56	1.50
2	A	500	HEM	CMC-C2C	2.32	1.57	1.51
2	В	500	HEM	CAA-C2A	2.31	1.55	1.52
2	A	500	HEM	FE-NB	2.28	2.08	1.96
2	В	500	HEM	C3C-CAC	2.24	1.52	1.47
2	A	500	HEM	CBD-CGD	2.22	1.55	1.50
2	В	500	HEM	C3B-C2B	-2.20	1.32	1.37
2	В	500	HEM	C1A-NA	2.10	1.40	1.36
2	В	500	HEM	FE-NB	2.09	2.07	1.96
2	A	500	HEM	CMD-C2D	2.01	1.55	1.50

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
2	В	500	HEM	C4D-ND-C1D	6.63	111.92	105.07
2	A	500	HEM	C4D-ND-C1D	5.31	110.55	105.07
2	В	500	HEM	C4C-CHD-C1D	4.14	128.03	122.56
2	В	500	HEM	C4B-C3B-C2B	3.77	110.11	107.11
2	A	500	HEM	CBD-CAD-C3D	-3.67	102.43	112.63
2	В	500	HEM	CBD-CAD-C3D	-3.54	102.79	112.63
2	A	500	HEM	CAD-CBD-CGD	-3.54	105.99	113.60
2	A	500	HEM	C1B-NB-C4B	3.30	108.48	105.07



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\mathrm{Ideal}(^{o})$
2	A	500	HEM	C1D-C2D-C3D	-3.28	103.51	106.96
2	A	500	HEM	O2D-CGD-CBD	3.01	123.70	114.03
2	В	500	HEM	CAD-CBD-CGD	-3.00	107.15	113.60
2	A	500	HEM	C4C-CHD-C1D	2.75	126.18	122.56
2	A	500	HEM	C4B-CHC-C1C	2.71	126.13	122.56
2	В	500	HEM	CMC-C2C-C3C	2.55	129.45	124.68
2	A	500	HEM	O2A-CGA-CBA	2.44	121.88	114.03
2	В	500	HEM	C2C-C3C-C4C	2.42	108.58	106.90
2	A	500	HEM	O1D-CGD-CBD	-2.39	115.41	123.08
2	A	500	HEM	C4B-C3B-C2B	2.37	109.00	107.11
2	В	500	HEM	CAD-C3D-C4D	2.32	128.72	124.66
2	A	500	HEM	CHC-C4B-NB	2.30	126.93	124.43
2	A	500	HEM	C2C-C3C-C4C	2.26	108.47	106.90
2	В	500	HEM	C1B-NB-C4B	2.09	107.23	105.07

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	HEM	C2B-C3B-CAB-CBB
2	A	500	HEM	C4B-C3B-CAB-CBB
2	В	500	HEM	CAD-CBD-CGD-O1D
2	В	500	HEM	CAD-CBD-CGD-O2D
2	A	500	HEM	CAD-CBD-CGD-O2D
2	A	500	HEM	CAA-CBA-CGA-O2A
2	A	500	HEM	CAD-CBD-CGD-O1D

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	500	HEM	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and



any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	456/472 (96%)	0.00	12 (2%) 56 59	12, 25, 43, 59	0
1	В	$456/472 \ (96\%)$	0.14	20 (4%) 34 37	12, 25, 49, 78	0
All	All	912/944 (96%)	0.07	32 (3%) 44 47	12, 25, 46, 78	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	191	ALA	6.3
1	A	191	ALA	6.2
1	В	197	ALA	6.0
1	В	2	ILE	5.5
1	В	225	ALA	4.3
1	A	196	PRO	4.2
1	В	198	TYR	3.7
1	A	197	ALA	3.6
1	В	201	ASN	3.5
1	В	200	GLU	3.3
1	В	203	ARG	3.0
1	В	192	ASN	3.0
1	A	195	ASP	2.9
1	В	193	PRO	2.8
1	A	149	THR	2.8
1	A	198	TYR	2.8
1	В	227	GLY	2.7
1	В	136	ASP	2.6
1	A	2	ILE	2.6
1	A	231	ASP	2.5
1	В	199	ASP	2.4
1	A	457	GLY	2.4
1	A	192	ASN	2.3
1	В	369	ASP	2.2



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	228	GLU	2.2
1	В	384	ALA	2.2
1	В	3	LYS	2.1
1	A	269	THR	2.1
1	В	457	GLY	2.1
1	В	229	GLN	2.1
1	A	203	ARG	2.0
1	В	194	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	HEM	A	500	43/43	0.98	0.12	5,14,18,22	0
2	HEM	В	500	43/43	0.99	0.12	7,14,18,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

