



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 9, 2020 – 02:56 PM BST

PDB ID : 6DVH
Title : Lactate Monooxygenase from Mycobacterium smegmatis - C203A mutant
Authors : Kean, K.M.; Karplus, P.A.
Deposited on : 2018-06-23
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.3.dev2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.3.dev2


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	5% (Poor fit) 91% (0 outliers) 8% (1 outlier) •
1	B	394	13% (Poor fit) 89% (0 outliers) 9% (1 outlier) ••
1	C	394	11% (Poor fit) 92% (0 outliers) 6% (1 outlier) •
1	D	394	22% (Poor fit) 86% (0 outliers) 13% (1 outlier) •
1	E	394	61% (Poor fit) 85% (0 outliers) 13% (1 outlier) •
1	F	394	72% (Poor fit) 85% (0 outliers) 13% (1 outlier) •

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 20104 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactate 2-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	394	3080	1964	533	571	12	0	12	0
1	B	394	3166	2020	548	586	12	0	22	0
1	C	394	3066	1949	540	565	12	0	7	0
1	D	394	3062	1949	535	566	12	0	7	0
1	E	394	3043	1935	530	566	12	0	6	0
1	F	393	3037	1933	529	564	11	0	6	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	203	ALA	CYS	engineered mutation	UNP P21795
B	203	ALA	CYS	engineered mutation	UNP P21795
C	203	ALA	CYS	engineered mutation	UNP P21795
D	203	ALA	CYS	engineered mutation	UNP P21795
E	203	ALA	CYS	engineered mutation	UNP P21795
F	203	ALA	CYS	engineered mutation	UNP P21795

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	C	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	D	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	E	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	F	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	1
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	1
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	1
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	499	Total	O	0	4
			499	499		
4	B	358	Total	O	0	4
			358	358		
4	C	313	Total	O	0	2
			313	313		
4	D	182	Total	O	0	2
			182	182		
4	E	13	Total	O	0	0
			13	13		
4	F	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lactate 2-monooxygenase




- Molecule 1: Lactate 2-monooxygenase




- Molecule 1: Lactate 2-monooxygenase




- Molecule 1: Lactate 2-monooxygenase




• Molecule 1: Lactate 2-monooxygenase



• Molecule 1: Lactate 2-monooxygenase





4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	149.62Å 149.62Å 274.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.90 – 1.70 30.90 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.90-1.70) 100.0 (30.90-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 1.70Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.169 , 0.194 0.176 , 0.199	Depositor DCC
R_{free} test set	34003 reflections (10.06%)	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtrriage
Anisotropy	0.223	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	20104	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	1/3197 (0.0%)	0.86	3/4353 (0.1%)
1	B	0.70	1/3286 (0.0%)	0.80	6/4473 (0.1%)
1	C	0.63	0/3168	0.71	1/4311 (0.0%)
1	D	0.51	0/3164	0.60	0/4306
1	E	0.36	0/3140	0.49	0/4275
1	F	0.34	0/3134	0.51	0/4268
All	All	0.60	2/19089 (0.0%)	0.68	10/25986 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	30	GLU	CD-OE2	-8.02	1.16	1.25
1	B	78	ARG	CG-CD	-5.07	1.39	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ARG	NE-CZ-NH1	7.00	123.80	120.30
1	B	78	ARG	NE-CZ-NH2	-6.41	117.10	120.30
1	A	139	ARG	NE-CZ-NH1	-6.09	117.25	120.30
1	A	111	ASP	CB-CG-OD1	6.04	123.74	118.30
1	B	78	ARG	CA-CB-CG	-5.82	100.59	113.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	239[A]	TRP	Peptide
1	B	242[A]	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3080	0	3026	24	0
1	B	3166	0	3083	44	0
1	C	3066	0	3002	28	0
1	D	3062	0	2997	38	0
1	E	3043	0	2978	33	0
1	F	3037	0	2975	45	0
2	A	31	0	19	0	0
2	B	31	0	19	0	0
2	C	31	0	19	1	0
2	D	31	0	19	1	0
2	E	31	0	19	0	0
2	F	31	0	19	1	0
3	A	20	0	0	0	0
3	B	25	0	0	0	0
3	C	15	0	0	0	0
3	D	10	0	0	0	0
3	E	10	0	0	0	0
3	F	10	0	0	0	0
4	A	499	0	0	5	2
4	B	358	0	0	6	1
4	C	313	0	0	1	1
4	D	182	0	0	5	0
4	E	13	0	0	0	0
4	F	9	0	0	0	0
All	All	20104	0	18175	213	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 213 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:383:GLU:OE2	4:A:501:HOH:O	1.82	0.97
1:A:139:ARG:NH1	4:A:502:HOH:O	1.97	0.96
1:B:240[A]:HIS:O	1:B:243[A]:PHE:N	2.02	0.93
1:C:181[B]:THR:HG22	1:C:294:GLN:HG3	1.55	0.89
1:B:237[A]:ASP:O	1:B:239[A]:TRP:N	2.06	0.88

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:584:HOH:O	4:B:604:HOH:O[7_555]	2.13	0.07
4:A:992:HOH:O	4:C:813:HOH:O[3_545]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/394 (102%)	392 (97%)	10 (2%)	2 (0%)	29	13
1	B	414/394 (105%)	397 (96%)	12 (3%)	5 (1%)	13	3
1	C	399/394 (101%)	389 (98%)	9 (2%)	1 (0%)	41	24
1	D	399/394 (101%)	388 (97%)	10 (2%)	1 (0%)	41	24
1	E	398/394 (101%)	384 (96%)	13 (3%)	1 (0%)	41	24
1	F	397/394 (101%)	373 (94%)	20 (5%)	4 (1%)	15	4
All	All	2411/2364 (102%)	2323 (96%)	74 (3%)	14 (1%)	25	11

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1	SER
1	B	321	SER
1	C	321	SER
1	D	321	SER
1	A	321	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	316/304 (104%)	304 (96%)	12 (4%)	33	14
1	B	322/304 (106%)	312 (97%)	10 (3%)	40	21
1	C	311/304 (102%)	300 (96%)	11 (4%)	36	17
1	D	311/304 (102%)	298 (96%)	13 (4%)	30	12
1	E	310/304 (102%)	296 (96%)	14 (4%)	27	10
1	F	309/304 (102%)	296 (96%)	13 (4%)	30	12
All	All	1879/1824 (103%)	1806 (96%)	73 (4%)	34	13

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	0	MET
1	D	196	PHE
1	F	214	GLN
1	D	159	LEU
1	D	215	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	403	-	4,4,4	0.56	0	6,6,6	0.70	0
3	SO4	E	403	-	4,4,4	0.12	0	6,6,6	0.21	0
2	FMN	E	401	-	31,33,33	2.51	5 (16%)	40,50,50	2.12	8 (20%)
3	SO4	B	404[A]	-	4,4,4	0.20	0	6,6,6	0.12	0
3	SO4	D	402	-	4,4,4	0.27	0	6,6,6	0.69	0
2	FMN	A	401	-	31,33,33	2.56	8 (25%)	40,50,50	2.92	10 (25%)
3	SO4	A	402	-	4,4,4	0.54	0	6,6,6	0.83	0
3	SO4	C	402	-	4,4,4	0.29	0	6,6,6	0.74	0
2	FMN	B	401	-	31,33,33	2.70	10 (32%)	40,50,50	2.44	11 (27%)
3	SO4	B	403	-	4,4,4	0.20	0	6,6,6	0.38	0
3	SO4	B	405	-	4,4,4	0.08	0	6,6,6	0.14	0
2	FMN	D	401	-	31,33,33	2.60	8 (25%)	40,50,50	2.29	9 (22%)
3	SO4	B	406[B]	-	4,4,4	0.42	0	6,6,6	0.36	0
3	SO4	F	403	-	4,4,4	0.16	0	6,6,6	0.26	0
3	SO4	C	404	-	4,4,4	0.18	0	6,6,6	0.16	0
3	SO4	C	403	-	4,4,4	0.16	0	6,6,6	1.00	0
3	SO4	A	405	-	4,4,4	0.26	0	6,6,6	0.14	0
2	FMN	F	401	-	31,33,33	2.43	5 (16%)	40,50,50	2.35	10 (25%)
3	SO4	D	403	-	4,4,4	0.41	0	6,6,6	0.42	0
3	SO4	F	402	-	4,4,4	0.11	0	6,6,6	0.19	0
2	FMN	C	401	-	31,33,33	2.73	9 (29%)	40,50,50	2.32	10 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	402	-	4,4,4	0.55	0	6,6,6	0.47	0
3	SO4	E	402	-	4,4,4	0.21	0	6,6,6	0.30	0
3	SO4	A	404[A]	-	4,4,4	0.26	0	6,6,6	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	E	401	-	-	3/18/18/18	0/3/3/3
2	FMN	C	401	-	-	1/18/18/18	0/3/3/3
2	FMN	A	401	-	-	1/18/18/18	0/3/3/3
2	FMN	F	401	-	-	5/18/18/18	0/3/3/3
2	FMN	D	401	-	-	1/18/18/18	0/3/3/3
2	FMN	B	401	-	-	1/18/18/18	0/3/3/3

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	401	FMN	C4A-C10	10.73	1.49	1.38
2	F	401	FMN	C4A-C10	10.29	1.49	1.38
2	B	401	FMN	C4A-C10	10.17	1.49	1.38
2	D	401	FMN	C4A-C10	9.86	1.48	1.38
2	A	401	FMN	C4A-C10	9.11	1.47	1.38

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	FMN	C4-N3-C2	9.18	122.89	115.14
2	A	401	FMN	C4-N3-C2	8.99	122.73	115.14
2	A	401	FMN	C4-C4A-C10	-8.63	114.24	119.95
2	F	401	FMN	C4-N3-C2	8.63	122.43	115.14
2	B	401	FMN	C4-N3-C2	8.32	122.17	115.14

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	401	FMN	C2'-C3'-C4'-O4'
2	F	401	FMN	C2'-C3'-C4'-O4'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	F	401	FMN	C2'-C3'-C4'-C5'
2	E	401	FMN	C4'-C5'-O5'-P
2	B	401	FMN	C4'-C5'-O5'-P


There are no ring outliers.


3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	FMN	1	0
2	F	401	FMN	1	0
2	C	401	FMN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	394/394 (100%)	0.28	21 (5%) 26 29	19, 30, 49, 77	0
1	B	394/394 (100%)	0.66	52 (13%) 3 3	25, 38, 88, 131	0
1	C	394/394 (100%)	0.47	45 (11%) 5 5	30, 43, 70, 100	0
1	D	394/394 (100%)	1.16	85 (21%) 0 0	46, 62, 103, 160	0
1	E	394/394 (100%)	3.40	241 (61%) 0 0	72, 103, 158, 190	394 (100%)
1	F	393/394 (99%)	4.05	282 (71%) 0 0	87, 113, 169, 204	393 (100%)
All	All	2363/2364 (99%)	1.67	726 (30%) 0 0	19, 57, 137, 204	787 (33%)

The worst 5 of 726 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	223	VAL	28.1
1	E	238	PHE	23.5
1	E	234	LEU	21.9
1	F	222	GLY	19.6
1	F	226	GLU	18.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	406[B]	5/5	0.71	0.23	50,56,56,59	5
3	SO4	A	404[A]	5/5	0.78	0.31	50,56,58,59	5
2	FMN	F	401	31/31	0.81	0.18	74,79,82,82	31
3	SO4	F	402	5/5	0.84	0.14	88,89,89,90	5
3	SO4	E	403	5/5	0.85	0.14	94,95,95,96	5
2	FMN	E	401	31/31	0.86	0.20	74,78,81,81	31
3	SO4	B	404[A]	5/5	0.87	0.22	67,67,68,70	5
3	SO4	C	404	5/5	0.90	0.24	66,66,67,68	5
3	SO4	F	403	5/5	0.93	0.23	72,72,74,74	5
3	SO4	A	405	5/5	0.94	0.19	63,64,64,65	5
3	SO4	B	405	5/5	0.94	0.26	99,99,101,104	0
2	FMN	C	401	31/31	0.96	0.16	29,33,37,39	0
3	SO4	E	402	5/5	0.96	0.26	98,98,99,99	0
2	FMN	D	401	31/31	0.96	0.10	39,46,51,52	0
2	FMN	B	401	31/31	0.97	0.12	23,29,33,36	0
2	FMN	A	401	31/31	0.98	0.15	19,23,26,30	0
3	SO4	D	403	5/5	0.98	0.11	64,64,69,71	0
3	SO4	C	403	5/5	0.99	0.08	36,42,43,45	0
3	SO4	B	403	5/5	0.99	0.09	38,40,44,45	0
3	SO4	B	402	5/5	0.99	0.08	38,39,43,43	0
3	SO4	A	403	5/5	0.99	0.12	35,35,39,39	0
3	SO4	D	402	5/5	0.99	0.11	51,51,55,56	0
3	SO4	A	402	5/5	1.00	0.06	30,34,35,37	0
3	SO4	C	402	5/5	1.00	0.08	39,44,46,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FMN F 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FMN E 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FMN C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FMN D 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FMN B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.