



wwPDB X-ray Structure Validation Summary Report

Oct 18, 2023 – 02:02 PM EDT

PDB ID : 2E2G
Title : Crystal structure of archaeal peroxiredoxin, thioredoxin peroxidase from *Aeropyrum pernix* K1 (pre-oxidation form)
Authors : Nakamura, T.; Yamamoto, T.; Abe, M.; Matsumura, H.; Hagihara, Y.; Goto, T.; Yamaguchi, T.; Inoue, T.
Deposited on : 2006-11-13
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	 70% 21% 5% 5%
1	B	250	 64% 25% 5% 5%
1	C	250	 64% 26% 5% 5%
1	D	250	 69% 22% 5% 5%
1	E	250	 66% 22% 5% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	250	 66% 22% 5% • 5%
1	G	250	 67% 23% • • 5%
1	H	250	 66% 22% 5% • 5%
1	I	250	 63% 29% • • 5%
1	J	250	 69% 22% • 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 19878 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable peroxiredoxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	238	Total 1932	C 1244	N 339	O 343	S 6	0	0	0
1	B	239	Total 1942	C 1250	N 342	O 344	S 6	0	0	0
1	C	239	Total 1938	C 1247	N 340	O 345	S 6	0	0	0
1	D	238	Total 1932	C 1244	N 339	O 343	S 6	0	0	0
1	E	238	Total 1932	C 1244	N 339	O 343	S 6	0	0	0
1	F	238	Total 1932	C 1244	N 339	O 343	S 6	0	0	0
1	G	238	Total 1932	C 1244	N 339	O 343	S 6	0	0	0
1	H	238	Total 1932	C 1244	N 339	O 343	S 6	0	0	0
1	I	238	Total 1932	C 1244	N 339	O 343	S 6	0	0	0
1	J	238	Total 1932	C 1244	N 339	O 343	S 6	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	207	SER	CYS	engineered mutation	UNP Q9Y9L0
B	207	SER	CYS	engineered mutation	UNP Q9Y9L0
C	207	SER	CYS	engineered mutation	UNP Q9Y9L0
D	207	SER	CYS	engineered mutation	UNP Q9Y9L0
E	207	SER	CYS	engineered mutation	UNP Q9Y9L0
F	207	SER	CYS	engineered mutation	UNP Q9Y9L0
G	207	SER	CYS	engineered mutation	UNP Q9Y9L0
H	207	SER	CYS	engineered mutation	UNP Q9Y9L0
I	207	SER	CYS	engineered mutation	UNP Q9Y9L0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	207	SER	CYS	engineered mutation	UNP Q9Y9L0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	57	Total O 57 57	0	0
2	B	60	Total O 60 60	0	0
2	C	39	Total O 39 39	0	0
2	D	57	Total O 57 57	0	0
2	E	65	Total O 65 65	0	0
2	F	59	Total O 59 59	0	0
2	G	61	Total O 61 61	0	0
2	H	65	Total O 65 65	0	0
2	I	30	Total O 30 30	0	0
2	J	49	Total O 49 49	0	0


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable peroxiredoxin



- Molecule 1: Probable peroxiredoxin




Chain D:  69% 22% 5%




• Molecule 1: Probable peroxiredoxin

Chain E:  66% 22% 5% 5%




• Molecule 1: Probable peroxiredoxin

Chain F:  66% 22% 5% 5%



• Molecule 1: Probable peroxiredoxin


Chain G:  67% 23% 5% 5%







- Molecule 1: Probable peroxiredoxin

Chain H: 66% 22% 5% • 5%



- Molecule 1: Probable peroxiredoxin

Chain I: 63% 29% •• 5%



- Molecule 1: Probable peroxiredoxin

Chain J: 69% 22% • 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	76.20Å 103.35Å 104.63Å 105.79° 105.19° 92.68°	Depositor
Resolution (Å)	19.99 – 2.40 49.48 – 2.39	Depositor EDS
% Data completeness (in resolution range)	86.8 (19.99-2.40) 86.8 (49.48-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.153 , 0.229 0.160 , 0.158	Depositor DCC
R_{free} test set	5083 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtrriage
Anisotropy	0.222	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	19878	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.16	1/1984 (0.1%)	1.08	7/2696 (0.3%)
1	B	1.17	3/1995 (0.2%)	1.13	12/2711 (0.4%)
1	C	1.16	5/1990 (0.3%)	1.12	10/2704 (0.4%)
1	D	1.12	3/1984 (0.2%)	1.08	6/2696 (0.2%)
1	E	1.22	7/1984 (0.4%)	1.13	12/2696 (0.4%)
1	F	1.26	15/1984 (0.8%)	1.13	15/2696 (0.6%)
1	G	1.15	5/1984 (0.3%)	1.05	5/2696 (0.2%)
1	H	1.19	8/1984 (0.4%)	1.08	7/2696 (0.3%)
1	I	1.11	3/1984 (0.2%)	1.08	9/2696 (0.3%)
1	J	1.12	2/1984 (0.1%)	1.03	6/2696 (0.2%)
All	All	1.17	52/19857 (0.3%)	1.09	89/26983 (0.3%)

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	183	GLU	CD-OE2	10.14	1.36	1.25
1	H	225	GLU	CG-CD	8.59	1.64	1.51
1	F	62	GLU	CG-CD	8.18	1.64	1.51
1	F	225	GLU	CG-CD	7.85	1.63	1.51
1	E	225	GLU	CG-CD	7.60	1.63	1.51

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	206	ARG	NE-CZ-NH1	11.66	126.13	120.30
1	C	206	ARG	NE-CZ-NH1	10.83	125.72	120.30
1	C	126	ARG	NE-CZ-NH1	10.80	125.70	120.30
1	E	228	ARG	NE-CZ-NH1	10.50	125.55	120.30
1	B	149	ARG	NE-CZ-NH2	-9.89	115.36	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1932	0	1924	58	0
1	B	1942	0	1931	69	0
1	C	1938	0	1929	63	0
1	D	1932	0	1924	48	0
1	E	1932	0	1924	63	0
1	F	1932	0	1924	55	1
1	G	1932	0	1924	64	0
1	H	1932	0	1924	60	1
1	I	1932	0	1924	54	0
1	J	1932	0	1924	44	0
2	A	57	0	0	0	0
2	B	60	0	0	2	0
2	C	39	0	0	4	0
2	D	57	0	0	4	0
2	E	65	0	0	9	0
2	F	59	0	0	6	0
2	G	61	0	0	5	0
2	H	65	0	0	6	0
2	I	30	0	0	2	0
2	J	49	0	0	1	0
All	All	19878	0	19252	524	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 524 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:206:ARG:HG3	1:F:206:ARG:HH11	1.10	1.15
1:A:123:HIS:HA	1:A:145:MET:HE1	1.33	1.10
1:I:129:PHE:CE2	1:I:140:MET:HE3	1.88	1.08
1:A:206:ARG:HG3	1:A:206:ARG:HH11	1.18	1.07
1:E:220:ARG:HG2	1:E:220:ARG:HH11	0.96	1.06

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:228:ARG:NH1	1:H:228:ARG:CB[1_544]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/250 (94%)	224 (96%)	7 (3%)	3 (1%)	12	17
1	B	235/250 (94%)	228 (97%)	6 (3%)	1 (0%)	34	48
1	C	235/250 (94%)	219 (93%)	15 (6%)	1 (0%)	34	48
1	D	234/250 (94%)	219 (94%)	13 (6%)	2 (1%)	17	25
1	E	234/250 (94%)	223 (95%)	11 (5%)	0	100	100
1	F	234/250 (94%)	224 (96%)	8 (3%)	2 (1%)	17	25
1	G	234/250 (94%)	217 (93%)	15 (6%)	2 (1%)	17	25
1	H	234/250 (94%)	221 (94%)	10 (4%)	3 (1%)	12	17
1	I	234/250 (94%)	221 (94%)	10 (4%)	3 (1%)	12	17
1	J	234/250 (94%)	211 (90%)	21 (9%)	2 (1%)	17	25
All	All	2342/2500 (94%)	2207 (94%)	116 (5%)	19 (1%)	19	29

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	THR
1	C	121	ALA
1	F	32	SER
1	F	122	THR
1	I	122	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/216 (95%)	189 (92%)	17 (8%)	11	17
1	B	207/216 (96%)	182 (88%)	25 (12%)	5	6
1	C	207/216 (96%)	191 (92%)	16 (8%)	13	20
1	D	206/216 (95%)	188 (91%)	18 (9%)	10	15
1	E	206/216 (95%)	185 (90%)	21 (10%)	7	10
1	F	206/216 (95%)	188 (91%)	18 (9%)	10	15
1	G	206/216 (95%)	191 (93%)	15 (7%)	14	22
1	H	206/216 (95%)	185 (90%)	21 (10%)	7	10
1	I	206/216 (95%)	188 (91%)	18 (9%)	10	15
1	J	206/216 (95%)	190 (92%)	16 (8%)	12	19
All	All	2062/2160 (96%)	1877 (91%)	185 (9%)	9	14

5 of 185 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	24	ILE
1	H	197	ARG
1	G	88	TRP
1	H	11	ARG
1	H	242	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	204	GLN
1	G	92	HIS
1	I	195	GLN
1	I	92	HIS
1	I	106	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	238/250 (95%)	-0.75	2 (0%) 86 84	12, 22, 45, 72	0
1	B	239/250 (95%)	-0.63	0 100 100	14, 24, 44, 62	0
1	C	239/250 (95%)	-0.59	0 100 100	15, 25, 44, 71	0
1	D	238/250 (95%)	-0.66	0 100 100	13, 25, 46, 76	0
1	E	238/250 (95%)	-0.59	2 (0%) 86 84	13, 25, 44, 65	0
1	F	238/250 (95%)	-0.70	0 100 100	11, 22, 41, 65	0
1	G	238/250 (95%)	-0.63	0 100 100	14, 24, 44, 59	0
1	H	238/250 (95%)	-0.69	0 100 100	13, 24, 45, 63	0
1	I	238/250 (95%)	-0.56	1 (0%) 92 91	18, 30, 49, 68	0
1	J	238/250 (95%)	-0.69	1 (0%) 92 91	15, 27, 48, 63	0
All	All	2382/2500 (95%)	-0.65	6 (0%) 94 93	11, 25, 46, 76	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	238	ALA	4.3
1	A	245	ALA	2.8
1	A	244	GLU	2.7
1	E	242	TYR	2.4
1	E	244	GLU	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.