

# wwPDB X-ray Structure Validation Summary Report (i)

Oct 15, 2023 – 09:21 PM EDT

PDB ID	:	8E6M
Title	:	X-ray structure of the Deinococcus radiodurans Nramp/MntH divalent tran-
		sition metal transporter WT in an inward-open, cadmium-bound state
Authors	:	Ray, S.; Gaudet, R.
Deposited on	:	2022-08-22
Resolution	:	2.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity Mogul Xtriage (Phenix) EDS	:	4.02b-467 1.8.5 (274361), CSD as541be (2020) 1.13 2.36
buster-report Percentile statistics Refmac	: : :	1.1.7 (2018) 20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
			7%						
1	А	414	84%	9%	6%				

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	OLC	А	515	-	-	-	Х



$\alpha$ $\cdot$ $\cdot$ $\cdot$	C		
Continued	trom	nremons	naae
Continucu	JION	precious	page

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	OLC	А	517	-	-	-	Х
4	OLC	А	520	-	-	-	Х



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 3321 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Divalent metal cation transporter MntH.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	388	Total 2905	C 1914	N 483	0 491	S 17	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	23	MET	-	expression tag	UNP Q9RTP8
А	24	HIS	-	expression tag	UNP Q9RTP8
А	25	HIS	-	expression tag	UNP Q9RTP8
А	26	HIS	-	expression tag	UNP Q9RTP8
А	27	HIS	-	expression tag	UNP Q9RTP8
А	28	HIS	-	expression tag	UNP Q9RTP8
А	29	HIS	-	expression tag	UNP Q9RTP8
А	30	HIS	-	expression tag	UNP Q9RTP8
А	31	HIS	-	expression tag	UNP Q9RTP8

• Molecule 2 is CADMIUM ION (three-letter code: CD) (formula: Cd).


Mo	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Cd 2 2	0	0

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	2	Total Cl 2 2	0	0

• Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula:  $C_{21}H_{40}O_4$ ).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total         C         O           25         21         4	0	0
4	А	1	Total         C         O           14         10         4	0	0
4	А	1	Total         C         O           22         18         4	0	0
4	А	1	Total         C         O           11         7         4	0	0
4	А	1	Total         C         O           25         21         4	0	0
4	А	1	Total         C         O           25         21         4	0	0
4	А	1	Total         C         O           24         20         4	0	0
4	А	1	Total         C         O           23         19         4	0	0
4	А	1	Total         C         O           21         17         4	0	0
4	А	1	Total         C         O           19         15         4	0	0
4	А	1	Total         C         O           19         15         4	0	0
4	А	1	Total         C         O           15         11         4	0	0
4	А	1	Total         C         O           23         19         4	0	0
4	А	1	Total C 11 11	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total C O 15 11 4	0	0
4	А	1	Total         C           16         16	0	0
4	А	1	Total C 13 13	0	0
4	А	1	Total C O 14 10 4	0	0
4	А	1	$\begin{array}{cc} \text{Total} & \text{C} \\ 7 & 7 \end{array}$	0	0
4	А	1	TotalC1212	0	0

• Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0


• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	54	$\begin{array}{cc} \text{Total} & \text{O} \\ 54 & 54 \end{array}$	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants	59.14Å 71.37Å 99.05Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	45.54 - 2.48	Depositor
Resolution (A)	45.54 - 2.48	EDS
% Data completeness	97.3 (45.54-2.48)	Depositor
(in resolution range)	$84.6\ (45.54-2.48)$	EDS
R <sub>merge</sub>	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.70 (at 2.48 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
D D.	0.203 , $0.250$	Depositor
$R, R_{free}$	0.206 , $0.251$	DCC
$R_{free}$ test set	1507  reflections  (10.03%)	wwPDB-VP
Wilson B-factor $(Å^2)$	40.2	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32 , $64.6$	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3321	wwPDB-VP
Average B, all atoms $(Å^2)$	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.29% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PEG, CD, OLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.24	0/2969	0.44	0/4058	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2905	0	3040	29	0
2	А	2	0	0	0	0
3	А	2	0	0	0	0
4	А	354	0	513	21	0
5	А	4	0	5	0	0
6	А	54	0	0	1	0
All	All	3321	0	3558	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:LYS:HB3	1:A:405:LEU:HD11	1.73	0.71
1:A:115:LEU:HD21	4:A:508:OLC:H2A	1.77	0.67
1:A:114:PRO:HB3	4:A:522:OLC:H21	1.77	0.66
1:A:413:VAL:HG22	4:A:505:OLC:H5A	1.76	0.65
1:A:192:VAL:HA	4:A:507:OLC:H2A	1.79	0.65

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	А	384/414~(93%)	380~(99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	297/322 ( $92%$ )	296 (100%)	1 (0%)	92 97		

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	117	TRP



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	82	ASN
1	А	232	HIS
1	А	237	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 25 ligands modelled in this entry, 4 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain Res Link		Link	Bo	ond leng	ths	Bond angles		
10101	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	OLC	А	522	-	13,13,24	1.06	2 (15%)	$14,\!14,\!25$	0.98	1 (7%)
5	PEG	А	525	-	3,3,6	0.19	0	2,2,5	0.22	0
4	OLC	А	514	-	18,18,24	0.93	2 (11%)	$18,\!19,\!25$	1.02	1 (5%)
4	OLC	А	511	-	23,23,24	0.83	2 (8%)	$24,\!24,\!25$	0.96	1 (4%)
4	OLC	А	506	-	13,13,24	1.05	2 (15%)	14,14,25	1.11	1 (7%)
4	OLC	А	513	-	20,20,24	0.87	2 (10%)	21,21,25	1.02	1 (4%)
4	OLC	А	509	-	24,24,24	0.82	2 (8%)	$25,\!25,\!25$	0.95	1 (4%)
4	OLC	А	512	-	22,22,24	0.84	2 (9%)	23,23,25	1.03	1 (4%)



Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
10101	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	OLC	А	518	-	10,10,24	0.38	0	8,9,25	0.75	0
4	OLC	А	523	-	6,6,24	0.31	0	$5,\!5,\!25$	0.67	0
4	OLC	А	524	-	11,11,24	0.37	0	$9,\!10,\!25$	0.80	0
4	OLC	А	516	-	14,14,24	1.01	2 (14%)	$15,\!15,\!25$	1.00	1 (6%)
4	OLC	А	521	-	12,12,24	0.34	0	11,11,25	0.72	0
4	OLC	А	520	-	15,15,24	0.36	0	14,14,25	0.64	0
4	OLC	А	519	-	14,14,24	1.03	2 (14%)	$15,\!15,\!25$	1.07	1 (6%)
4	OLC	А	505	-	24,24,24	0.79	2 (8%)	$25,\!25,\!25$	0.96	1 (4%)
4	OLC	А	507	-	21,21,24	0.86	2 (9%)	22,22,25	0.96	1 (4%)
4	OLC	А	515	-	18,18,24	0.94	2 (11%)	18,19,25	1.05	1 (5%)
4	OLC	А	510	-	24,24,24	0.81	1 (4%)	25,25,25	0.98	1 (4%)
4	OLC	А	517	-	22,22,24	0.83	2 (9%)	23,23,25	0.91	1 (4%)
4	OLC	А	508	-	10,10,24	1.21	2 (20%)	11,11,25	1.13	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	А	522	-	-	9/13/13/24	-
5	PEG	А	525	-	-	1/1/1/4	-
4	OLC	А	514	-	-	10/18/18/24	-
4	OLC	А	511	-	-	9/23/23/24	-
4	OLC	А	506	-	-	0/13/13/24	-
4	OLC	А	513	-	-	8/20/20/24	-
4	OLC	А	509	-	-	12/24/24/24	-
4	OLC	А	512	-	-	11/22/22/24	-
4	OLC	А	518	-	-	6/8/8/24	-
4	OLC	А	523	-	-	2/4/4/24	-
4	OLC	А	524	-	-	4/9/9/24	-
4	OLC	А	516	-	-	7/14/14/24	-
4	OLC	А	521	-	-	4/10/10/24	-
4	OLC	А	520	-	-	6/13/13/24	-
4	OLC	А	519	-	-	9/14/14/24	-
4	OLC	А	505	-	-	10/24/24/24	-
4	OLC	А	507	-	-	6/21/21/24	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	А	515	-	-	13/18/18/24	-
4	OLC	А	510	-	-	11/24/24/24	-
4	OLC	А	517	-	-	10/22/22/24	-
4	OLC	А	508	-	-	4/10/10/24	-

Continued from previous page...

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	А	512	OLC	O20-C1	2.56	1.40	1.33
4	А	515	OLC	O20-C1	2.55	1.40	1.33
4	А	510	OLC	O20-C1	2.49	1.40	1.33
4	А	513	OLC	O20-C1	2.49	1.40	1.33
4	А	508	OLC	O20-C1	2.46	1.40	1.33

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
4	А	515	OLC	O20-C1-C2	2.94	121.14	111.91
4	А	512	OLC	O20-C1-C2	2.87	120.92	111.91
4	А	510	OLC	O20-C1-C2	2.82	120.75	111.91
4	А	506	OLC	O20-C1-C2	2.79	120.65	111.91
4	А	513	OLC	O20-C1-C2	2.73	120.48	111.91

There are no chirality outliers.

5 of 152 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	505	OLC	C21-C22-C24-O25
4	А	508	OLC	O20-C21-C22-C24
4	А	510	OLC	C2-C1-O20-C21
4	А	510	OLC	O19-C1-O20-C21
4	А	512	OLC	C21-C22-C24-O25

There are no ring outliers.

14 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	А	522	OLC	2	0
4	А	514	OLC	1	0
4	А	511	OLC	1	0




Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	А	506	OLC	2	0
4	А	509	OLC	1	0
4	А	512	OLC	3	0
4	А	524	OLC	2	0
4	А	516	OLC	1	0
4	А	519	OLC	1	0
4	А	505	OLC	3	0
4	А	507	OLC	2	0
4	А	515	OLC	1	0
4	А	510	OLC	3	0
4	А	508	OLC	1	0

Continued from previous page...


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	А	388/414~(93%)	0.20	30 (7%) 13 13	32, 53, 115, 182	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	49	ILE	6.1
1	А	170	ARG	5.9
1	А	173	ARG	5.1
1	А	174	PRO	5.1
1	А	248	ASP	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{\AA}^2)$	Q<0.9
4	OLC	А	522	14/25	0.44	0.40	$65,\!84,\!98,\!112$	0
4	OLC	А	517	23/25	0.60	0.49	60,101,114,124	0




	0	m previoi	1 0		Daga	DOD	<b>D a</b> (82)	
Mol	Type	Chain	$\mathbf{Res}$	Atoms	RSCC	$\mathbf{RSR}$	$\operatorname{B-factors}(\operatorname{\AA}^2)$	Q < 0.9
4	OLC	А	520	16/25	0.62	0.46	$69,\!84,\!99,\!100$	0
4	OLC	А	514	19/25	0.67	0.33	74,93,104,104	0
4	OLC	А	515	19/25	0.69	0.42	$65,\!81,\!95,\!105$	0
4	OLC	А	521	13/25	0.70	0.39	66,76,82,84	0
4	OLC	А	511	24/25	0.72	0.32	$67,\!83,\!98,\!103$	0
4	OLC	А	512	23/25	0.72	0.28	60,79,98,113	0
4	OLC	А	513	21/25	0.74	0.31	56,76,88,93	0
5	PEG	А	525	4/7	0.78	0.13	62,70,72,72	0
4	OLC	А	509	25/25	0.79	0.33	$67,\!83,\!102,\!111$	0
4	OLC	А	524	12/25	0.81	0.20	$65,\!79,\!90,\!92$	0
4	OLC	А	505	25/25	0.81	0.24	63,87,97,99	0
4	OLC	А	519	15/25	0.82	0.25	57,89,106,108	0
4	OLC	А	510	25/25	0.82	0.20	59,74,85,96	0
4	OLC	А	516	15/25	0.83	0.33	64,79,92,94	0
4	OLC	А	518	11/25	0.83	0.22	74,87,90,90	0
4	OLC	А	507	22/25	0.85	0.28	47,66,92,105	0
4	OLC	А	508	11/25	0.85	0.21	59,68,72,83	0
2	CD	А	501	1/1	0.85	0.10	83,83,83,83	1
4	OLC	А	506	14/25	0.86	0.21	52,73,97,102	0
4	OLC	А	523	7/25	0.87	0.24	76,79,89,90	0
3	CL	А	504	1/1	0.93	0.08	63,63,63,63	0
3	CL	А	503	1/1	0.93	0.12	70,70,70,70	0
2	CD	А	502	1/1	0.99	0.12	$65,\!65,\!65,\!65$	0


Continued from previous page..

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















































## 6.5 Other polymers (i)

There are no such residues in this entry.

