



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 06:52 pm BST

PDB ID : 3ECO
Title : Crystal structure of MepR, a transcription regulator of the *Staphylococcus aureus* multidrug efflux pump MepA
Authors : Brennan, R.G.; Kumaraswami, M.
Deposited on : 2008-09-01
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	139	 71% 25% ••
1	B	139	 69% 20% • 7%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MepR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	135	1080	676	186	213	5	0	0	0
1	B	129	1046	656	179	206	5	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	B	1	5	4	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	97	97	97	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	99	Total	O	0	0
			99	99		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MepR

Chain A: 



- Molecule 1: MepR

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	32.20Å 96.57Å 110.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.89 – 2.40 47.89 – 1.49	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.89-2.40) 53.1 (47.89-1.49)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.50Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.242 , 0.289 0.241 , 0.290	Depositor DCC
R_{free} test set	1395 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtrriage
Anisotropy	0.209	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	2327	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/1092	0.53	0/1466
1	B	0.38	0/1058	0.55	0/1419
All	All	0.36	0/2150	0.54	0/2885

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1080	0	1071	30	0
1	B	1046	0	1050	21	0
2	B	5	0	0	0	0
3	A	97	0	0	3	0
3	B	99	0	0	4	0
All	All	2327	0	2121	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:44:GLN:NE2	1:B:93:LEU:H	1.74	0.85
1:A:135:LEU:O	1:A:139:GLN:HG3	1.77	0.83
1:A:4:THR:HG23	1:A:7:TYR:H	1.47	0.80
1:A:45:GLN:HB2	3:A:180:HOH:O	1.88	0.72
1:A:4:THR:CG2	1:A:7:TYR:H	2.03	0.70
1:A:53:ILE:O	1:A:57:LEU:HB2	1.93	0.69
1:A:121:GLU:O	1:A:125:GLU:HG3	1.94	0.67
1:A:120:SER:OG	1:A:123:GLU:HG3	1.96	0.65
1:A:4:THR:HG22	1:A:7:TYR:HB3	1.79	0.65
1:A:16:MET:CE	1:B:6:SER:HA	2.28	0.63
1:A:134:MET:O	1:A:138:LEU:HD23	1.99	0.63
1:A:40:LEU:HD23	1:A:48:LEU:HD12	1.79	0.63
1:A:133:LYS:HD2	3:B:234:HOH:O	2.00	0.61
1:A:44:GLN:HB2	3:A:144:HOH:O	2.01	0.61
1:A:133:LYS:HB3	3:B:234:HOH:O	2.03	0.58
1:B:118:GLN:HG3	3:B:234:HOH:O	2.03	0.58
1:B:43:HIS:O	1:B:48:LEU:HD11	2.04	0.57
1:B:23:LYS:HD2	1:B:107:ILE:HG23	1.87	0.57
1:B:115:LEU:O	1:B:118:GLN:HB2	2.06	0.54
1:A:138:LEU:HD21	1:B:115:LEU:HD21	1.89	0.54
1:A:68:LEU:O	1:A:72:GLU:HB2	2.09	0.53
1:A:54:ALA:HA	1:A:64:VAL:HG21	1.92	0.52
1:A:4:THR:HG22	1:A:7:TYR:CB	2.39	0.52
1:B:40:LEU:O	1:B:44:GLN:HG3	2.10	0.51
1:B:10:ARG:HG3	1:B:11:MET:N	2.24	0.51
1:A:16:MET:HE2	1:B:6:SER:HA	1.93	0.49
1:A:30:THR:OG1	1:A:33:GLN:HG3	2.12	0.49
1:B:23:LYS:O	1:B:26:GLN:HB2	2.13	0.49
1:B:41:TYR:HE1	1:B:98:ILE:HG23	1.78	0.49
3:A:193:HOH:O	1:B:133:LYS:HD3	2.12	0.48
1:B:30:THR:H	1:B:33:GLN:NE2	2.14	0.45
1:A:43:HIS:HB2	1:A:48:LEU:HD11	1.98	0.45
1:B:61:GLY:N	1:B:62:PRO:HD2	2.32	0.45
1:A:48:LEU:O	1:A:90:ASN:HA	2.16	0.45
1:A:116:VAL:C	1:A:118:GLN:H	2.21	0.44
1:B:66:ASN:ND2	3:B:248:HOH:O	2.50	0.44
1:A:115:LEU:O	1:A:118:GLN:HB2	2.18	0.43
1:A:130:ASN:O	1:A:134:MET:HG3	2.19	0.43
1:A:16:MET:HE1	1:B:6:SER:HA	1.99	0.43
1:B:24:LEU:HD22	1:B:29:ILE:HB	2.00	0.43
1:B:72:GLU:OE2	1:B:79:ARG:HD3	2.19	0.42
1:B:98:ILE:O	1:B:102:GLU:HG3	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:GLY:N	1:A:62:PRO:HD2	2.34	0.42
1:A:64:VAL:O	1:A:68:LEU:HG	2.20	0.42
1:A:19:LYS:HD3	1:A:111:MET:CE	2.50	0.41
1:A:116:VAL:HA	1:A:124:ASN:HD21	1.86	0.41
1:B:114:THR:O	1:B:118:GLN:NE2	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/139 (94%)	128 (98%)	3 (2%)	0	100	100
1	B	125/139 (90%)	123 (98%)	1 (1%)	1 (1%)	19	29
All	All	256/278 (92%)	251 (98%)	4 (2%)	1 (0%)	34	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	45	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/126 (94%)	114 (97%)	4 (3%)	37	56
1	B	117/126 (93%)	107 (92%)	10 (8%)	10	16
All	All	235/252 (93%)	221 (94%)	14 (6%)	19	31

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	MET
1	A	65	SER
1	A	72	GLU
1	A	100	LEU
1	B	10	ARG
1	B	24	LEU
1	B	45	GLN
1	B	57	LEU
1	B	70	ASN
1	B	71	LEU
1	B	79	ARG
1	B	100	LEU
1	B	118	GLN
1	B	138	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	50	GLN
1	A	90	ASN
1	A	124	ASN
1	B	26	GLN
1	B	33	GLN
1	B	44	GLN
1	B	45	GLN
1	B	50	GLN
1	B	66	ASN
1	B	70	ASN
1	B	118	GLN
1	B	124	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	140	-	4,4,4	0.28	0	6,6,6	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.