

wwPDB X-ray Structure Validation Summary Report (i)

Oct 26, 2023 – 05:44 AM EDT

PDB ID : 3ED1

Title : Crystal Structure of Rice GID1 complexed with GA3

Authors: Shimada, A.; Nakatsu, T.; Ueguchi-Tanaka, M.; Kato, H.; Matsuoka, M.

Deposited on : 2008-09-02

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS: 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	Λ	265	2%			
1	A	365	81% 6%		7%	12%
1	В	365	82%		5%	13%
1	С	365	74%	8%	_	18%
1	D	365	80%	6	5% •	14%
1	Е	365	<mark>6%</mark> 78%	7%	6 •	14%



Continued from previous page...

Mo	l Chain	Length	Quality of chain	• •				
			2%					
1	F	365	79%	5%	16%			



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 15936 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Gibberellin receptor GID1.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	A	321	Total	С	N	О	S	0	4	0
1	Λ	321	2510	1591	447	462	10	U	4	
1	В	316	Total	С	N	О	S	0	5	0
1	Ъ	310	2476	1572	441	453	10	U	3	
1	С	301	Total	С	N	О	S	0	3	0
1		301	2365	1507	416	432	10	U		
1	D	315	Total	С	N	О	O S 0 3	3	0	
1	D	313	2455	1561	429	455	10	U	3	
1	Е	313	Total	С	N	О	S	0	2	0
1	l L	313	2390	1526	416	437	11	U		
1	E	305	Total	С	N	О	S	0	2	0
1	1 F	300	2378	1518	416	434	10	U		

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	355	GLY	-	expression tag	UNP Q6L545
A	356	SER	-	expression tag	UNP Q6L545
A	357	HIS	-	expression tag	UNP Q6L545
A	358	HIS	-	expression tag	UNP Q6L545
A	359	HIS	-	expression tag	UNP Q6L545
A	360	HIS	-	expression tag	UNP Q6L545
A	361	HIS	-	expression tag	UNP Q6L545
A	362	HIS	-	expression tag	UNP Q6L545
A	363	HIS	-	expression tag	UNP Q6L545
A	364	HIS	-	expression tag	UNP Q6L545
A	365	HIS	-	expression tag	UNP Q6L545
A	366	HIS	-	expression tag	UNP Q6L545
В	355	GLY	-	expression tag	UNP Q6L545
В	356	SER	-	expression tag	UNP Q6L545
В	357	HIS	=	expression tag	UNP Q6L545
В	358	HIS	-	expression tag	UNP Q6L545
В	359	HIS	-	expression tag	UNP Q6L545



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled	Actual	Comment	Reference
В	360	HIS	-	expression tag	UNP Q6L545
В	361	HIS	-	expression tag	UNP Q6L545
В	362	HIS	-	expression tag	UNP Q6L545
В	363	HIS	-	expression tag	UNP Q6L545
В	364	HIS	-	expression tag	UNP Q6L545
В	365	HIS	-	expression tag	UNP Q6L545
В	366	HIS	-	expression tag	UNP Q6L545
С	355	GLY	-	expression tag	UNP Q6L545
С	356	SER	-	expression tag	UNP Q6L545
С	357	HIS	-	expression tag	UNP Q6L545
С	358	HIS	-	expression tag	UNP Q6L545
С	359	HIS	-	expression tag	UNP Q6L545
С	360	HIS	_	expression tag	UNP Q6L545
С	361	HIS	-	expression tag	UNP Q6L545
С	362	HIS	-	expression tag	UNP Q6L545
С	363	HIS	-	expression tag	UNP Q6L545
С	364	HIS	-	expression tag	UNP Q6L545
С	365	HIS	-	expression tag	UNP Q6L545
С	366	HIS	-	expression tag	UNP Q6L545
D	355	GLY	-	expression tag	UNP Q6L545
D	356	SER	-	expression tag	UNP Q6L545
D	357	HIS	-	expression tag	UNP Q6L545
D	358	HIS	-	expression tag	UNP Q6L545
D	359	HIS	-	expression tag	UNP Q6L545
D	360	HIS	-	expression tag	UNP Q6L545
D	361	HIS	-	expression tag	UNP Q6L545
D	362	HIS	-	expression tag	UNP Q6L545
D	363	HIS	-	expression tag	UNP Q6L545
D	364	HIS	-	expression tag	UNP Q6L545
D	365	HIS	-	expression tag	UNP Q6L545
D	366	HIS	-	expression tag	UNP Q6L545
Е	355	GLY	-	expression tag	UNP Q6L545
Е	356	SER	-	expression tag	UNP Q6L545
Е	357	HIS	-	expression tag	UNP Q6L545
Е	358	HIS	-	expression tag	UNP Q6L545
Е	359	HIS	-	expression tag	UNP Q6L545
Е	360	HIS		expression tag	UNP Q6L545
Е	361	HIS		expression tag	UNP Q6L545
Е	362	HIS		expression tag	UNP Q6L545
Е	363	HIS	-	expression tag	UNP Q6L545
Е	364	HIS	-	expression tag	UNP Q6L545
Е	365	HIS	-	expression tag	UNP Q6L545



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Е	366	HIS	-	expression tag	UNP Q6L545
F	355	GLY	-	expression tag	UNP Q6L545
F	356	SER	-	expression tag	UNP Q6L545
F	357	HIS	-	expression tag	UNP Q6L545
F	358	HIS	-	expression tag	UNP Q6L545
F	359	HIS	ı	expression tag	UNP Q6L545
F	360	HIS	-	expression tag	UNP Q6L545
F	361	HIS	-	expression tag	UNP Q6L545
F	362	HIS	-	expression tag	UNP Q6L545
F	363	HIS	-	expression tag	UNP Q6L545
F	364	HIS	=	expression tag	UNP Q6L545
F	365	HIS	-	expression tag	UNP Q6L545
F	366	HIS	-	expression tag	UNP Q6L545

 \bullet Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2).$




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 8 6 2	0	0
2	В	1	Total C O 8 6 2	0	0
2	С	1	Total C O 8 6 2	0	0
2	D	1	Total C O 8 6 2	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	E	1	Total C O 8 6 2	0	0
2	F	1	Total C O 8 6 2	0	0

 \bullet Molecule 3 is NITRATE ION (three-letter code: NO3) (formula: NO3).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total N O 4 1 3	0	0
3	A	1	Total N O 4 1 3	0	0
3	A	1	Total N O 4 1 3	0	0
3	В	1	Total N O 4 1 3	0	0
3	В	1	Total N O 4 1 3	0	0
3	В	1	Total N O 4 1 3	0	0
3	С	1	Total N O 4 1 3	0	0
3	С	1	Total N O 4 1 3	0	0
3	D	1	Total N O 4 1 3	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total N O 4 1 3	0	0
3	F	1	Total N O 4 1 3	0	0
3	F	1	Total N O 4 1 3	0	0


 \bullet Molecule 4 is GIBBERELLIN A3 (three-letter code: GA3) (formula: $\mathrm{C_{19}H_{22}O_6}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O	0	0
4	A	1	25 19 6	0	U
4	В	1	Total C O	0	0
4	Ъ	1	25 19 6	0	U
4	C	1	Total C O	0	0
4		1	25 19 6	U	U
4	D	1	Total C O	0	0
4	D	1	25 19 6	O	0
4	E	1	Total C O	0	0
4	Ш	1	25 19 6	O	0
4	F	1	Total C O	0	0
T	I.	1	25 19 6		

 \bullet Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total O P 5 4 1	0	0
5	E	1	Total O P 5 4 1	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	239	Total O 239 239	0	0
6	В	193	Total O 193 193	0	0
6	С	173	Total O 173 173	0	0
6	D	204	Total O 204 204	0	0
6	Е	123	Total O 123 123	0	0
6	F	174	Total O 174 174	0	0





3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Gibberellin receptor GID1









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	82.72Å 134.14Å 118.87Å	Donositor
a, b, c, α , β , γ	90.00° 105.20° 90.00°	Depositor
Resolution (Å)	20.00 - 1.90	Depositor
Resolution (A)	41.20 - 1.90	EDS
% Data completeness	100.0 (20.00-1.90)	Depositor
(in resolution range)	97.4 (41.20-1.90)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.53 (at 1.89Å)	Xtriage
Refinement program	REFMAC	Depositor
D D.	0.197 , 0.239	Depositor
R, R_{free}	0.199 , 0.242	DCC
R_{free} test set	9653 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.420	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 50.5	EDS
L-test for twinning ²	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15936	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, PO4, GA3, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.75	0/2593	0.70	0/3524
1	В	0.69	0/2561	0.67	0/3480
1	С	0.65	0/2440	0.67	0/3313
1	D	0.67	0/2533	0.67	0/3446
1	Е	0.58	0/2461	0.63	0/3352
1	F	0.64	0/2449	0.65	0/3328
All	All	0.66	0/15037	0.67	0/20443

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2510	0	2387	17	0
1	В	2476	0	2367	12	0
1	С	2365	0	2245	17	0
1	D	2455	0	2315	15	0
1	Е	2390	0	2242	18	0
1	F	2378	0	2246	8	0
2	A	8	0	14	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	8	0	14	2	0
2	С	8	0	14	0	0
2	D	8	0	14	0	0
2	Ε	8	0	14	1	0
2	F	8	0	14	1	0
3	A	12	0	0	1	0
3	В	12	0	0	1	0
3	С	8	0	0	2	0
3	D	8	0	0	0	0
3	F	8	0	0	0	0
4	A	25	0	21	0	0
4	В	25	0	21	0	0
4	С	25	0	21	0	0
4	D	25	0	21	0	0
4	Е	25	0	21	0	0
4	F	25	0	21	0	0
5	В	5	0	0	0	0
5	Е	5	0	0	0	0
6	A	239	0	0	1	0
6	В	193	0	0	1	0
6	С	173	0	0	1	0
6	D	204	0	0	1	0
6	Е	123	0	0	0	0
6	F	174	0	0	0	0
All	All	15936	0	14012	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:26:ASN:HD22	1:B:26:ASN:HD22	1.29	0.81
1:A:15:VAL:HG12	1:A:16:VAL:H	1.52	0.73
2:E:501:MPD:H13	2:F:501:MPD:H13	1.77	0.67
1:E:290:ILE:HG13	1:E:290:ILE:O	1.93	0.67
1:A:28[A]:LYS:HE3	1:A:245:PHE:O	1.98	0.63

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	321/365 (88%)	310 (97%)	11 (3%)	0	100	100
1	В	315/365 (86%)	303 (96%)	12 (4%)	0	100	100
1	С	298/365~(82%)	288 (97%)	10 (3%)	0	100	100
1	D	314/365 (86%)	305 (97%)	8 (2%)	1 (0%)	41	31
1	E	309/365~(85%)	297 (96%)	11 (4%)	1 (0%)	41	31
1	F	301/365 (82%)	291 (97%)	9 (3%)	1 (0%)	41	31
All	All	1858/2190 (85%)	1794 (97%)	61 (3%)	3 (0%)	47	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	72	GLN
1	Е	110	ALA
1	F	73	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	$_{ m ntiles}$
1	A	260/302~(86%)	258 (99%)	2 (1%)	81	82
1	В	$258/302\ (85\%)$	256 (99%)	2 (1%)	81	82
1	С	244/302 (81%)	243 (100%)	1 (0%)	91	91
1	D	251/302 (83%)	249 (99%)	2 (1%)	81	82



Continued from previous page...

Mol	Chain	Analysed	Rotameric Outliers		Percentiles	
1	E	$239/302 \ (79\%)$	236 (99%)	3 (1%)	69	68
1	F	241/302 (80%)	238 (99%)	3 (1%)	71	70
All	All	1493/1812 (82%)	1480 (99%)	13 (1%)	78	79

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	144	LEU
1	Е	231	THR
1	F	283	LEU
1	F	128	SER
1	F	177	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	19	HIS
1	F	68	HIS
1	С	26	ASN
1	С	179	GLN
1	D	68	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

26 ligands are modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / - 1	Т	Clasia	Das	T :1-	В	ond leng	$_{ m gths}$	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NO3	A	601	_	1,3,3	3.41	1 (100%)	0,3,3	-	-
5	PO4	В	701	-	4,4,4	0.86	0	6,6,6	0.46	0
3	NO3	D	603	-	1,3,3	3.57	1 (100%)	0,3,3	_	_
3	NO3	A	603	-	1,3,3	3.59	1 (100%)	0,3,3	-	-
4	GA3	D	401	-	28,29,29	1.31	3 (10%)	36,52,52	1.95	5 (13%)
3	NO3	В	601	-	1,3,3	3.83	1 (100%)	0,3,3	-	_
4	GA3	В	401	-	28,29,29	1.45	3 (10%)	36,52,52	1.94	8 (22%)
3	NO3	В	604	-	1,3,3	3.56	1 (100%)	0,3,3	-	-
3	NO3	С	602	-	1,3,3	3.56	1 (100%)	0,3,3	-	_
3	NO3	A	602	-	1,3,3	3.89	1 (100%)	0,3,3	-	_
2	MPD	D	501	-	7,7,7	0.47	0	9,10,10	0.40	0
3	NO3	С	604	-	1,3,3	3.40	1 (100%)	0,3,3	-	-
3	NO3	В	602	-	1,3,3	3.54	1 (100%)	0,3,3	-	-
3	NO3	F	603	-	1,3,3	3.54	1 (100%)	0,3,3	-	-
3	NO3	D	601	-	1,3,3	3.74	1 (100%)	0,3,3	-	_
5	PO4	Е	701	_	4,4,4	0.88	0	6,6,6	0.50	0
3	NO3	F	602	-	1,3,3	3.54	1 (100%)	0,3,3	-	-
2	MPD	F	501	-	7,7,7	0.27	0	9,10,10	0.51	0
4	GA3	F	401	_	28,29,29	1.23	1 (3%)	36,52,52	1.82	9 (25%)
2	MPD	С	501	-	7,7,7	0.36	0	9,10,10	0.49	0
4	GA3	С	401	-	28,29,29	1.15	1 (3%)	36,52,52	2.23	12 (33%)
2	MPD	A	501	-	7,7,7	0.42	0	9,10,10	0.60	0
2	MPD	В	501	-	7,7,7	0.34	0	9,10,10	0.44	0
2	MPD	Е	501	-	7,7,7	0.30	0	9,10,10	0.38	0
4	GA3	Е	401	-	28,29,29	1.35	3 (10%)	36,52,52	1.96	12 (33%)
4	GA3	A	401	_	28,29,29	1.14	1 (3%)	36,52,52	2.09	13 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	С	501	-	-	0/5/5/5	-
4	GA3	В	401	-	-	0/4/84/84	-
2	MPD	F	501	-	-	0/5/5/5	-
4	GA3	С	401	_	-	0/4/84/84	-
4	GA3	F	401	_	-	0/4/84/84	-
4	GA3	D	401	-	-	0/4/84/84	-
2	MPD	A	501	_	-	0/5/5/5	-
2	MPD	В	501	_	-	0/5/5/5	-
2	MPD	D	501	_	-	0/5/5/5	-
2	MPD	E	501	_	-	0/5/5/5	-
4	GA3	Е	401	_	-	2/4/84/84	-
4	GA3	A	401	-	-	1/4/84/84	-

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
4	В	401	GA3	O92-C19	4.89	1.46	1.36
4	Е	401	GA3	O92-C19	4.71	1.45	1.36
4	F	401	GA3	O92-C19	4.36	1.44	1.36
4	D	401	GA3	O92-C19	4.00	1.44	1.36
3	A	602	NO3	O1-N	3.89	1.42	1.24

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
4	С	401	GA3	C15-C16-C13	7.41	113.72	107.58
4	D	401	GA3	C14-C13-C16	-7.01	98.30	102.09
4	С	401	GA3	C14-C13-C16	-6.90	98.36	102.09
4	D	401	GA3	C15-C16-C13	6.48	112.95	107.58
4	В	401	GA3	C15-C16-C13	6.38	112.87	107.58

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	401	GA3	C5-C6-C7-O71
4	Е	401	GA3	C5-C6-C7-O71
4	Е	401	GA3	C5-C6-C7-O72

There are no ring outliers.


8 monomers are involved in 7 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	604	NO3	1	0
3	С	602	NO3	1	0
3	A	602	NO3	1	0
3	С	604	NO3	1	0
2	F	501	MPD	1	0
2	A	501	MPD	1	0
2	В	501	MPD	2	0
2	Е	501	MPD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	A	321/365~(87%)	-0.21	8 (2%) 57 60	15, 24, 42, 54	0
1	В	316/365~(86%)	0.04	22 (6%) 16 18	16, 28, 49, 66	0
1	С	301/365~(82%)	-0.05	7 (2%) 60 63	18, 31, 48, 57	0
1	D	315/365~(86%)	-0.03	10 (3%) 47 50	16, 28, 48, 56	0
1	E	313/365~(85%)	0.42	23 (7%) 15 16	22, 39, 56, 68	0
1	F	305/365~(83%)	-0.19	6 (1%) 65 68	19, 31, 50, 58	0
All	All	1871/2190 (85%)	-0.00	76 (4%) 37 40	15, 30, 51, 68	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	107	ALA	6.1
1	Е	110	ALA	5.7
1	В	14	THR	5.4
1	Е	109	ALA	4.7
1	В	107	ALA	4.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q<0.9
3	NO3	A	602	4/4	0.77	0.19	42,43,44,45	0
3	NO3	D	601	4/4	0.83	0.14	61,61,61,61	0
5	PO4	В	701	5/5	0.84	0.18	89,89,89,90	0
3	NO3	A	601	4/4	0.86	0.18	45,45,46,46	0
3	NO3	В	601	4/4	0.87	0.19	49,50,50,51	0
5	PO4	Е	701	5/5	0.88	0.22	97,98,98,98	0
3	NO3	С	602	4/4	0.90	0.26	42,43,43,43	0
3	NO3	F	602	4/4	0.91	0.27	51,51,51,53	0
3	NO3	В	604	4/4	0.92	0.20	50,51,51,51	0
2	MPD	D	501	8/8	0.92	0.15	34,38,38,40	0
3	NO3	A	603	4/4	0.92	0.14	61,61,62,62	0
3	NO3	С	604	4/4	0.92	0.10	44,45,45,46	0
2	MPD	F	501	8/8	0.93	0.10	32,34,35,35	0
3	NO3	F	603	4/4	0.93	0.09	62,62,62,62	0
2	MPD	В	501	8/8	0.93	0.12	33,36,36,37	0
2	MPD	A	501	8/8	0.93	0.11	28,31,37,37	0
3	NO3	D	603	4/4	0.94	0.11	51,51,51,52	0
4	GA3	Е	401	25/25	0.94	0.09	24,28,30,32	0
2	MPD	С	501	8/8	0.95	0.13	31,33,35,35	0
2	MPD	Е	501	8/8	0.95	0.09	38,41,42,43	0
3	NO3	В	602	4/4	0.95	0.18	45,46,46,46	0
4	GA3	F	401	25/25	0.96	0.07	20,23,26,27	0
4	GA3	С	401	25/25	0.97	0.08	16,19,22,23	0
4	GA3	D	401	25/25	0.97	0.10	19,21,23,26	0
4	GA3	В	401	25/25	0.97	0.10	15,17,21,21	0
4	GA3	A	401	25/25	0.98	0.07	15,17,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


























6.5 Other polymers (i)

There are no such residues in this entry.

