



wwPDB X-ray Structure Validation Summary Report ⓘ

May 24, 2020 – 10:50 pm BST

PDB ID : 2EHO
Title : Crystal structure of human GINS complex
Authors : Choi, J.M.; Lim, H.S.; Kim, J.J.; Song, O.K.; Cho, Y.
Deposited on : 2007-03-07
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	203	 27% 47% 5% 21%
1	E	203	 43% 45% 5% 6%
1	I	203	 41% 45% 6% 5%
2	B	152	 41% 45% 9% 5%
2	F	152	 49% 42% 5% 4%
2	J	152	 50% 38% 7% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	186	<p>%</p> <p>26% 57% 6% 10%</p>
3	G	186	<p>%</p> <p>35% 53% 7% 5%</p>
3	K	186	<p>40% 47% 8% 6%</p>
4	D	216	<p>%</p> <p>31% 44% 9% 16%</p>
4	H	216	<p>41% 35% 8% 15%</p>
4	L	216	<p>37% 38% 8% 16%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	B	1016	-	-	X	-
5	SO4	F	1010	-	-	X	-
5	SO4	K	1007	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 16937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GINS complex subunit 4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	161	Total 1340	C 857	N 227	O 246	S 2	Se 8	0	0	0
1	E	190	Total 1583	C 1005	N 269	O 299	S 2	Se 8	0	0	0
1	I	192	Total 1604	C 1016	N 278	O 300	S 2	Se 8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
A	55	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
A	61	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
A	80	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
A	82	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
A	98	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
A	134	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
A	149	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
E	35	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
E	55	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
E	61	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
E	80	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
E	82	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
E	98	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
E	134	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
E	149	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
I	35	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
I	55	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
I	61	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
I	80	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
I	82	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
I	98	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9
I	134	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	149	MSE	MET	MODIFIED RESIDUE	UNP Q9BRT9

- Molecule 2 is a protein called DNA replication complex GINS protein PSF1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	B	145	Total	C	N	O	S	Se	0	0	0
			1188	744	217	217	3	7			
2	F	146	Total	C	N	O	S	Se	0	0	0
			1196	749	218	218	3	8			
2	J	145	Total	C	N	O	S	Se	0	0	0
			1188	744	217	217	3	7			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MSE	-	CLONING ARTIFACT	UNP Q14691
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q14691
B	7	MSE	MET	MODIFIED RESIDUE	UNP Q14691
B	36	MSE	MET	MODIFIED RESIDUE	UNP Q14691
B	97	ILE	VAL	SEE REMARK 999	UNP Q14691
B	106	MSE	MET	MODIFIED RESIDUE	UNP Q14691
B	111	MSE	MET	MODIFIED RESIDUE	UNP Q14691
B	125	MSE	MET	MODIFIED RESIDUE	UNP Q14691
B	140	MSE	MET	MODIFIED RESIDUE	UNP Q14691
F	0	MSE	-	CLONING ARTIFACT	UNP Q14691
F	1	MSE	MET	MODIFIED RESIDUE	UNP Q14691
F	7	MSE	MET	MODIFIED RESIDUE	UNP Q14691
F	36	MSE	MET	MODIFIED RESIDUE	UNP Q14691
F	97	ILE	VAL	SEE REMARK 999	UNP Q14691
F	106	MSE	MET	MODIFIED RESIDUE	UNP Q14691
F	111	MSE	MET	MODIFIED RESIDUE	UNP Q14691
F	125	MSE	MET	MODIFIED RESIDUE	UNP Q14691
F	140	MSE	MET	MODIFIED RESIDUE	UNP Q14691
J	0	MSE	-	CLONING ARTIFACT	UNP Q14691
J	1	MSE	MET	MODIFIED RESIDUE	UNP Q14691
J	7	MSE	MET	MODIFIED RESIDUE	UNP Q14691
J	36	MSE	MET	MODIFIED RESIDUE	UNP Q14691
J	97	ILE	VAL	SEE REMARK 999	UNP Q14691
J	106	MSE	MET	MODIFIED RESIDUE	UNP Q14691
J	111	MSE	MET	MODIFIED RESIDUE	UNP Q14691
J	125	MSE	MET	MODIFIED RESIDUE	UNP Q14691
J	140	MSE	MET	MODIFIED RESIDUE	UNP Q14691

- Molecule 3 is a protein called DNA replication complex GINS protein PSF2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	168	Total	C	N	O	S	Se	0	0	0
			1369	876	230	253	1	9			
3	G	176	Total	C	N	O	S	Se	0	0	0
			1432	916	242	264	1	9			
3	K	175	Total	C	N	O	S	Se	0	0	0
			1423	911	240	262	1	9			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MSE	-	CLONING ARTIFACT	UNP Q9Y248
C	1	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
C	66	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
C	74	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
C	87	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
C	93	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
C	120	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
C	151	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
C	167	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
G	0	MSE	-	CLONING ARTIFACT	UNP Q9Y248
G	1	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
G	66	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
G	74	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
G	87	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
G	93	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
G	120	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
G	151	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
G	167	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
K	0	MSE	-	CLONING ARTIFACT	UNP Q9Y248
K	1	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
K	66	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
K	74	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
K	87	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
K	93	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
K	120	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
K	151	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248
K	167	MSE	MET	MODIFIED RESIDUE	UNP Q9Y248


- Molecule 4 is a protein called GINS complex subunit 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
4	D	182	Total 1462	C 931	N 251	O 274	S 1	Se 5	0	0	0
4	H	183	Total 1470	C 935	N 253	O 276	S 1	Se 5	0	0	0
4	L	182	Total 1462	C 931	N 252	O 273	S 1	Se 5	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
D	27	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
D	40	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
D	112	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
D	150	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
D	171	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
H	1	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
H	27	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
H	40	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
H	112	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
H	150	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
H	171	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
L	1	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
L	27	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
L	40	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
L	112	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
L	150	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5
L	171	MSE	MET	MODIFIED RESIDUE	UNP Q9BRX5

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	I	1	Total	O	S	0	0
			5	4	1		
5	I	1	Total	O	S	0	0
			5	4	1		
5	I	1	Total	O	S	0	0
			5	4	1		
5	J	1	Total	O	S	0	0
			5	4	1		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	K	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	8	Total	O	0	0
			8	8		
6	B	5	Total	O	0	0
			5	5		
6	C	2	Total	O	0	0
			2	2		
6	D	1	Total	O	0	0
			1	1		
6	E	22	Total	O	0	0
			22	22		
6	F	11	Total	O	0	0
			11	11		
6	G	6	Total	O	0	0
			6	6		
6	H	7	Total	O	0	0
			7	7		
6	I	18	Total	O	0	0
			18	18		
6	J	23	Total	O	0	0
			23	23		
6	K	13	Total	O	0	0
			13	13		
6	L	19	Total	O	0	0
			19	19		




• Molecule 2: DNA replication complex GINS protein PSF1




• Molecule 2: DNA replication complex GINS protein PSF1




• Molecule 2: DNA replication complex GINS protein PSF1




• Molecule 3: DNA replication complex GINS protein PSF2







• Molecule 3: DNA replication complex GINS protein PSF2



• Molecule 3: DNA replication complex GINS protein PSF2




• Molecule 4: GINS complex subunit 3




• Molecule 4: GINS complex subunit 3





- Molecule 4: GINS complex subunit 3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	164.94Å 177.00Å 126.51Å 90.00° 105.91° 90.00°	Depositor
Resolution (Å)	29.90 – 3.00 50.08 – 2.81	Depositor EDS
% Data completeness (in resolution range)	91.9 (29.90-3.00) 96.0 (50.08-2.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.01	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.81Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.221 , 0.274 0.216 , 0.266	Depositor DCC
R_{free} test set	4073 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	52.9	Xtrriage
Anisotropy	0.240	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 66.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	16937	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1359	0.56	1/1814 (0.1%)
1	E	0.42	0/1606	0.67	1/2150 (0.0%)
1	I	0.43	0/1627	0.85	9/2177 (0.4%)
2	B	0.37	0/1205	0.60	0/1608
2	F	0.42	0/1212	0.62	0/1615
2	J	0.44	0/1205	0.64	0/1608
3	C	0.32	0/1389	0.58	0/1866
3	G	0.37	0/1454	0.59	0/1955
3	K	0.42	0/1445	0.64	0/1943
4	D	0.35	0/1491	0.59	0/2003
4	H	0.46	0/1499	0.70	0/2014
4	L	0.76	4/1491 (0.3%)	1.04	12/2003 (0.6%)
All	All	0.44	4/16983 (0.0%)	0.69	23/22756 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	L	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	L	58	ASN	C-O	18.93	1.59	1.23
4	L	60	VAL	C-O	10.76	1.43	1.23
4	L	41	PRO	N-CD	8.00	1.59	1.47
4	L	38	THR	C-O	-6.51	1.10	1.23

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	L	38	THR	O-C-N	-15.98	97.13	122.70
4	L	60	VAL	O-C-N	13.99	147.68	121.10
4	L	60	VAL	CA-C-O	-12.51	93.82	120.10
1	I	211	ILE	CB-CA-C	-12.32	86.97	111.60
4	L	38	THR	CA-C-O	11.87	145.02	120.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	L	58	ASN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1340	0	1353	130	0
1	E	1583	0	1587	118	0
1	I	1604	0	1616	134	0
2	B	1188	0	1180	110	0
2	F	1196	0	1189	83	0
2	J	1188	0	1180	81	0
3	C	1369	0	1391	153	0
3	G	1432	0	1458	112	0
3	K	1423	0	1450	117	0
4	D	1462	0	1418	126	0
4	H	1470	0	1424	115	0
4	L	1462	0	1420	115	0
5	A	5	0	0	0	0
5	B	5	0	0	5	0
5	E	20	0	0	0	0
5	F	20	0	0	5	0
5	I	15	0	0	1	0
5	J	5	0	0	0	0
5	K	5	0	0	0	0
5	L	10	0	0	0	0
6	A	8	0	0	2	0
6	B	5	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	2	0	0	0	0
6	D	1	0	0	0	0
6	E	22	0	0	5	0
6	F	11	0	0	1	0
6	G	6	0	0	0	0
6	H	7	0	0	0	0
6	I	18	0	0	6	0
6	J	23	0	0	1	0
6	K	13	0	0	2	0
6	L	19	0	0	5	0
All	All	16937	0	16666	1234	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

The worst 5 of 1234 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:211:ILE:CG2	1:I:211:ILE:O	1.92	1.18
1:I:144:VAL:HG22	2:J:109:GLU:HG3	1.29	1.13
4:L:13:LEU:HD21	4:L:171:MSE:HE2	1.27	1.11
1:I:149:MSE:HE3	2:J:36:MSE:HE1	1.15	1.10
4:L:90:LYS:HE3	4:L:90:LYS:H	1.12	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	155/203 (76%)	119 (77%)	28 (18%)	8 (5%)	2 12
1	E	186/203 (92%)	162 (87%)	20 (11%)	4 (2%)	6 31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	190/203 (94%)	159 (84%)	25 (13%)	6 (3%)	4	22
2	B	143/152 (94%)	117 (82%)	18 (13%)	8 (6%)	2	10
2	F	144/152 (95%)	127 (88%)	10 (7%)	7 (5%)	2	13
2	J	143/152 (94%)	125 (87%)	14 (10%)	4 (3%)	5	25
3	C	164/186 (88%)	126 (77%)	30 (18%)	8 (5%)	2	13
3	G	174/186 (94%)	148 (85%)	19 (11%)	7 (4%)	3	17
3	K	173/186 (93%)	152 (88%)	17 (10%)	4 (2%)	6	30
4	D	178/216 (82%)	136 (76%)	29 (16%)	13 (7%)	1	5
4	H	179/216 (83%)	160 (89%)	14 (8%)	5 (3%)	5	25
4	L	178/216 (82%)	150 (84%)	18 (10%)	10 (6%)	2	10
All	All	2007/2271 (88%)	1681 (84%)	242 (12%)	84 (4%)	3	16

5 of 84 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	LEU
1	A	72	ASP
1	A	116	GLU
2	B	17	PRO
3	C	68	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	151/181 (83%)	143 (95%)	8 (5%)	22	58
1	E	179/181 (99%)	167 (93%)	12 (7%)	16	49
1	I	180/181 (99%)	164 (91%)	16 (9%)	9	35
2	B	126/125 (101%)	114 (90%)	12 (10%)	8	32
2	F	127/125 (102%)	120 (94%)	7 (6%)	21	57
2	J	126/125 (101%)	116 (92%)	10 (8%)	12	41

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	154/161 (96%)	140 (91%)	14 (9%)	9	34
3	G	160/161 (99%)	148 (92%)	12 (8%)	13	43
3	K	159/161 (99%)	140 (88%)	19 (12%)	5	22
4	D	157/180 (87%)	144 (92%)	13 (8%)	11	39
4	H	158/180 (88%)	134 (85%)	24 (15%)	3	14
4	L	157/180 (87%)	137 (87%)	20 (13%)	4	19
All	All	1834/1941 (94%)	1667 (91%)	167 (9%)	9	34

5 of 167 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	153	ILE
4	H	139	LEU
4	L	90	LYS
3	G	175	GLN
4	H	57	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 80 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	106	ASN
4	H	93	GLN
4	L	122	GLN
3	G	139	GLN
4	H	18	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	F	1014	-	4,4,4	0.32	0	6,6,6	0.12	0
5	SO4	I	1001	-	4,4,4	0.27	0	6,6,6	0.10	0
5	SO4	F	1010	-	4,4,4	0.27	0	6,6,6	0.13	0
5	SO4	E	1015	-	4,4,4	0.34	0	6,6,6	0.09	0
5	SO4	K	1007	-	4,4,4	0.35	0	6,6,6	0.18	0
5	SO4	I	1003	-	4,4,4	0.28	0	6,6,6	0.07	0
5	SO4	E	1011	-	4,4,4	0.32	0	6,6,6	0.06	0
5	SO4	L	1008	-	4,4,4	0.35	0	6,6,6	0.06	0
5	SO4	F	1005	-	4,4,4	0.30	0	6,6,6	0.05	0
5	SO4	L	1009	-	4,4,4	0.32	0	6,6,6	0.12	0
5	SO4	B	1016	-	4,4,4	0.31	0	6,6,6	0.18	0
5	SO4	A	1017	-	4,4,4	0.33	0	6,6,6	0.09	0
5	SO4	F	1004	-	4,4,4	0.29	0	6,6,6	0.10	0
5	SO4	J	1006	-	4,4,4	0.33	0	6,6,6	0.05	0
5	SO4	E	1013	-	4,4,4	0.33	0	6,6,6	0.08	0
5	SO4	I	1002	-	4,4,4	0.27	0	6,6,6	0.09	0
5	SO4	E	1012	-	4,4,4	0.34	0	6,6,6	0.10	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	1014	SO4	1	0
5	I	1001	SO4	1	0
5	F	1010	SO4	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1016	SO4	5	0
5	F	1004	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	153/203 (75%)	-0.06	2 (1%) 77 51	48, 77, 108, 115	0
1	E	182/203 (89%)	-0.51	1 (0%) 91 75	21, 47, 79, 108	0
1	I	184/203 (90%)	-0.38	1 (0%) 91 75	21, 44, 87, 97	0
2	B	138/152 (90%)	-0.12	2 (1%) 75 49	33, 65, 92, 106	0
2	F	138/152 (90%)	-0.29	0 100 100	18, 44, 77, 98	0
2	J	138/152 (90%)	-0.42	0 100 100	19, 40, 69, 86	0
3	C	159/186 (85%)	-0.03	1 (0%) 89 72	43, 78, 115, 123	0
3	G	167/186 (89%)	-0.21	1 (0%) 89 72	22, 65, 103, 112	0
3	K	166/186 (89%)	-0.56	0 100 100	22, 41, 63, 72	0
4	D	177/216 (81%)	0.11	3 (1%) 70 41	40, 83, 104, 113	0
4	H	178/216 (82%)	-0.50	0 100 100	18, 40, 76, 88	0
4	L	177/216 (81%)	-0.41	0 100 100	19, 43, 85, 97	0
All	All	1957/2271 (86%)	-0.29	11 (0%) 89 72	18, 54, 101, 123	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	65	LEU	4.4
4	D	49	GLU	3.8
4	D	160	ASP	3.2
1	I	69	LYS	2.9
2	B	131	ASP	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	K	1007	5/5	0.73	0.69	114,114,115,116	0
5	SO4	A	1017	5/5	0.75	0.21	115,115,116,116	0
5	SO4	F	1004	5/5	0.76	0.31	137,138,138,138	0
5	SO4	L	1008	5/5	0.79	0.35	118,118,118,119	0
5	SO4	I	1001	5/5	0.79	0.20	124,124,125,125	0
5	SO4	L	1009	5/5	0.83	0.35	102,102,102,103	0
5	SO4	E	1015	5/5	0.83	0.22	125,125,126,126	0
5	SO4	I	1003	5/5	0.86	0.25	137,137,138,138	0
5	SO4	F	1005	5/5	0.87	0.21	111,112,112,112	0
5	SO4	E	1012	5/5	0.88	0.18	107,107,108,108	0
5	SO4	F	1014	5/5	0.89	0.18	115,116,116,116	0
5	SO4	F	1010	5/5	0.90	0.29	106,107,107,107	0
5	SO4	E	1011	5/5	0.90	0.30	107,107,108,108	0
5	SO4	E	1013	5/5	0.91	0.25	117,118,118,118	0
5	SO4	B	1016	5/5	0.91	0.29	95,97,98,98	0
5	SO4	I	1002	5/5	0.92	0.14	95,96,96,97	0
5	SO4	J	1006	5/5	0.94	0.35	96,96,96,97	0

6.5 Other polymers [i](#)

There are no such residues in this entry.