



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 18, 2024 – 08:27 AM EST

PDB ID : 4EJ3  
Title : Crystal structure of a CRISPR associated protein from *Thermus thermophilus* HB8  
Authors : Huang, Q.Q.  
Deposited on : 2012-04-06  
Resolution : 2.52 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	514	 4% 78% 12% • 9%
1	B	514	 % 79% 11% • 9%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR associated protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	467	3693	2370	659	659	5	0	0	0
1	B	467	3695	2371	662	657	5	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP Q53VY1
A	-10	ARG	-	expression tag	UNP Q53VY1
A	-9	GLY	-	expression tag	UNP Q53VY1
A	-8	SER	-	expression tag	UNP Q53VY1
A	-7	HIS	-	expression tag	UNP Q53VY1
A	-6	HIS	-	expression tag	UNP Q53VY1
A	-5	HIS	-	expression tag	UNP Q53VY1
A	-4	HIS	-	expression tag	UNP Q53VY1
A	-3	HIS	-	expression tag	UNP Q53VY1
A	-2	HIS	-	expression tag	UNP Q53VY1
A	-1	GLY	-	expression tag	UNP Q53VY1
A	0	SER	-	expression tag	UNP Q53VY1
B	-11	MET	-	expression tag	UNP Q53VY1
B	-10	ARG	-	expression tag	UNP Q53VY1
B	-9	GLY	-	expression tag	UNP Q53VY1
B	-8	SER	-	expression tag	UNP Q53VY1
B	-7	HIS	-	expression tag	UNP Q53VY1
B	-6	HIS	-	expression tag	UNP Q53VY1
B	-5	HIS	-	expression tag	UNP Q53VY1
B	-4	HIS	-	expression tag	UNP Q53VY1
B	-3	HIS	-	expression tag	UNP Q53VY1
B	-2	HIS	-	expression tag	UNP Q53VY1
B	-1	GLY	-	expression tag	UNP Q53VY1
B	0	SER	-	expression tag	UNP Q53VY1


- Molecule 2 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	A	61	Total 61	O 61	0	0
2	B	96	Total 96	O 96	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CRISPR associated protein



- Molecule 1: CRISPR associated protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.78Å 47.83Å 129.18Å 90.00° 97.34° 90.00°	Depositor
Resolution (Å)	49.86 – 2.52 49.86 – 2.52	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.86-2.52) 94.8 (49.86-2.52)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.46 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.218 , 0.277 0.215 , 0.272	Depositor DCC
$R_{free}$ test set	2000 reflections (5.42%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.4	Xtrriage
Anisotropy	0.305	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 34.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7545	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/3784	0.58	0/5141
1	B	0.48	0/3786	0.64	0/5143
All	All	0.46	0/7570	0.61	0/10284

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3693	0	3706	40	0
1	B	3695	0	3713	46	0
2	A	61	0	0	4	0
2	B	96	0	0	4	0
All	All	7545	0	7419	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:THR:HG23	1:B:272:THR:HG22	1.42	0.99

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:THR:HG23	1:B:272:THR:CG2	2.11	0.80
1:B:228:ALA:HB1	1:B:267:GLU:HG3	1.63	0.79
1:A:33:LEU:HD22	1:A:84:GLN:HG3	1.65	0.79
1:A:191:THR:HG21	1:A:340:ARG:HD3	1.75	0.66
1:B:33:LEU:HD22	1:B:84:GLN:HG3	1.80	0.63
1:B:325:GLN:HB2	1:B:337:ILE:HD11	1.80	0.63
1:A:413:LEU:HD12	1:A:416:LEU:HD12	1.80	0.62
1:B:228:ALA:HB1	1:B:267:GLU:CG	2.28	0.61
1:B:424:PRO:HB2	1:B:428:LEU:HD12	1.83	0.60
1:A:395:VAL:HG13	1:A:481:ALA:HB1	1.83	0.59
1:A:217:VAL:HG13	1:A:218:PRO:HD2	1.84	0.59
1:B:448:ALA:O	1:B:452:LEU:HD12	2.02	0.58
1:B:413:LEU:HD23	1:B:416:LEU:HD12	1.84	0.58
1:A:55:LEU:HD22	1:A:158:LEU:HD11	1.85	0.58
1:A:341:VAL:C	1:A:342:LEU:HD12	2.23	0.57
1:A:64:LEU:HD13	1:A:76:TRP:CH2	2.40	0.57
1:A:228:ALA:HB1	1:A:267:GLU:HG3	1.85	0.57
1:B:193:GLN:HG3	1:B:336:ARG:CZ	2.35	0.56
1:A:337:ILE:CG2	1:A:361:LEU:HD13	2.34	0.56
1:B:274:ARG:NH1	2:B:603:HOH:O	2.39	0.56
1:B:209:GLU:HG3	2:B:676:HOH:O	2.04	0.55
1:B:496:LEU:HD12	1:B:497:GLY:N	2.22	0.55
1:A:191:THR:HG21	1:A:340:ARG:CD	2.38	0.54
1:B:174:VAL:HG21	1:B:264:PRO:HG3	1.88	0.54
1:A:307:LEU:HD11	1:A:374:LEU:HD23	1.90	0.54
1:B:182:VAL:HG11	1:B:187:LEU:HD11	1.89	0.53
1:A:454:ARG:HD2	2:A:656:HOH:O	2.09	0.52
1:B:9:LEU:HD22	1:B:14:TRP:CE2	2.44	0.52
1:B:182:VAL:CG1	1:B:182:VAL:O	2.58	0.52
1:B:217:VAL:HG13	1:B:218:PRO:HD2	1.91	0.52
1:B:182:VAL:CG1	1:B:187:LEU:HD11	2.40	0.52
1:B:496:LEU:HD12	1:B:497:GLY:H	1.75	0.51
1:A:337:ILE:HG21	1:A:361:LEU:HD13	1.91	0.51
1:B:49:GLU:OE1	1:B:356:ARG:NH1	2.43	0.51
1:B:182:VAL:O	1:B:182:VAL:HG13	2.11	0.51
1:A:49:GLU:OE1	1:A:356:ARG:NH1	2.43	0.51
1:B:232:TRP:CD2	1:B:266:VAL:HG11	2.45	0.50
1:A:12:GLU:HG3	1:A:13:PRO:HD2	1.92	0.50
1:A:113:LEU:O	1:A:249:ARG:NH1	2.44	0.50
1:B:182:VAL:HG22	1:B:187:LEU:HD21	1.92	0.50
1:A:270:GLU:HA	2:A:646:HOH:O	2.11	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:325:GLN:HB2	1:B:337:ILE:CD1	2.43	0.49
1:A:46:PRO:HD2	1:A:354:ILE:HD12	1.94	0.48
1:B:61:HIS:CE1	1:B:203:LEU:HD23	2.47	0.48
1:B:291:LEU:HD11	1:B:306:MET:HB3	1.96	0.48
1:B:391:LEU:HD22	1:B:425:LEU:CD2	2.43	0.48
1:A:342:LEU:HD12	1:A:342:LEU:N	2.29	0.47
1:B:296:GLU:N	1:B:296:GLU:OE1	2.47	0.47
1:A:232:TRP:CD2	1:A:266:VAL:HG11	2.48	0.47
1:B:234:LEU:HD12	1:B:243:TRP:CZ3	2.49	0.47
1:A:182:VAL:HG13	1:A:182:VAL:O	2.15	0.47
1:A:452:LEU:HD11	2:A:654:HOH:O	2.15	0.47
1:A:380:MET:HB3	1:A:380:MET:HE2	1.80	0.46
1:B:55:LEU:O	1:B:58:ALA:HB3	2.15	0.46
1:A:324:LEU:O	1:A:328:LEU:HG	2.16	0.46
1:A:182:VAL:HG22	1:A:187:LEU:HD21	1.97	0.46
1:A:395:VAL:O	1:A:399:VAL:HG23	2.16	0.46
1:A:275:ASP:HB3	1:A:278:VAL:HG23	1.97	0.45
1:B:61:HIS:CE1	1:B:203:LEU:CD2	2.99	0.45
1:B:336:ARG:HA	2:B:660:HOH:O	2.15	0.45
1:B:145:LEU:HD12	1:B:146:PRO:HD2	1.99	0.45
1:B:282:LEU:HD23	1:B:288:LEU:HD23	1.99	0.45
1:A:230:THR:HG22	1:A:266:VAL:HG13	1.98	0.44
1:B:293:LEU:HD13	1:B:355:ARG:HB2	1.98	0.44
1:A:419:LEU:O	1:A:423:LEU:HD13	2.17	0.44
1:A:82:PHE:HB2	1:A:83:PRO:HD2	2.00	0.43
1:A:7:PHE:CE1	1:A:12:GLU:HG2	2.53	0.43
1:A:7:PHE:CD1	1:A:12:GLU:HG2	2.53	0.43
1:A:236:GLY:O	1:A:240:VAL:HG23	2.19	0.43
1:B:174:VAL:CG2	1:B:264:PRO:HG3	2.49	0.43
1:A:145:LEU:HD12	1:A:146:PRO:HD2	2.01	0.42
1:B:272:THR:O	1:B:272:THR:OG1	2.31	0.42
1:B:399:VAL:HG11	1:B:419:LEU:HD11	2.00	0.42
1:A:47:LEU:HD22	1:A:163:PHE:CE1	2.55	0.42
1:B:19:LYS:HE3	1:B:37:GLU:HG2	2.01	0.42
1:B:469:ARG:HG2	1:B:482:LEU:HD11	2.01	0.41
1:A:125:LEU:HD23	2:A:634:HOH:O	2.19	0.41
1:B:207:THR:HB	2:B:676:HOH:O	2.19	0.41
1:B:391:LEU:HD23	1:B:391:LEU:C	2.40	0.41
1:A:304:SER:CB	1:A:378:LEU:HD12	2.51	0.41
1:B:11:ASP:OD1	1:B:95:ARG:NH2	2.50	0.41
1:B:94:PHE:O	1:B:95:ARG:C	2.58	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:234:LEU:HD12	1:B:243:TRP:CE3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	461/514 (90%)	447 (97%)	14 (3%)	0	100	100
1	B	461/514 (90%)	438 (95%)	22 (5%)	1 (0%)	47	67
All	All	922/1028 (90%)	885 (96%)	36 (4%)	1 (0%)	51	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	95	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/418 (90%)	363 (96%)	13 (4%)	36	60
1	B	376/418 (90%)	366 (97%)	10 (3%)	44	69
All	All	752/836 (90%)	729 (97%)	23 (3%)	40	65

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	CYS
1	A	82	PHE
1	A	84	GLN
1	A	92	ASN
1	A	176	SER
1	A	182	VAL
1	A	314	VAL
1	A	369	LYS
1	A	413	LEU
1	A	419	LEU
1	A	446	GLU
1	A	451	ASP
1	A	491	ARG
1	B	5	GLU
1	B	37	GLU
1	B	84	GLN
1	B	217	VAL
1	B	249	ARG
1	B	348	GLN
1	B	419	LEU
1	B	439	ARG
1	B	491	ARG
1	B	496	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	373	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	467/514 (90%)	0.27	21 (4%) 33 36	28, 48, 69, 86	0
1	B	467/514 (90%)	-0.01	3 (0%) 89 90	21, 38, 54, 67	0
All	All	934/1028 (90%)	0.13	24 (2%) 56 59	21, 42, 66, 86	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	496	LEU	5.3
1	A	497	GLY	4.3
1	A	144	ASN	3.3
1	A	417	THR	3.1
1	A	364	GLY	3.0
1	A	330	ASP	2.9
1	A	451	ASP	2.7
1	A	452	LEU	2.7
1	A	495	GLU	2.7
1	A	493	LEU	2.6
1	B	336	ARG	2.6
1	A	455	GLU	2.6
1	A	367	THR	2.5
1	A	384	LEU	2.4
1	A	477	ARG	2.4
1	A	472	LEU	2.4
1	A	494	GLY	2.3
1	A	418	LYS	2.2
1	B	451	ASP	2.2
1	A	484	GLN	2.2
1	B	497	GLY	2.1
1	A	253	GLU	2.1
1	A	173	GLY	2.0
1	A	446	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.