



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2023 – 01:13 AM EDT

PDB ID : 8ELF
Title : Structure of Get3d, a homolog of Get3, from Arabidopsis thaliana
Authors : Manu, M.S.; Barlow, A.N.; Clemons Jr., W.M.; Ramasamy, S.
Deposited on : 2022-09-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	
1	B	407	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12253 atoms, of which 6141 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein At1g26090, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	381	6004	1894	3036	508	550	16	0	1	0
1	B	384	6033	1904	3046	511	556	16	0	1	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	MET	-	initiating methionine	UNP Q6DYE4
A	456	LEU	-	expression tag	UNP Q6DYE4
A	457	GLU	-	expression tag	UNP Q6DYE4
A	458	HIS	-	expression tag	UNP Q6DYE4
A	459	HIS	-	expression tag	UNP Q6DYE4
A	460	HIS	-	expression tag	UNP Q6DYE4
A	461	HIS	-	expression tag	UNP Q6DYE4
A	462	HIS	-	expression tag	UNP Q6DYE4
A	463	HIS	-	expression tag	UNP Q6DYE4
B	57	MET	-	initiating methionine	UNP Q6DYE4
B	456	LEU	-	expression tag	UNP Q6DYE4
B	457	GLU	-	expression tag	UNP Q6DYE4
B	458	HIS	-	expression tag	UNP Q6DYE4
B	459	HIS	-	expression tag	UNP Q6DYE4
B	460	HIS	-	expression tag	UNP Q6DYE4
B	461	HIS	-	expression tag	UNP Q6DYE4
B	462	HIS	-	expression tag	UNP Q6DYE4
B	463	HIS	-	expression tag	UNP Q6DYE4

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is 1,2-DIMYRISTOYL-SN-GLYCERO-3-PHOSPHATE (three-letter code: XP4) (formula: C₃₁H₆₀O₈P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	P		
4	B	1	99	31	59	8	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	59	Total	O	0	0
			59	59		
5	B	41	Total	O	0	0
			41	41		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein At1g26090, chloroplastic



- Molecule 1: Uncharacterized protein At1g26090, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.25Å 67.05Å 99.40Å 90.00° 97.81° 90.00°	Depositor
Resolution (Å)	32.83 – 2.00 58.70 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (32.83-2.00) 99.5 (58.70-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.213 , 0.256 0.215 , 0.256	Depositor DCC
R_{free} test set	2559 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtrriage
Anisotropy	0.201	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12253	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PO4, XP4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3021	0.49	0/4075
1	B	0.26	0/3042	0.49	0/4106
All	All	0.26	0/6063	0.49	0/8181

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2968	3036	3034	24	0
1	B	2987	3046	3045	26	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	B	40	59	60	1	0
5	A	59	0	0	0	0
5	B	41	0	0	0	0
All	All	6112	6141	6139	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:THR:HG21	1:A:274:LEU:HD22	1.65	0.79
1:B:397:THR:HG23	1:B:451:ILE:HD13	1.65	0.77
1:B:288:ARG:HD3	1:B:376:THR:HG22	1.73	0.70
1:B:284:PRO:O	1:B:288:ARG:NH1	2.29	0.65
1:A:265:ALA:O	1:A:269:THR:HG23	1.98	0.63
1:B:272:ARG:O	1:B:276:THR:HG23	2.02	0.59
1:A:394:LYS:NZ	1:A:435:GLN:O	2.37	0.57
1:A:337:ASP:O	1:A:448[B]:ARG:NH1	2.36	0.57
1:A:279:SER:HA	1:A:282:ARG:NH2	2.20	0.56
1:B:178:VAL:HG11	1:B:180:PHE:CZ	2.41	0.55
1:A:178:VAL:HG11	1:A:180:PHE:CE2	2.42	0.54
1:A:279:SER:OG	1:A:282:ARG:NH2	2.41	0.53
1:B:290:PHE:CZ	1:B:376:THR:HG21	2.44	0.52
1:A:383:LEU:HD23	1:A:399:PHE:CD2	2.45	0.52
1:A:171:MET:SD	1:A:213:LEU:HD22	2.49	0.52
1:A:140:ALA:O	1:A:144:LEU:HD12	2.10	0.51
1:A:217:THR:CG2	1:A:274:LEU:HD22	2.39	0.51
1:B:322:PHE:CD1	1:B:369:ILE:HD12	2.47	0.50
1:A:333:ILE:HG23	1:A:343:PHE:CE1	2.48	0.49
1:B:288:ARG:CD	1:B:376:THR:HG22	2.42	0.48
1:B:389:PHE:CE1	1:B:396:VAL:HG13	2.49	0.48
1:A:346:ALA:HB2	1:A:369:ILE:HD13	1.94	0.48
1:B:394:LYS:HA	1:B:436:ILE:HD13	1.96	0.47
1:A:150:VAL:CG1	1:B:227:LEU:HD23	2.44	0.47
1:A:162:LEU:HD11	1:B:220:TYR:CD1	2.50	0.47
1:A:144:LEU:HD11	1:A:247:GLU:CG	2.46	0.46
1:A:156:GLY:HA2	1:A:159:LEU:HD12	1.97	0.46
1:B:371:GLU:O	1:B:375:GLU:HG3	2.17	0.45
1:B:391:THR:O	1:B:394:LYS:NZ	2.48	0.45
1:B:213:LEU:O	1:B:217:THR:HG23	2.17	0.44
1:B:444:LYS:HG3	1:B:446:VAL:HG23	1.99	0.44
1:B:103:LEU:HD21	1:B:350:PHE:CE1	2.54	0.43
1:A:322:PHE:CD1	1:A:369:ILE:HD12	2.54	0.42
1:A:236:LEU:HD23	1:B:236:LEU:HD21	2.00	0.42
1:B:250:ASN:OD1	1:B:250:ASN:N	2.53	0.42
1:B:437:GLN:O	1:B:456:LEU:HD22	2.20	0.42
1:A:168:ILE:CD1	4:B:503:XP4:H32	2.50	0.41
1:B:290:PHE:HZ	1:B:376:THR:HG21	1.83	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:THR:HG21	1:B:237:THR:HG22	2.01	0.41
1:B:346:ALA:HB2	1:B:369:ILE:HD13	2.01	0.41
1:A:236:LEU:HD23	1:B:236:LEU:HD11	2.02	0.41
1:B:395:LEU:HD23	1:B:453:THR:HA	2.02	0.41
1:A:284:PRO:O	1:A:288:ARG:NE	2.53	0.41
1:B:238:SER:N	1:B:239:PRO:HD2	2.36	0.41
1:A:108:ILE:CG2	1:A:123:ARG:HB2	2.50	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/407 (92%)	363 (97%)	11 (3%)	0	100	100
1	B	379/407 (93%)	371 (98%)	8 (2%)	0	100	100
All	All	753/814 (92%)	734 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/349 (93%)	322 (99%)	4 (1%)	71	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	329/349 (94%)	314 (95%)	15 (5%)	27	23
All	All	655/698 (94%)	636 (97%)	19 (3%)	42	43

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	218	ARG
1	A	261	MET
1	A	393	LYS
1	A	445	PHE
1	B	59	LYS
1	B	128	LYS
1	B	218	ARG
1	B	226	SER
1	B	249	MET
1	B	250	ASN
1	B	287	PHE
1	B	289	SER
1	B	325	SER
1	B	326	SER
1	B	359	LYS
1	B	368	SER
1	B	379	HIS
1	B	417	SER
1	B	434	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	501	3	4,4,4	0.93	0	6,6,6	0.46	0
4	XP4	B	503	-	39,39,39	1.52	6 (15%)	43,44,44	1.18	2 (4%)
2	PO4	A	503	-	4,4,4	0.93	0	6,6,6	0.46	0
2	PO4	B	501	3	4,4,4	0.90	0	6,6,6	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	XP4	B	503	-	-	20/41/41/41	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	503	XP4	O5-C4	4.08	1.45	1.33
4	B	503	XP4	O7-C18	3.52	1.44	1.34
4	B	503	XP4	P1-O3	2.90	1.66	1.54
4	B	503	XP4	O7-C2	-2.82	1.39	1.46
4	B	503	XP4	P1-O4	2.67	1.68	1.60
4	B	503	XP4	C19-C18	2.43	1.57	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	XP4	O7-C18-C19	4.50	121.19	111.50
4	B	503	XP4	O5-C4-C5	2.68	120.31	111.91

There are no chirality outliers.

All (20) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	503	XP4	C1-O4-P1-O1
4	B	503	XP4	C1-O4-P1-O2
4	B	503	XP4	C1-O4-P1-O3
4	B	503	XP4	O4-C1-C2-O7
4	B	503	XP4	O8-C18-O7-C2
4	B	503	XP4	C19-C18-O7-C2
4	B	503	XP4	C5-C4-O5-C3
4	B	503	XP4	C9-C10-C11-C12
4	B	503	XP4	C6-C7-C8-C9
4	B	503	XP4	O6-C4-O5-C3
4	B	503	XP4	O4-C1-C2-C3
4	B	503	XP4	C25-C26-C27-C28
4	B	503	XP4	C1-C2-C3-O5
4	B	503	XP4	O7-C2-C3-O5
4	B	503	XP4	C19-C20-C21-C22
4	B	503	XP4	C4-C5-C6-C7
4	B	503	XP4	C11-C10-C9-C8
4	B	503	XP4	C7-C8-C9-C10
4	B	503	XP4	C13-C14-C15-C16
4	B	503	XP4	C3-C2-O7-C18

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	503	XP4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	381/407 (93%)	0.13	11 (2%) 51 50	30, 48, 93, 116	0
1	B	384/407 (94%)	0.37	28 (7%) 15 14	31, 57, 110, 137	0
All	All	765/814 (93%)	0.25	39 (5%) 28 27	30, 52, 106, 137	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	249	MET	6.9
1	A	333	ILE	5.1
1	B	285	GLU	5.0
1	B	282	ARG	4.9
1	B	281	TRP	4.5
1	A	376	THR	4.3
1	B	236	LEU	4.2
1	A	383	LEU	3.7
1	B	270	LEU	3.6
1	B	287	PHE	3.5
1	B	143	ARG	3.5
1	A	377	VAL	3.5
1	B	379	HIS	3.5
1	A	332	GLN	3.5
1	B	273	PHE	3.3
1	B	333	ILE	3.2
1	A	273	PHE	3.0
1	A	143	ARG	2.9
1	B	350	PHE	2.9
1	B	456	LEU	2.7
1	A	456	LEU	2.7
1	B	266	MET	2.6
1	B	233	LEU	2.6
1	B	331	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	284	PRO	2.6
1	B	276	THR	2.5
1	A	224	LEU	2.5
1	B	241	ILE	2.5
1	B	267	TRP	2.4
1	B	274	LEU	2.3
1	B	262	THR	2.2
1	A	245	VAL	2.2
1	B	440	VAL	2.1
1	A	352	ILE	2.1
1	B	430	ILE	2.1
1	B	326	SER	2.1
1	B	264	PRO	2.0
1	B	334	PRO	2.0
1	B	245	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	XP4	B	503	40/40	0.67	0.35	52,91,120,127	0
3	MG	A	502	1/1	0.94	0.15	47,47,47,47	0
3	MG	B	502	1/1	0.96	0.07	49,49,49,49	0
2	PO4	A	503	5/5	0.97	0.18	65,69,86,106	0
2	PO4	B	501	5/5	0.99	0.15	31,34,36,39	0
2	PO4	A	501	5/5	0.99	0.17	34,36,37,40	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.