



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 03:27 pm BST

PDB ID : 4EOR
Title : Thr 160 phosphorylated CDK2 WT - human cyclin A3 complex with the inhibitor NU6102
Authors : Echalier, A.; Cot, E.; Camasses, A.; Hodimont, E.; Hoh, F.; Sheinerman, F.; Krasinska, L.; Fisher, D.
Deposited on : 2012-04-14
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	 4% (poor fit), 90% (0-1 outliers), 8% (2 outliers), 0% (3+ outliers)
1	C	298	 8% (poor fit), 75% (0-1 outliers), 14% (2 outliers), 9% (3+ outliers)
2	B	258	 3% (poor fit), 88% (0-1 outliers), 11% (2 outliers)
2	D	258	 10% (poor fit), 90% (0-1 outliers), 9% (2 outliers)

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	294	2401	1560	412	420	1	8	0	3	0
1	C	271	2171	1406	370	387	1	7	0	1	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	EXPRESSION TAG	UNP P24941
C	0	SER	-	EXPRESSION TAG	UNP P24941

- Molecule 2 is a protein called Cyclin-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	257	2083	1349	339	384	11	0	1	0
2	D	255	2059	1334	335	379	11	0	0	0

- Molecule 3 is O6-CYCLOHEXYLMETHOXY-2-(4'-SULPHAMOYLANILINO) PURINE (three-letter code: 4SP) (formula: $C_{18}H_{22}N_6O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			28	18	6	3	1		
3	C	1	Total	C	N	O	S	0	0
			28	18	6	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total	O	0	0
			52	52		
4	B	46	Total	O	0	0
			46	46		
4	C	19	Total	O	0	0
			19	19		
4	D	12	Total	O	0	0
			12	12		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclin-dependent kinase 2




- Molecule 1: Cyclin-dependent kinase 2



- Molecule 2: Cyclin-A2



- Molecule 2: Cyclin-A2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.97Å 135.44Å 149.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20 19.93 – 2.20	Depositor EDS
% Data completeness (in resolution range)	89.1 (20.00-2.20) 89.1 (19.93-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.79 (at 2.19Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.220 , 0.245 0.218 , 0.244	Depositor DCC
R_{free} test set	3443 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtrriage
Anisotropy	0.126	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8899	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, 4SP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/2452	0.52	0/3323
1	C	0.39	0/2210	0.54	1/2994 (0.0%)
2	B	0.37	0/2133	0.52	0/2896
2	D	0.34	0/2109	0.50	0/2864
All	All	0.37	0/8904	0.52	1/12077 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	220	GLY	N-CA-C	6.15	128.48	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2401	0	2444	12	0
1	C	2171	0	2207	29	0
2	B	2083	0	2105	14	0
2	D	2059	0	2079	9	0
3	A	28	0	22	2	0
3	C	28	0	22	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	52	0	0	0	0
4	B	46	0	0	0	0
4	C	19	0	0	1	0
4	D	12	0	0	0	0
All	All	8899	0	8879	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:218:THR:HB	1:C:219:LEU:CB	2.14	0.77
1:C:218:THR:CA	1:C:219:LEU:CB	2.70	0.68
1:C:219:LEU:H	1:C:269:TYR:HE2	1.42	0.68
1:C:60:HIS:HD2	1:C:62:ASN:H	1.40	0.67
1:A:60:HIS:HD2	1:A:62:ASN:H	1.40	0.67
1:A:60:HIS:CD2	1:A:62:ASN:H	2.15	0.65
1:C:60:HIS:CD2	1:C:62:ASN:H	2.15	0.64
1:C:218:THR:CB	1:C:219:LEU:CB	2.79	0.60
1:C:218:THR:HG21	1:C:249:SER:O	2.01	0.59
1:C:218:THR:HA	1:C:219:LEU:CB	2.34	0.57
2:D:374:GLU:HA	2:D:377:ILE:HD12	1.87	0.57
1:A:64:VAL:HG21	3:A:301:4SP:C8	2.38	0.53
2:B:361:HIS:CD2	2:B:391:LEU:HD21	2.43	0.53
1:A:177:CYS:HB2	1:A:233:MET:CE	2.40	0.52
1:C:64:VAL:HG21	3:C:301:4SP:C8	2.40	0.52
2:D:176:PRO:HA	2:D:179:HIS:CE1	2.46	0.51
2:B:214:LEU:HD22	2:B:253:LEU:HG	1.93	0.51
1:C:64:VAL:HG21	3:C:301:4SP:H8	1.93	0.50
1:C:253:PRO:HB2	1:C:254:PRO:HD3	1.94	0.49
2:D:346:PRO:O	2:D:349:LYS:HG2	2.11	0.49
1:A:49:ILE:HG23	2:B:306:LEU:HD12	1.93	0.49
1:C:155:PRO:HD2	2:D:316:THR:HB	1.96	0.48
1:C:216:PHE:O	1:C:219:LEU:HA	2.14	0.48
2:B:395:HIS:HE1	2:B:427:PRO:O	1.97	0.47
1:A:71:HIS:CE1	2:B:304:PHE:HZ	2.33	0.47
1:C:107:TYR:CZ	1:C:135:ILE:HD12	2.50	0.47
1:A:64:VAL:HG21	3:A:301:4SP:H8	1.95	0.47
2:D:344:ALA:HB1	2:D:348:LEU:HD22	1.97	0.47
2:B:219:VAL:HG22	2:B:232:LEU:HD11	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:268:HIS:HD2	1:C:270:ASP:H	1.62	0.46
1:A:88:LYS:HB2	1:A:130:PRO:HB2	1.97	0.46
1:C:252:VAL:H	1:C:260:ARG:HD3	1.81	0.46
1:A:71:HIS:CD2	1:A:76:LEU:HD13	2.52	0.45
2:B:345:ASP:HA	2:B:346:PRO:HA	1.71	0.45
1:C:87:LEU:O	1:C:91:MET:HG2	2.16	0.45
1:C:103:LEU:HD23	1:C:292:PRO:HG2	1.99	0.44
1:C:72:THR:HG22	1:C:73:GLU:H	1.82	0.44
2:D:372:TRP:HA	2:D:373:PRO:HD3	1.81	0.44
1:C:51:GLU:O	1:C:55:LEU:HB2	2.18	0.43
2:B:194:LYS:HA	2:B:195:PRO:HD3	1.93	0.42
2:D:345:ASP:HA	2:D:346:PRO:HA	1.82	0.42
2:B:281:ILE:H	2:B:281:ILE:HG13	1.75	0.42
1:C:88:LYS:HB2	1:C:130:PRO:HB2	2.02	0.42
2:D:347:TYR:OH	2:D:394:LEU:HA	2.20	0.42
2:B:346:PRO:O	2:B:349:LYS:HG2	2.19	0.42
1:C:133:LEU:HD11	1:C:192:ILE:HD13	2.02	0.42
1:C:34:LYS:HD3	1:C:75:LYS:HE3	2.01	0.42
2:B:388:LYS:HB3	2:B:389:PRO:HD3	2.02	0.42
1:A:99:ILE:HA	1:A:100:PRO:HD2	1.92	0.42
1:A:115:LEU:HD21	1:A:185:ASP:HB3	2.00	0.42
1:A:51:GLU:HG3	1:A:55:LEU:HD22	2.02	0.41
1:C:115:LEU:HD21	1:C:185:ASP:HB3	2.01	0.41
1:C:135:ILE:HD11	1:C:141:ILE:HD12	2.02	0.41
1:C:60:HIS:HB3	1:C:63:ILE:HD12	2.02	0.41
1:C:135:ILE:HG23	4:C:402:HOH:O	2.19	0.41
1:C:44:VAL:HA	1:C:45:PRO:HD3	1.92	0.41
2:B:203:GLN:HA	2:B:204:PRO:HD3	1.95	0.41
2:B:407:GLN:O	2:B:411:GLU:HG2	2.21	0.41
2:D:323:GLN:HA	2:D:324:PRO:HA	1.85	0.41
2:B:230:GLU:OE2	2:B:313:GLN:NE2	2.54	0.40
1:C:202:LEU:HD12	1:C:203:PHE:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/298 (98%)	284 (97%)	7 (2%)	1 (0%)	41	46
1	C	267/298 (90%)	253 (95%)	12 (4%)	2 (1%)	22	22
2	B	256/258 (99%)	255 (100%)	1 (0%)	0	100	100
2	D	253/258 (98%)	249 (98%)	4 (2%)	0	100	100
All	All	1068/1112 (96%)	1041 (98%)	24 (2%)	3 (0%)	41	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	164	VAL
1	A	164	VAL
1	C	216	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/262 (100%)	249 (95%)	12 (5%)	27	34
1	C	235/262 (90%)	224 (95%)	11 (5%)	26	33
2	B	232/232 (100%)	226 (97%)	6 (3%)	46	58
2	D	229/232 (99%)	220 (96%)	9 (4%)	32	41
All	All	957/988 (97%)	919 (96%)	38 (4%)	32	40

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	15[A]	TYR
1	A	15[B]	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	55	LEU
1	A	59	ASN
1	A	71	HIS
1	A	101	LEU
1	A	122	ARG
1	A	148	LEU
1	A	150	ARG
1	A	206	ASP
1	A	297	ARG
2	B	196	LYS
2	B	232	LEU
2	B	292	LEU
2	B	296	HIS
2	B	378	ARG
2	B	403	GLN
1	C	55	LEU
1	C	103	LEU
1	C	122	ARG
1	C	131	GLN
1	C	148	LEU
1	C	150	ARG
1	C	202	LEU
1	C	212	LEU
1	C	217	ARG
1	C	218	THR
1	C	278	LYS
2	D	177	ASP
2	D	196	LYS
2	D	197	VAL
2	D	232	LEU
2	D	245	SER
2	D	292	LEU
2	D	328	LYS
2	D	348	LEU
2	D	429	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	60	HIS
1	A	71	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	246	GLN
1	A	265	GLN
1	A	268	HIS
2	B	183	HIS
2	B	254	GLN
2	B	317	GLN
2	B	395	HIS
2	B	403	GLN
1	C	60	HIS
1	C	71	HIS
1	C	113	GLN
1	C	131	GLN
1	C	268	HIS
2	D	254	GLN
2	D	317	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	C	160	1	8,10,11	0.87	0	10,14,16	1.01	1 (10%)
1	TPO	A	160	1	8,10,11	0.82	0	10,14,16	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	C	160	1	-	0/9/11/13	-
1	TPO	A	160	1	-	0/9/11/13	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	160	TPO	O3P-P-O2P	2.13	115.76	107.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	4SP	A	301	-	28,31,31	3.39	10 (35%)	35,44,44	2.30	12 (34%)
3	4SP	C	301	-	28,31,31	2.38	7 (25%)	35,44,44	1.91	8 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4SP	A	301	-	-	4/15/23/23	0/4/4/4
3	4SP	C	301	-	-	3/15/23/23	0/4/4/4

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	4SP	C20-S23	-12.65	1.57	1.77
3	C	301	4SP	C20-S23	-10.06	1.61	1.77
3	A	301	4SP	S23-N26	-8.87	1.42	1.60
3	C	301	4SP	S23-N26	-4.05	1.52	1.60
3	A	301	4SP	C17-N2	-3.60	1.32	1.40
3	A	301	4SP	O24-S23	-3.31	1.37	1.43
3	A	301	4SP	O25-S23	-3.17	1.37	1.43
3	C	301	4SP	C6-N1	2.77	1.36	1.31
3	A	301	4SP	C4-N3	-2.71	1.31	1.36
3	A	301	4SP	C5-N7	-2.68	1.30	1.39
3	A	301	4SP	C4-N9	-2.66	1.30	1.34
3	C	301	4SP	O6-C6	2.59	1.37	1.35
3	A	301	4SP	O6-C6	-2.40	1.32	1.35
3	C	301	4SP	C5-C4	2.34	1.47	1.40
3	C	301	4SP	O25-S23	2.31	1.48	1.43
3	C	301	4SP	O24-S23	2.26	1.47	1.43
3	A	301	4SP	C2-N2	-2.07	1.32	1.36

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	4SP	O25-S23-C20	6.26	114.34	107.35
3	C	301	4SP	O25-S23-O24	-6.04	108.84	118.76
3	A	301	4SP	O24-S23-C20	5.62	113.62	107.35
3	A	301	4SP	O25-S23-O24	-5.00	110.53	118.76
3	C	301	4SP	C2-N1-C6	4.76	123.58	115.18
3	A	301	4SP	C2-N1-C6	4.30	122.77	115.18
3	A	301	4SP	C13-C12-C11	-3.61	105.33	112.15
3	C	301	4SP	N3-C2-N1	-3.55	120.62	126.23
3	A	301	4SP	C4-C5-N7	-3.09	106.17	109.40
3	C	301	4SP	C4-C5-N7	-3.09	106.18	109.40
3	C	301	4SP	O25-S23-C20	2.68	110.35	107.35
3	C	301	4SP	C2-N3-C4	2.59	118.22	115.28
3	A	301	4SP	C5-C6-N1	-2.41	118.66	123.26
3	A	301	4SP	N3-C2-N1	-2.41	122.41	126.23
3	A	301	4SP	C18-C19-C20	2.41	121.94	119.45
3	C	301	4SP	C10-O6-C6	2.40	119.79	117.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	301	4SP	C5-C6-N1	-2.38	118.72	123.26
3	A	301	4SP	O24-S23-N26	-2.21	104.08	107.36
3	A	301	4SP	C15-C16-C11	-2.05	108.28	112.15
3	A	301	4SP	C21-C20-C19	-2.04	117.60	120.44

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	301	4SP	C19-C20-S23-O24
3	A	301	4SP	C19-C20-S23-N26
3	A	301	4SP	C21-C20-S23-O24
3	A	301	4SP	C21-C20-S23-N26
3	C	301	4SP	N1-C2-N2-C17
3	C	301	4SP	N3-C2-N2-C17
3	C	301	4SP	C19-C20-S23-O24

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	4SP	2	0
3	C	301	4SP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/298 (98%)	0.02	12 (4%) 37 35	10, 22, 39, 51	2 (0%)
1	C	270/298 (90%)	0.42	24 (8%) 9 8	25, 34, 59, 81	5 (1%)
2	B	257/258 (99%)	0.02	9 (3%) 44 42	13, 25, 38, 49	3 (1%)
2	D	255/258 (98%)	0.41	27 (10%) 6 5	20, 37, 56, 67	0
All	All	1075/1112 (96%)	0.21	72 (6%) 17 16	10, 29, 51, 81	10 (0%)

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	248	PHE	11.6
1	C	249	SER	7.0
1	C	222	PRO	6.5
1	C	247	ASP	6.5
2	B	283	ASP	6.2
1	C	253	PRO	5.4
1	C	220	GLY	5.0
1	A	15[A]	TYR	4.6
1	A	0	SER	4.6
1	A	96	LEU	4.6
1	C	287	GLN	4.6
1	C	288	ASP	4.4
1	C	221	THR	4.4
2	D	325	ALA	4.3
1	C	219	LEU	4.2
2	D	176	PRO	4.2
2	D	284	ASP	4.2
1	C	73	GLU	4.1
2	D	428	GLU	4.0
2	B	284	ASP	3.9
2	B	431	ASN	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	293	VAL	3.8
2	D	283	ASP	3.7
2	D	175	VAL	3.6
2	D	323	GLN	3.6
2	D	332	LEU	3.6
1	A	95	ALA	3.6
1	A	162	GLU	3.5
2	B	323	GLN	3.5
2	D	371	SER	3.5
2	D	368	THR	3.4
2	D	197	VAL	3.4
1	C	218	THR	3.4
2	D	383	THR	3.3
2	B	432	LEU	3.2
2	D	425	ASN	2.9
1	C	97	THR	2.9
2	D	416	SER	2.9
1	C	162	GLU	2.9
2	D	378	ARG	2.9
1	C	269	TYR	2.9
2	D	177	ASP	2.8
1	A	36	ARG	2.8
1	A	73	GLU	2.8
2	B	201	LYS	2.8
2	B	176	PRO	2.7
1	C	251	VAL	2.7
2	B	177	ASP	2.7
1	A	74	ASN	2.6
1	A	32	LEU	2.6
1	C	252	VAL	2.6
1	C	36	ARG	2.5
2	D	280	TYR	2.5
2	D	419	HIS	2.5
1	A	71	HIS	2.4
2	D	392	MET	2.4
1	C	138	GLU	2.3
2	D	411	GLU	2.3
2	D	327	CYS	2.3
1	C	178	LYS	2.2
1	A	206	ASP	2.2
1	C	250	LYS	2.2
1	C	59	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	427	PRO	2.2
1	A	247	ASP	2.2
2	D	429	THR	2.2
2	D	336	LEU	2.1
2	D	367	VAL	2.1
2	D	324	PRO	2.1
2	B	224	GLU	2.0
1	C	256	ASP	2.0
2	D	370	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	C	160	11/12	0.96	0.09	27,30,32,33	0
1	TPO	A	160	11/12	0.98	0.09	15,18,19,20	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	4SP	C	301	28/28	0.88	0.17	36,38,40,40	0
3	4SP	A	301	28/28	0.92	0.12	26,28,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 4SP C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.