

Full wwPDB X-ray Structure Validation Report (i)

Mar 30, 2025 - 03:59 am BST

PDB ID	:	$9\mathrm{ESY} \ / \ \mathrm{pdb}_00009\mathrm{esy}$
Title	:	CDK2-cyclin A in complex with FragLite 16
Authors	:	Hope, I.; Martin, M.P.; Waring, M.J.; Noble, M.E.M.; Endicott, J.A.; Tatum,
		N.J.
Deposited on	:	2024-03-26
Resolution	:	2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	?.? (???), CSD ??CSD?? (????)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
wietric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158(2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	302	<u>6%</u> 86%	12% ••
1	С	302	3% 80%	8% • 12%
2	В	268	% 90%	8% •
2	D	268	83%	14% •



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 9169 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace	
1	А	299	Total	С	N	0	Р	S	0	2	0
			2417	1567	411	430	1	8	_		
1	C	266	Total	\mathbf{C}	Ν	0	Р	\mathbf{S}	0	0	0
1		200	2137	1386	364	379	1	7	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-3	GLY	-	expression tag	UNP P24941
А	-2	PRO	-	expression tag	UNP P24941
А	-1	GLY	-	expression tag	UNP P24941
А	0	SER	-	expression tag	UNP P24941
С	-3	GLY	-	expression tag	UNP P24941
С	-2	PRO	-	expression tag	UNP P24941
С	-1	GLY	-	expression tag	UNP P24941
C	0	SER	-	expression tag	UNP P24941

• Molecule 2 is a protein called Cyclin-A2.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
0	Р	າຄາ	Total	С	Ν	0	\mathbf{S}	0	0	0
	D	202	2110	1366	344	390	10	0	0	0
0	П	262	Total	С	Ν	0	S	0	0	0
	D	202	2110	1366	344	390	10	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	171	GLY	-	expression tag	UNP P30274
В	433	HIS	-	expression tag	UNP P30274
В	434	HIS	-	expression tag	UNP P30274
В	435	HIS	-	expression tag	UNP P30274



Chain	Residue	Modelled	Actual	Comment	Reference
В	436	HIS	-	expression tag	UNP P30274
В	437	HIS	-	expression tag	UNP P30274
В	438	HIS	-	expression tag	UNP P30274
D	171	GLY	-	expression tag	UNP P30274
D	433	HIS	-	expression tag	UNP P30274
D	434	HIS	-	expression tag	UNP P30274
D	435	HIS	-	expression tag	UNP P30274
D	436	HIS	-	expression tag	UNP P30274
D	437	HIS	-	expression tag	UNP P30274
D	438	HIS	-	expression tag	UNP P30274

• Molecule 3 is 4-bromanyl-1,8-naphthyridine (CCD ID: HH8) (formula: $C_8H_5BrN_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	Λ	1	Total	Br	С	Ν	0	0	
5	Л	I	11	1	8	2	0	0	
2	Δ	Δ	1	Total	Br	С	Ν	0	0
5	A	1	11	1	8	2	0	0	
9	D	1	Total	Br	С	Ν	0	0	
0	3 B		11	1	8	2	0	0	
3	C	1	Total	Br	С	Ν	0	0	
			11	1	8	2	0		

• Molecule 4 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	124	Total O 124 124	0	0
4	В	107	Total O 107 107	0	0
4	С	72	Total O 72 72	0	0
4	D	48	Total O 48 48	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Cyclin-dependent kinase 2





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	74.07Å 133.63Å 148.06Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	74.14 - 2.39	Depositor
	74.14 - 2.39	EDS
% Data completeness	99.3 (74.14-2.39)	Depositor
(in resolution range)	99.3(74.14-2.39)	EDS
R_{merge}	0.19	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.93 (at 2.40 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
B B.	0.256 , 0.262	Depositor
II, II, <i>free</i>	0.238 , 0.242	DCC
R_{free} test set	2995 reflections $(5.11%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	57.7	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.36 , 32.8	EDS
L-test for $twinning^2$	$ < L >=0.51, < L^2>=0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9169	wwPDB-VP
Average B, all atoms $(Å^2)$	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HH8, TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Chain		Bond lengths		Bond angles	
	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.52	0/2469	0.88	0/3350
1	С	0.46	0/2174	0.85	0/2944
2	В	0.52	0/2160	0.88	1/2937~(0.0%)
2	D	0.49	0/2160	0.84	1/2937~(0.0%)
All	All	0.50	0/8963	0.86	2/12168~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	3
1	С	0	1
2	В	0	1
2	D	0	1
All	All	0	6

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	346	PRO	N-CA-CB	-11.52	89.48	103.30
2	D	346	PRO	N-CA-CB	-10.22	91.04	103.30

There are no chirality outliers.

All (6) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	А	122	ARG	Sidechain
1	А	150	ARG	Sidechain
1	А	36	ARG	Sidechain
2	В	293	ARG	Sidechain
1	С	150	ARG	Sidechain
2	D	293	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2417	0	2451	36	0
1	С	2137	0	2193	22	0
2	В	2110	0	2128	20	0
2	D	2110	0	2128	27	0
3	А	22	0	0	0	0
3	В	11	0	0	3	0
3	С	11	0	0	0	0
4	А	124	0	0	5	0
4	В	107	0	0	5	0
4	С	72	0	0	2	0
4	D	48	0	0	1	0
All	All	9169	0	8900	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:175:VAL:CG2	2:D:178:TYR:HB2	1.96	0.96
1:C:71:HIS:HD2	2:D:296:HIS:NE2	1.64	0.95
1:A:38:ASP:OD1	1:A:39:THR:HG23	1.74	0.88
1:C:59:ASN:HD21	1:C:65:LYS:HZ2	1.15	0.88
2:D:175:VAL:HG22	2:D:178:TYR:HB2	1.58	0.83
1:A:36:ARG:O	1:A:40:GLU:CG	2.29	0.81
1:A:36:ARG:O	1:A:40:GLU:HG2	1.83	0.79



		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:C:59:ASN:HD21	1:C:65:LYS:NZ	1.82	0.78
2:B:305:ASP:HB2	4:B:604:HOH:O	1.87	0.75
1:C:59:ASN:ND2	1:C:65:LYS:NZ	2.36	0.74
1:A:97:THR:HG23	1:A:98:GLY:O	1.89	0.73
1:A:177:CYS:SG	1:A:233:MET:HG2	2.29	0.71
1:A:59[A]:ASN:ND2	1:A:65:LYS:NZ	2.38	0.71
4:C:711:HOH:O	2:D:305:ASP:HB2	1.91	0.70
4:A:757:HOH:O	2:B:189:MET:HE2	1.94	0.67
1:C:83:LEU:CD1	1:C:134:LEU:HB2	2.25	0.66
2:D:233:HIS:HD2	4:D:535:HOH:O	1.77	0.66
2:B:241:ARG:HH22	2:B:305:ASP:HB3	1.60	0.65
2:B:395:HIS:HE1	2:B:427:PRO:O	1.79	0.65
1:A:71:HIS:CD2	1:A:76:LEU:HD13	2.32	0.64
2:B:305:ASP:CB	4:B:604:HOH:O	2.43	0.64
2:B:225:TYR:HE2	2:B:281:ILE:HG21	1.63	0.64
1:A:38:ASP:HB3	1:A:74:ASN:HB3	1.81	0.62
1:C:83:LEU:HD12	1:C:134:LEU:HB2	1.81	0.62
1:A:59[A]:ASN:ND2	1:A:65:LYS:HZ2	1.98	0.60
1:A:97:THR:HG23	1:A:98:GLY:N	2.15	0.60
1:C:71:HIS:CD2	2:D:296:HIS:NE2	2.57	0.60
1:C:83:LEU:HD12	1:C:134:LEU:CB	2.32	0.59
2:B:221:VAL:HG21	2:B:281:ILE:HD12	1.84	0.59
1:A:36:ARG:O	1:A:40:GLU:HG3	2.03	0.58
1:A:97:THR:HG21	1:A:295:HIS:HE1	1.68	0.57
1:A:38:ASP:OD1	1:A:39:THR:N	2.39	0.56
2:D:430:LEU:O	2:D:431:ASN:C	2.44	0.56
1:A:97:THR:CG2	1:A:98:GLY:N	2.68	0.55
1:A:59[A]:ASN:ND2	1:A:65:LYS:HZ3	2.03	0.55
1:A:283:HIS:ND1	1:A:284:PRO:HD2	2.22	0.55
2:D:322:GLN:HG3	2:D:325:ALA:HA	1.90	0.53
1:A:38:ASP:OD1	1:A:39:THR:CG2	2.53	0.53
1:C:60:HIS:CG	1:C:61:PRO:HD2	2.44	0.52
1:A:56:LYS:HB2	4:A:810:HOH:O	2.10	0.51
1:C:83:LEU:CD1	1:C:134:LEU:CB	2.87	0.51
1:C:88:LYS:HA	1:C:91:MET:HE2	1.93	0.51
1:C:59:ASN:ND2	1:C:65:LYS:HZ3	2.08	0.51
1:A:163:VAL:HG23	1:A:164:VAL:HG23	1.93	0.50
1:A:107:TYR:O	1:A:111:LEU:HG	2.12	0.50
2:B:221:VAL:CG2	2:B:281:ILE:HD12	2.42	0.49
1:C:88:LYS:HB2	1:C:130:PRO:HB2	1.95	0.49
3:B:501:HH8:C5	4:B:608:HOH:O	2.62	0.48



	lo de page	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:C:255:LEU:O	1:C:260:ARG:NH1	2.47	0.48	
2:D:175:VAL:O	2:D:175:VAL:HG13	2.13	0.48	
4:C:711:HOH:O	2:D:305:ASP:CB	2.58	0.47	
1:C:15:TYR:OH	1:C:51:GLU:OE1	2.21	0.47	
2:D:241:ARG:HH22	2:D:305:ASP:HB3	1.79	0.47	
1:A:230:VAL:HA	1:A:233:MET:SD	2.55	0.46	
2:D:216:ASP:HB2	2:D:406:GLN:HG2	1.96	0.46	
4:A:757:HOH:O	2:B:189:MET:CE	2.60	0.46	
2:D:421:VAL:HA	2:D:424:LEU:HG	1.97	0.46	
1:C:219:LEU:HB2	1:C:269:TYR:OH	2.15	0.46	
2:D:323:GLN:HA	2:D:324:PRO:HA	1.70	0.46	
1:A:60:HIS:CG	1:A:61:PRO:HD2	2.51	0.46	
1:C:121:HIS:C	1:C:122:ARG:HG3	2.36	0.45	
2:D:175:VAL:HG21	2:D:178:TYR:HB2	1.91	0.45	
2:B:241:ARG:NH2	2:B:305:ASP:HB3	2.28	0.45	
2:B:321:HIS:HE1	3:B:501:HH8:C5	2.30	0.45	
2:D:322:GLN:CG	2:D:325:ALA:HA	2.47	0.45	
1:A:49:ILE:HG23	2:B:306:LEU:HD12	1.99	0.44	
1:A:44:VAL:HA	1:A:45:PRO:HD3	1.88	0.44	
1:A:177:CYS:HB2	4:A:714:HOH:O	2.17	0.44	
2:D:338:GLU:HB3	2:D:409:ILE:HD13	1.99	0.44	
1:C:83:LEU:HD13	1:C:134:LEU:HB2	1.96	0.44	
1:A:186:ILE:HD13	1:A:186:ILE:HA	1.87	0.44	
1:A:71:HIS:HD2	1:A:76:LEU:HD13	1.78	0.44	
2:D:346:PRO:O	2:D:349:LYS:HG2	2.17	0.44	
2:D:275:VAL:HG21	2:D:292:LEU:HD21	2.00	0.43	
1:A:37:LEU:HA	1:A:40:GLU:CG	2.48	0.43	
2:B:183:HIS:HE1	4:B:629:HOH:O	2.00	0.43	
1:C:83:LEU:HD12	1:C:134:LEU:CD1	2.48	0.43	
2:B:345:ASP:HA	2:B:346:PRO:HA	1.75	0.43	
2:B:401:ALA:N	2:B:402:PRO:CD	2.82	0.43	
2:D:194:LYS:HA	2:D:195:PRO:HD3	1.85	0.43	
1:C:12:GLU:HG3	1:C:16:GLY:O	2.19	0.42	
2:D:326:ASN:HB3	2:D:329:VAL:HB	2.02	0.42	
2:D:346:PRO:HD2	2:D:347:TYR:CD2	2.53	0.42	
2:B:321:HIS:CE1	3:B:501:HH8:C5	3.03	0.42	
2:B:396:GLN:HE21	2:B:400:ARG:CZ	2.33	0.42	
2:B:233:HIS:HD2	4:B:650:HOH:O	2.01	0.42	
2:D:404:HIS:O	2:D:407:GLN:NE2	2.45	0.42	
1:A:177:CYS:SG	1:A:233:MET:CG	3.06	0.42	
1:C:15:TYR:CG	1:C:35:ILE:HG12	2.54	0.42	



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:D:401:ALA:HB3	2:D:402:PRO:HD3	2.01	0.42
1:A:88:LYS:HA	1:A:91:MET:HE2	2.02	0.41
1:A:290:THR:HA	4:A:811:HOH:O	2.21	0.41
1:A:195:GLU:O	1:A:199:ARG:N	2.52	0.41
2:B:346:PRO:HD2	2:B:347:TYR:CD2	2.56	0.41
2:D:414:LYS:HG2	2:D:423:LEU:HG	2.03	0.41
1:A:88:LYS:HA	1:A:91:MET:CE	2.51	0.41
1:A:157:ARG:HE	1:A:157:ARG:HB2	1.67	0.40
1:A:283:HIS:CE1	1:A:284:PRO:HD2	2.56	0.40
2:D:184:THR:O	2:D:188:GLU:HG3	2.21	0.40
1:A:78:LEU:N	1:A:78:LEU:HD23	2.36	0.40
1:C:91:MET:HE2	1:C:130:PRO:HB3	2.03	0.40
2:D:394:LEU:HD12	2:D:394:LEU:HA	1.87	0.40
2:B:299:LEU:HD21	2:B:306:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	298/302~(99%)	289~(97%)	8 (3%)	1 (0%)	37	51
1	С	261/302~(86%)	253~(97%)	7 (3%)	1 (0%)	30	44
2	В	260/268~(97%)	254 (98%)	6(2%)	0	100	100
2	D	260/268~(97%)	252~(97%)	7 (3%)	1 (0%)	30	44
All	All	1079/1140~(95%)	1048 (97%)	28(3%)	3 (0%)	37	51

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	164	VAL
	a r.	1	1



 $Continued \ from \ previous \ page...$

Mol	Chain	Res	Type
1	А	164	VAL
2	D	420	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	264/264~(100%)	263~(100%)	1 (0%)	89 95
1	С	233/264~(88%)	229~(98%)	4 (2%)	56 75
2	В	234/240~(98%)	232~(99%)	2 (1%)	75 88
2	D	234/240~(98%)	231~(99%)	3 (1%)	65 81
All	All	965/1008~(96%)	955~(99%)	10 (1%)	73 86

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	122	ARG
2	В	199	TYR
2	В	346	PRO
1	С	47	THR
1	С	72	THR
1	С	122	ARG
1	С	150	ARG
2	D	205	ASP
2	D	293	ARG
2	D	346	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	295	HIS
2	В	183	HIS
2	В	233	HIS
2	В	254	GLN



	9	1	1 0
Mol	Chain	Res	Type
2	В	317	GLN
2	В	378	GLN
2	В	395	HIS
2	В	396	GLN
1	С	59	ASN
1	С	71	HIS
2	D	179	HIS
2	D	233	HIS
2	D	254	GLN

Continued from previous page...

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.There are no bond length outliers.There are no bond angle outliers.There are no chirality outliers.There are no torsion outliers.



There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and similar rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	А	298/302~(98%)	0.18	19 (6%) 27 24	12, 28, 67, 110	2~(0%)
1	С	265/302~(87%)	0.24	9 (3%) 48 45	17, 30, 54, 92	0
2	В	262/268~(97%)	-0.02	3 (1%) 77 75	16, 28, 49, 75	0
2	D	262/268~(97%)	0.70	27 (10%) 13 11	23, 32, 48, 75	0
All	All	1087/1140~(95%)	0.27	58 (5%) 33 30	12, 30, 54, 110	2~(0%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	251	VAL	7.5
2	D	432	VAL	6.0
1	А	15	TYR	5.6
1	А	162	GLU	5.0
1	А	96	LEU	4.8
1	А	-2	PRO	4.7
1	А	71	HIS	4.4
2	D	324	PRO	4.2
2	D	390	CYS	3.7
1	А	36	ARG	3.5
1	С	282	ALA	3.3
2	D	373	PRO	3.2
1	А	13	GLY	3.2
2	D	372	TRP	3.0
1	А	161	HIS	2.9
2	D	424	LEU	2.9
2	D	376	LEU	2.9
2	В	432	VAL	2.8
1	С	73	GLU	2.8
2	В	296	HIS	2.7
2	D	327	CYS	2.7



Mol	Chain	Res	Type	RSRZ
1	А	14	THR	2.6
2	D	320	LEU	2.6
2	D	384	LEU	2.6
1	А	0	SER	2.6
1	А	94	SER	2.6
1	А	39	THR	2.5
1	С	252	VAL	2.5
2	D	175	VAL	2.5
1	С	295	HIS	2.5
2	D	392	LEU	2.5
2	D	418	TYR	2.4
1	A	74	ASN	2.4
2	D	429	THR	2.4
2	D	363	ALA	2.4
2	D	365	TYR	2.4
2	D	319	PHE	2.3
1	А	72	THR	2.3
1	А	73	GLU	2.3
2	D	368	THR	2.3
2	D	381	GLY	2.3
2	D	326	ASN	2.3
2	D	375	SER	2.2
1	А	157	ARG	2.2
1	А	17	VAL	2.2
2	В	192	LYS	2.2
2	D	360	PHE	2.2
2	D	425	ASN	2.1
2	D	171	GLY	2.1
2	D	420	GLY	2.1
1	А	206	ASP	2.1
1	С	258	ASP	2.1
1	C	219	LEU	2.1
2	D	332	LEU	2.1
1	C	216	PHE	2.0
1	A	296	LEU	2.0
1	C	273	LYS	2.0
2	D	422	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
1	TPO	С	160	11/12	0.96	0.07	$26,\!29,\!37,\!40$	0
1	TPO	А	160	11/12	0.98	0.07	19,27,31,32	0

median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
3	HH8	В	501	11/11	0.74	0.23	80,92,128,133	0
3	HH8	С	601	11/11	0.85	0.12	35,41,60,122	0
3	HH8	А	602	11/11	0.88	0.21	47,65,79,87	0
3	HH8	А	601	11/11	0.94	0.13	21,28,35,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















6.5 Other polymers (i)

There are no such residues in this entry.

