



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 12:25 PM EDT

PDB ID : 1EUC
Title : CRYSTAL STRUCTURE OF DEPHOSPHORYLATED PIG HEART, GTP-SPECIFIC SUCCINYL-COA SYNTHETASE
Authors : Fraser, M.E.; James, M.N.G.; Bridger, W.A.; Wolodko, W.T.
Deposited on : 2000-04-14
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	
2	B	396	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUCCINYL-COA SYNTHETASE, ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	306	2258	1421	401	423	13	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	MET	-	cloning artifact	UNP O19069
A	-3	ALA	-	cloning artifact	UNP O19069

- Molecule 2 is a protein called SUCCINYL-COA SYNTHETASE, BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	394	2981	1883	503	580	15	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	cloning artifact	UNP P53590
B	1	VAL	LEU	engineered mutation	UNP P53590
B	332	CME	CYS	modified residue	UNP P53590

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
3	A	1	5	4	1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0
4	B	1	5	4	1	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total 1	Zn 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	124	Total 124	O 124	0	0
6	B	80	Total 80	O 80	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SUCCINYL-COA SYNTHETASE, ALPHA CHAIN



- Molecule 2: SUCCINYL-COA SYNTHETASE, BETA CHAIN



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.80Å 81.97Å 49.89Å 90.00° 104.53° 90.00°	Depositor
Resolution (Å)	59.00 – 2.10 58.67 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.4 (59.00-2.10) 95.4 (58.67-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.10Å)	Xtrriage
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.158 , 0.220 0.160 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtrriage
Anisotropy	0.115	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.030 for -h-1,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5459	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CME, SO4, ZN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.86	4/2305 (0.2%)	0.89	0/3115
2	B	0.77	1/3007 (0.0%)	0.90	3/4058 (0.1%)
All	All	0.81	5/5312 (0.1%)	0.90	3/7173 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	268	CYS	CB-SG	-6.15	1.71	1.82
1	A	183[A]	CYS	CB-SG	-5.94	1.72	1.81
1	A	183[B]	CYS	CB-SG	-5.94	1.72	1.81
1	A	102	CYS	CB-SG	-5.40	1.73	1.81
1	A	201	GLU	CG-CD	-5.02	1.44	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	299	GLY	N-CA-C	-6.80	96.11	113.10
2	B	208	PHE	N-CA-C	-6.12	94.47	111.00
2	B	352	LEU	CA-CB-CG	5.23	127.33	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2306	37	0
2	B	2981	0	3035	79	0
3	A	5	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	B	1	0	0	0	0
6	A	124	0	0	2	0
6	B	80	0	0	4	0
All	All	5459	0	5341	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:GLY:HA2	2:B:258:LYS:HD3	1.54	0.90
2:B:246:GLU:HB2	2:B:249:GLU:HG3	1.66	0.78
2:B:361:VAL:O	2:B:365:GLN:HG2	1.85	0.76
2:B:344:ARG:CB	2:B:344:ARG:HH11	2.00	0.75
2:B:139:ASP:OD2	2:B:141:GLU:HB2	1.87	0.74
2:B:269:PHE:HZ	2:B:309:PHE:CE2	2.07	0.73
1:A:136:ILE:HG22	1:A:183[B]:CYS:HB3	1.74	0.69
2:B:252:ALA:HB2	2:B:311:LEU:HD21	1.75	0.69
1:A:136:ILE:HG22	1:A:183[A]:CYS:HB2	1.75	0.68
1:A:65:LYS:O	1:A:69:GLU:HG2	1.93	0.68
2:B:275:LEU:HD21	2:B:324:ASN:ND2	2.09	0.67
2:B:41:LYS:HG3	2:B:42:GLU:N	2.09	0.67
1:A:256:ARG:HH21	2:B:0:MET:HE1	1.59	0.66
2:B:0:MET:HG2	6:B:467:HOH:O	1.97	0.65
1:A:191:PHE:CE1	2:B:123:ARG:HA	2.33	0.63
2:B:147:ASN:HA	2:B:149:GLU:OE2	1.99	0.62
1:A:221:ASN:H	1:A:221:ASN:HD22	1.46	0.62
1:A:245:PHE:HB2	1:A:292:LEU:HD21	1.82	0.62
1:A:299:GLU:O	1:A:303:ARG:HG2	2.00	0.61
2:B:269:PHE:CD1	2:B:296:LEU:HD11	2.36	0.60
2:B:258:LYS:HD2	2:B:295:ASP:HB3	1.83	0.60
2:B:369:THR:HG23	6:B:475:HOH:O	2.02	0.60
2:B:210:GLU:HA	2:B:215:GLN:O	2.03	0.59
2:B:41:LYS:HG3	2:B:42:GLU:H	1.67	0.58
2:B:269:PHE:HD1	2:B:296:LEU:HD11	1.68	0.58
2:B:352:LEU:HD12	2:B:375:ILE:HG12	1.86	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:300:VAL:O	2:B:300:VAL:HG13	2.04	0.57
2:B:246:GLU:O	2:B:249:GLU:HB2	2.05	0.57
2:B:85:GLY:HA2	2:B:99:LYS:HE3	1.86	0.56
2:B:340:THR:O	2:B:344:ARG:HD3	2.05	0.56
2:B:35:ALA:HB1	2:B:43:ILE:HD13	1.86	0.56
2:B:35:ALA:HB1	2:B:43:ILE:CD1	2.35	0.56
2:B:90:THR:C	2:B:92:GLN:H	2.09	0.56
2:B:230:GLU:HG3	2:B:237:PHE:CG	2.40	0.55
2:B:213:GLU:CD	2:B:213:GLU:H	2.10	0.55
1:A:116:LYS:NZ	6:A:391:HOH:O	2.26	0.55
2:B:383:ASP:OD1	2:B:387:LYS:HE3	2.07	0.54
2:B:344:ARG:HH11	2:B:344:ARG:CG	2.21	0.54
2:B:269:PHE:HB3	2:B:294:LEU:HD23	1.89	0.54
2:B:358:GLY:O	2:B:361:VAL:HG23	2.09	0.52
1:A:256:ARG:HG3	1:A:263:ILE:HD12	1.91	0.52
2:B:111:ILE:HG21	2:B:114:GLU:HG2	1.92	0.52
2:B:131:VAL:HG22	2:B:153:LYS:HG2	1.90	0.52
1:A:256:ARG:NH2	2:B:0:MET:HE1	2.25	0.52
2:B:135:GLN:O	2:B:135:GLN:HG3	2.10	0.52
2:B:139:ASP:HB3	2:B:142:GLU:OE2	2.11	0.51
2:B:122:ASP:OD2	2:B:125:CYS:SG	2.68	0.51
2:B:242:LYS:HA	2:B:245:ASN:ND2	2.26	0.51
1:A:166:THR:HG21	1:A:183[B]:CYS:SG	2.50	0.51
2:B:269:PHE:CZ	2:B:309:PHE:CE2	2.95	0.51
2:B:344:ARG:HH11	2:B:344:ARG:HB2	1.75	0.51
1:A:221:ASN:HD22	1:A:221:ASN:N	2.07	0.50
2:B:371:SER:HB2	2:B:373:LEU:HD12	1.94	0.50
1:A:136:ILE:CG2	1:A:183[A]:CYS:HB2	2.42	0.49
2:B:325:ILE:HD12	2:B:335:ILE:HG21	1.95	0.49
2:B:52:GLY:H	2:B:92:GLN:NE2	2.12	0.48
2:B:147:ASN:CA	2:B:149:GLU:OE2	2.61	0.48
2:B:130:LEU:O	2:B:153:LYS:HA	2.14	0.48
2:B:18:LYS:NZ	2:B:183:GLN:OE1	2.46	0.48
2:B:248:ILE:HG23	2:B:311:LEU:HG	1.96	0.47
1:A:275:ILE:HG12	1:A:285:VAL:HG11	1.97	0.47
2:B:50:LEU:HD11	2:B:241:ASP:HB3	1.96	0.47
2:B:206:ASN:HA	2:B:207:PRO:HA	1.67	0.47
1:A:53:LYS:HD3	6:A:405:HOH:O	2.15	0.46
2:B:29:ASN:O	2:B:32:LEU:HB3	2.16	0.46
1:A:10:LEU:CD2	1:A:128:ILE:HD11	2.45	0.46
1:A:294:THR:O	1:A:298:LYS:HG2	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:321:ILE:HB	2:B:352:LEU:HD23	1.98	0.45
2:B:263:ASP:OD1	2:B:289:LYS:HE2	2.17	0.45
1:A:171:HIS:HE1	6:B:456:HOH:O	1.98	0.45
2:B:117:LEU:HD11	2:B:130:LEU:CD2	2.46	0.45
2:B:354:VAL:HG12	2:B:356:LEU:HG	1.99	0.45
2:B:255:TYR:CE2	2:B:307:GLN:NE2	2.85	0.45
2:B:285:LEU:HD23	2:B:285:LEU:HA	1.76	0.45
1:A:118:ARG:HE	1:A:118:ARG:HB3	1.48	0.44
2:B:322:LEU:C	2:B:322:LEU:HD23	2.38	0.44
1:A:164:THR:HG21	2:B:271:ASN:O	2.17	0.44
1:A:132:CYS:HB2	1:A:185:GLY:O	2.18	0.44
2:B:122:ASP:OD2	2:B:125:CYS:HB2	2.18	0.44
2:B:322:LEU:HD23	2:B:323:VAL:N	2.33	0.44
2:B:96:GLU:H	2:B:96:GLU:HG2	1.47	0.43
2:B:251:GLU:HA	2:B:254:LYS:HD3	1.99	0.43
1:A:210:GLU:O	1:A:241:PRO:HD2	2.18	0.43
2:B:222:LYS:HG2	6:B:402:HOH:O	2.18	0.43
2:B:72:ASP:HA	2:B:73:PRO:HD3	1.86	0.43
2:B:275:LEU:HD21	2:B:324:ASN:HD22	1.81	0.43
1:A:19:ILE:HG23	1:A:73:ALA:HB2	1.99	0.43
1:A:147:PRO:HG2	1:A:150:ILE:CD1	2.49	0.43
2:B:157:ASP:HB3	2:B:160:GLU:HG2	2.00	0.43
2:B:344:ARG:CG	2:B:344:ARG:NH1	2.80	0.43
1:A:30:PHE:CE2	2:B:329:ILE:HB	2.54	0.42
2:B:147:ASN:HB3	2:B:149:GLU:OE2	2.19	0.42
2:B:282:ILE:HG12	2:B:286:ASN:ND2	2.34	0.42
1:A:166:THR:CG2	1:A:183[B]:CYS:SG	3.07	0.42
2:B:138:VAL:CG1	2:B:139:ASP:N	2.82	0.42
1:A:2:SER:O	1:A:5:ALA:HB3	2.20	0.42
2:B:10:LYS:HD2	2:B:14:ASP:OD2	2.19	0.42
1:A:65:LYS:O	1:A:69:GLU:CG	2.65	0.42
1:A:256:ARG:NE	2:B:0:MET:HE2	2.35	0.42
2:B:133:SER:HA	2:B:134:PRO:HD3	1.76	0.41
1:A:94:ASP:OD1	1:A:118:ARG:NH2	2.44	0.41
2:B:90:THR:C	2:B:92:GLN:N	2.72	0.41
1:A:121:ARG:HH11	1:A:121:ARG:HG3	1.86	0.41
1:A:221:ASN:N	1:A:221:ASN:ND2	2.69	0.41
2:B:212:PRO:C	2:B:214:GLY:H	2.24	0.41
1:A:214:LEU:HD23	1:A:214:LEU:C	2.41	0.40
1:A:221:ASN:ND2	1:A:221:ASN:C	2.75	0.40
1:A:10:LEU:HD23	1:A:10:LEU:HA	1.93	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:138:VAL:HG12	2:B:139:ASP:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/311 (98%)	301 (99%)	4 (1%)	0	100	100
2	B	391/396 (99%)	381 (97%)	9 (2%)	1 (0%)	41	41
All	All	696/707 (98%)	682 (98%)	13 (2%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	91	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/243 (99%)	233 (97%)	7 (3%)	42	46
2	B	317/319 (99%)	299 (94%)	18 (6%)	20	18
All	All	557/562 (99%)	532 (96%)	25 (4%)	27	27

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	TYR
1	A	65	LYS
1	A	69	GLU
1	A	152	LYS
1	A	165	LEU
1	A	221	ASN
1	A	296	ILE
2	B	0	MET
2	B	10	LYS
2	B	71	LYS
2	B	112	SER
2	B	120	LEU
2	B	130	LEU
2	B	147	ASN
2	B	149	GLU
2	B	155	GLN
2	B	180	LEU
2	B	186	ASP
2	B	222	LYS
2	B	251	GLU
2	B	313	THR
2	B	326	PHE
2	B	337	ASN
2	B	344	ARG
2	B	355	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	171	HIS
1	A	221	ASN
1	A	225	ASN
1	A	233	HIS
1	A	279	GLN
2	B	92	GLN
2	B	168	GLN
2	B	181	GLN
2	B	182	ASN
2	B	215	GLN
2	B	224	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	245	ASN
2	B	307	GLN
2	B	360	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CME	B	332	2	8,9,10	0.73	0	5,9,11	1.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CME	B	332	2	-	1/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	332	CME	CZ-CE-SD-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	B	396	-	4,4,4	0.13	0	6,6,6	0.23	0
3	PO4	A	307	-	4,4,4	1.53	1 (25%)	6,6,6	0.42	0
4	SO4	A	308	-	4,4,4	0.20	0	6,6,6	0.16	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	307	PO4	P-O3	-2.05	1.48	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/311 (98%)	-0.72	0 100 100	10, 18, 39, 64	0
2	B	393/396 (99%)	-0.46	2 (0%) 91 92	12, 28, 56, 73	0
All	All	699/707 (98%)	-0.57	2 (0%) 94 94	10, 23, 51, 73	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	255	TYR	2.4
2	B	0	MET	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CME	B	332	10/11	0.95	0.08	28,30,40,41	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	A	308	5/5	0.92	0.19	32,32,35,35	5
4	SO4	B	396	5/5	0.96	0.22	24,25,28,29	5
5	ZN	B	400	1/1	0.96	0.09	34,34,34,34	0
3	PO4	A	307	5/5	0.99	0.08	22,24,31,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.