



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 18, 2024 – 12:52 PM EST

PDB ID : 4EZV
Title : Crystal structure of the substrate binding domain of E.coli DnaK in complex with PR-bombesin in space group P21212
Authors : Zahn, M.; Straeter, N.
Deposited on : 2012-05-03
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	 89% 11%
1	B	219	 92% 5% ..
2	C	17	 18% 71% 6% 24%
2	D	17	 6% 53% 12% 35%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3696 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chaperone protein DnaK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	219	Total	C	N	O	S	0	5	0
			1692	1041	298	345	8			
1	B	216	Total	C	N	O	S	0	4	0
			1660	1021	292	340	7			

- Molecule 2 is a protein called Proline rich bombesin-related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	13	Total	C	N	O	S	0	1	1
			108	72	22	13	1			
2	D	11	Total	C	N	O	S	0	1	1
			90	61	17	11	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	GLU	-	expression tag	UNP Q8QFP2
C	17	MET	-	expression tag	UNP Q8QFP2
D	1	GLU	-	expression tag	UNP Q8QFP2
D	17	MET	-	expression tag	UNP Q8QFP2


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	62	Total	O	0	0
			62	62		
3	B	81	Total	O	0	0
			81	81		
3	C	1	Total	O	0	0
			1	1		
3	D	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chaperone protein DnaK



- Molecule 1: Chaperone protein DnaK



- Molecule 2: Proline rich bombesin-related protein



- Molecule 2: Proline rich bombesin-related protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	65.13Å 148.76Å 64.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.10 24.50 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (25.00-2.10) 99.8 (24.50-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.210 , 0.259 0.206 , 0.256	Depositor DCC
R_{free} test set	1151 reflections (3.10%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtrriage
Anisotropy	0.247	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.042 for l,-k,h	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3696	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.90	0/1723	0.91	0/2320
1	B	0.88	0/1686	0.96	1/2270 (0.0%)
2	C	1.27	1/118 (0.8%)	0.90	0/161
2	D	1.08	0/99	1.06	0/135
All	All	0.91	1/3626 (0.0%)	0.94	1/4886 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	10	TRP	CD2-CE2	5.44	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	391	LEU	CA-CB-CG	5.18	127.21	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1692	0	1714	25	0
1	B	1660	0	1688	23	0
2	C	108	0	101	0	0
2	D	90	0	81	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	62	0	0	1	0
3	B	81	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	0	0
All	All	3696	0	3584	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:541:HIS:HE1	1:B:469:MET:HE1	1.29	0.98
1:B:467:ARG:HH22	1:B:541:HIS:HB2	1.39	0.87
1:A:541:HIS:CE1	1:B:469:MET:HE1	2.12	0.84
1:A:541:HIS:HE1	1:B:469:MET:CE	1.91	0.83
1:A:467:ARG:NH2	1:A:544[B]:HIS:ND1	2.27	0.82
1:A:541:HIS:CE1	1:B:469:MET:CE	2.74	0.70
1:B:404[A]:MET:HE2	1:B:467:ARG:HH11	1.60	0.67
1:B:404[A]:MET:HE1	1:B:467:ARG:HE	1.66	0.60
1:A:597:LYS:HE3	1:A:601:ILE:HD11	1.82	0.60
1:A:469[B]:MET:HE1	1:B:541:HIS:HD2	1.69	0.57
1:A:593:GLN:HA	1:B:492:ASN:ND2	2.21	0.56
1:A:469[B]:MET:CE	1:B:541:HIS:HD2	2.19	0.55
1:A:549:GLN:HE22	1:A:599:MET:HE1	1.74	0.53
1:A:389:VAL:HG13	3:A:741:HOH:O	2.07	0.53
1:A:469[B]:MET:CE	1:B:541:HIS:CD2	2.91	0.53
1:A:593:GLN:HA	1:B:492:ASN:HD22	1.75	0.51
1:B:404[A]:MET:CE	1:B:467:ARG:HH11	2.21	0.51
1:B:467:ARG:NH2	1:B:541:HIS:HB2	2.18	0.49
1:A:469[B]:MET:HE2	1:B:541:HIS:CD2	2.48	0.49
1:B:498:LYS:HE2	1:B:498:LYS:HB3	1.42	0.47
1:A:549:GLN:HE22	1:A:599:MET:CE	2.27	0.47
1:A:543:LEU:HD12	1:A:569:LEU:HD22	1.97	0.47
1:A:553:ALA:O	1:A:556:LYS:HB2	2.16	0.45
1:A:592:ALA:HB2	1:B:471:GLN:NE2	2.31	0.45
1:A:469[B]:MET:HE1	1:B:541:HIS:CD2	2.49	0.44
1:A:549:GLN:HE21	1:A:549:GLN:HB2	1.59	0.43
1:A:446:LYS:HE3	1:A:530:GLU:OE1	2.19	0.43
1:A:541:HIS:CE1	1:B:469:MET:HE3	2.53	0.42
1:A:595:SER:O	1:A:599:MET:HG2	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:543:LEU:HD12	1:A:569:LEU:CD2	2.50	0.42
1:B:547[B]:ARG:HG2	1:B:569:LEU:HD13	2.01	0.41
1:B:430:GLU:HB2	2:D:9:GLN:HG2	2.02	0.41
1:B:543:LEU:HD12	1:B:569:LEU:HD22	2.02	0.41
1:A:431:ASP:OD1	1:A:467:ARG:HG2	2.20	0.41
1:B:435:ALA:HB3	2:D:10:TRP:CZ3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/219 (101%)	219 (99%)	3 (1%)	0	100	100
1	B	218/219 (100%)	213 (98%)	4 (2%)	1 (0%)	29	26
2	C	12/17 (71%)	11 (92%)	1 (8%)	0	100	100
2	D	10/17 (59%)	10 (100%)	0	0	100	100
All	All	462/472 (98%)	453 (98%)	8 (2%)	1 (0%)	47	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	492	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/181 (103%)	181 (97%)	5 (3%)	44	48
1	B	183/181 (101%)	181 (99%)	2 (1%)	73	79
2	C	11/15 (73%)	11 (100%)	0	100	100
2	D	9/15 (60%)	9 (100%)	0	100	100
All	All	389/392 (99%)	382 (98%)	7 (2%)	57	65

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	391	LEU
1	A	510	ASP
1	A	555	ASP
1	A	596	GLN
1	A	605	GLN
1	B	391	LEU
1	B	604	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	415	ASN
1	A	432	ASN
1	A	541	HIS
1	A	549	GLN
1	A	603	GLN
1	B	471	GLN
1	B	492	ASN
1	B	513	GLN
1	B	541	HIS
1	B	603	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/219 (100%)	-0.02	3 (1%) 75 78	27, 42, 64, 79	0
1	B	216/219 (98%)	-0.03	5 (2%) 60 65	26, 40, 63, 88	0
2	C	13/17 (76%)	0.80	3 (23%) 0 0	29, 39, 74, 88	0
2	D	11/17 (64%)	0.33	1 (9%) 9 12	29, 37, 63, 64	0
All	All	459/472 (97%)	0.01	12 (2%) 56 61	26, 41, 64, 88	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	6	ARG	6.0
1	A	607	ALA	4.6
2	C	7	PRO	3.7
2	C	5	PRO	3.4
1	B	555	ASP	3.3
1	A	559	ALA	2.8
1	B	441	LEU	2.8
1	B	554	GLY	2.7
1	B	556	LYS	2.6
1	A	441	LEU	2.5
2	D	7	PRO	2.2
1	B	440	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.