



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 06:09 AM JST

PDB ID : 7F0V
Title : Structure of the M305I mutant of CueO
Authors : Kawano, T.K.; Takata, S.T.; Sakai, N.S.; Imaizumi, R.I.; Nakata, S.N.;
Takeshita, K.T.; Yamashita, S.Y.; Sakurai, T.S.; Kataoka, K.K.
Deposited on : 2021-06-07
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	516	 3% 84% 8% 8%
1	B	516	 3% 80% 11% 8%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7966 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blue copper oxidase CueO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	475	3695	2343	645	678	29	0	5	0
1	B	474	3658	2321	638	671	28	0	3	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	305	ILE	MET	engineered mutation	UNP W8T2X0
B	305	ILE	MET	engineered mutation	UNP W8T2X0

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	0
			2	2		
2	B	2	Total	Cu	0	0
			2	2		

- Molecule 3 is CU-O-CU LINKAGE (three-letter code: C2O) (formula: Cu₂O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Cu	O	0	0
			3	2	1		
3	B	1	Total	Cu	O	0	0
			3	2	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0

- Molecule 5 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total N O 4 1 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	314	Total O 314 314	0	0
6	B	257	Total O 257 257	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Blue copper oxidase CueO



- Molecule 1: Blue copper oxidase CueO



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.80Å 49.92Å 95.93Å 86.98° 80.63° 70.53°	Depositor
Resolution (Å)	47.37 – 1.49 47.33 – 1.49	Depositor EDS
% Data completeness (in resolution range)	95.1 (47.37-1.49) 95.1 (47.33-1.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.187 , 0.232 0.195 , 0.242	Depositor DCC
R_{free} test set	6644 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtrriage
Anisotropy	0.126	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7966	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: C2O, NO3, CU, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/3785	0.86	0/5134
1	B	0.69	0/3749	0.88	0/5090
All	All	0.70	0/7534	0.87	0/10224

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3695	0	3657	24	0
1	B	3658	0	3627	53	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	A	14	0	20	1	0
4	B	14	0	20	2	0
5	B	4	0	0	0	0
6	A	314	0	0	2	0
6	B	257	0	0	6	0
All	All	7966	0	7324	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:322[B]:ILE:CD1	1:B:322[B]:ILE:O	1.77	1.31
1:B:322[B]:ILE:O	1:B:322[B]:ILE:HD12	1.17	1.27
1:B:227:PRO:HA	1:B:322[B]:ILE:HD11	1.40	1.04
1:B:322[B]:ILE:CD1	1:B:322[B]:ILE:C	2.27	0.98
1:B:355:MET:HE2	6:B:1076:HOH:O	1.73	0.87
1:B:322[B]:ILE:HD12	1:B:322[B]:ILE:C	1.88	0.87
1:B:322[B]:ILE:O	1:B:322[B]:ILE:HD13	1.73	0.86
1:B:322[B]:ILE:C	1:B:322[B]:ILE:HD13	2.01	0.81
1:B:227:PRO:HA	1:B:322[B]:ILE:CD1	2.11	0.80
1:A:303[A]:MET:SD	1:A:364:MET:SD	2.80	0.80
1:A:380:ASP:H	1:A:383:GLN:HE21	1.28	0.79
1:B:355:MET:CE	1:B:505:HIS:CE1	2.69	0.75
1:A:355:MET:HG2	1:A:408:ASN:ND2	2.03	0.74
1:B:225:ALA:HB1	1:B:322[A]:ILE:HD11	1.71	0.71
1:B:355:MET:SD	6:B:1076:HOH:O	2.49	0.70
1:B:322[B]:ILE:HD13	1:B:324:ILE:HG12	1.72	0.69
1:B:85:ALA:HB2	1:B:130:ASN:HD22	1.57	0.69
1:B:55:GLY:HA3	1:B:72:ASN:HD22	1.59	0.68
1:B:100:LEU:O	1:B:113:GLY:HA3	1.96	0.66
1:A:380:ASP:H	1:A:383:GLN:NE2	1.94	0.64
1:B:85:ALA:HB2	1:B:130:ASN:ND2	2.12	0.64
1:B:355:MET:CE	6:B:1076:HOH:O	2.39	0.64
1:B:206:ALA:HA	1:B:359:LEU:HD21	1.79	0.63
1:B:355:MET:SD	1:B:439:ASP:OD2	2.58	0.62
1:B:141:HIS:HB2	1:B:142:PRO:HD2	1.83	0.61
1:B:228:ARG:HG3	1:B:322[A]:ILE:O	2.02	0.59
1:A:355:MET:HG2	1:A:408:ASN:HD22	1.67	0.59
1:A:202:VAL:HG21	1:A:362:MET:HB3	1.84	0.59
1:B:355:MET:HE1	1:B:505:HIS:CE1	2.38	0.58
1:B:355:MET:HG2	1:B:408:ASN:ND2	2.19	0.57
1:B:358:MET:SD	6:B:1149:HOH:O	2.56	0.57
1:B:499:HIS:HD2	1:B:500:CYS:O	1.88	0.56
1:A:227:PRO:HB2	1:A:326:ALA:HB2	1.87	0.56
1:B:289:ASP:HB2	1:B:291:LYS:HD3	1.87	0.56
1:B:199:GLN:HG2	1:B:199:GLN:O	2.07	0.54
1:A:225:ALA:HB1	1:A:322:ILE:HG13	1.91	0.53
1:B:450:THR:HG21	1:B:484:VAL:CG2	2.40	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:352:GLN:HG2	4:B:806:PEG:H42	1.91	0.52
1:B:242:ARG:HA	1:B:305:ILE:HG12	1.93	0.51
1:B:344:GLU:HA	1:B:344:GLU:OE2	2.11	0.50
1:A:450:THR:HG21	1:A:484:VAL:CG2	2.42	0.50
1:A:276:LEU:HD11	1:A:474:LYS:HB2	1.96	0.48
1:B:85:ALA:HA	1:B:129:LEU:O	2.14	0.48
1:A:227:PRO:HB2	1:A:326:ALA:CB	2.43	0.48
1:A:202:VAL:CG2	1:A:362:MET:HB3	2.44	0.47
1:B:55:GLY:HA3	1:B:72:ASN:ND2	2.26	0.47
1:B:288:ASN:ND2	1:B:291:LYS:HE3	2.30	0.47
1:A:251:ASN:OD1	4:A:805:PEG:H11	2.15	0.46
1:A:84:LYS:HE3	6:A:932:HOH:O	2.14	0.46
4:B:804:PEG:O1	4:B:804:PEG:H32	2.16	0.46
1:A:242:ARG:HA	1:A:305:ILE:HG12	1.97	0.46
1:B:99:THR:O	1:B:142:PRO:HA	2.16	0.45
1:B:147:LYS:NZ	6:B:910:HOH:O	2.48	0.45
1:B:512:LEU:C	1:B:512:LEU:HD12	2.38	0.44
1:A:428:GLN:HE22	1:B:84:LYS:C	2.21	0.44
1:B:47:ARG:HD3	1:B:89:ASP:OD2	2.18	0.44
1:B:102:TRP:CB	1:B:105:LEU:HD12	2.48	0.44
1:A:405:HIS:HE1	6:A:902:HOH:O	2.01	0.43
1:B:102:TRP:HB3	1:B:105:LEU:HD12	2.01	0.43
1:B:175:GLN:HG2	6:B:1116:HOH:O	2.18	0.43
1:B:358:MET:O	1:B:362:MET:HB2	2.19	0.43
1:A:227:PRO:CB	1:A:326:ALA:HB2	2.48	0.43
1:B:225:ALA:HA	1:B:320:GLN:O	2.19	0.43
1:B:379:MET:HA	1:B:379:MET:HE2	2.01	0.42
1:A:292:PRO:HB2	1:A:318[A]:ARG:HD2	2.01	0.42
1:A:431:ARG:NH1	1:A:481:GLU:OE2	2.48	0.42
1:A:42:THR:HB	1:A:46:ASN:HA	2.01	0.41
1:B:203:MET:HB2	1:B:366:MET:SD	2.60	0.41
1:B:64:ALA:HB2	1:B:194:GLY:O	2.20	0.41
1:B:355:MET:SD	1:B:441:MET:HG3	2.60	0.41
1:B:355:MET:CE	1:B:443:HIS:HE1	2.33	0.41
1:B:355:MET:HE2	1:B:505:HIS:NE2	2.36	0.41
1:A:360:ASP:O	1:A:364:MET:HG2	2.20	0.41
1:B:147:LYS:O	1:B:151:GLN:HG3	2.21	0.40
1:A:47:ARG:HA	1:A:87:THR:O	2.22	0.40
1:B:337:LEU:HD21	1:B:453:ARG:NH2	2.35	0.40
1:A:99:THR:O	1:A:142:PRO:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/516 (92%)	458 (97%)	16 (3%)	0	100	100
1	B	473/516 (92%)	458 (97%)	15 (3%)	0	100	100
All	All	947/1032 (92%)	916 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/422 (94%)	383 (97%)	13 (3%)	38	10
1	B	392/422 (93%)	384 (98%)	8 (2%)	55	25
All	All	788/844 (93%)	767 (97%)	21 (3%)	46	15

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	ARG
1	A	63[A]	THR
1	A	63[B]	THR
1	A	143	HIS
1	A	171	MET
1	A	174	LYS
1	A	280	ARG
1	A	288	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	350	LYS
1	A	362	MET
1	A	370	LYS
1	A	381	HIS
1	A	470	LYS
1	B	120	PRO
1	B	266	GLU
1	B	280	ARG
1	B	288	ASN
1	B	290	ASN
1	B	291	LYS
1	B	336	SER
1	B	380	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	ASN
1	A	103	HIS
1	A	130	ASN
1	A	199	GLN
1	A	223	GLN
1	A	383	GLN
1	A	408	ASN
1	A	428	GLN
1	B	56	GLN
1	B	72	ASN
1	B	103	HIS
1	B	130	ASN
1	B	223	GLN
1	B	288	ASN
1	B	408	ASN
1	B	499	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	B	806	-	6,6,6	0.35	0	5,5,5	0.22	0
4	PEG	A	804	-	6,6,6	0.25	0	5,5,5	0.20	0
4	PEG	B	804	-	6,6,6	0.37	0	5,5,5	0.43	0
3	C2O	B	803	1	0,2,2	-	-	-	-	-
4	PEG	A	805	-	6,6,6	0.30	0	5,5,5	0.21	0
5	NO3	B	805	-	1,3,3	0.11	0	0,3,3	-	-
3	C2O	A	803	1	0,2,2	-	-	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	B	806	-	-	2/4/4/4	-
4	PEG	A	805	-	-	3/4/4/4	-
4	PEG	A	804	-	-	1/4/4/4	-
4	PEG	B	804	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	804	PEG	O1-C1-C2-O2
4	A	804	PEG	O2-C3-C4-O4
4	A	805	PEG	O1-C1-C2-O2
4	B	806	PEG	O2-C3-C4-O4
4	A	805	PEG	O2-C3-C4-O4
4	B	804	PEG	O2-C3-C4-O4
4	B	806	PEG	O1-C1-C2-O2
4	A	805	PEG	C1-C2-O2-C3
4	B	804	PEG	C1-C2-O2-C3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	806	PEG	1	0
4	B	804	PEG	1	0
4	A	805	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	475/516 (92%)	-0.18	3 (0%) 89 91	22, 35, 57, 72	0
1	B	474/516 (91%)	0.07	13 (2%) 54 59	24, 41, 67, 86	0
All	All	949/1032 (91%)	-0.06	16 (1%) 70 75	22, 37, 63, 86	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	324	ILE	4.0
1	A	402	PHE	4.0
1	B	290	ASN	3.7
1	B	404	PHE	3.3
1	B	44	ALA	3.3
1	B	358	MET	3.2
1	B	323	ALA	3.2
1	B	380	ASP	3.2
1	B	289	ASP	2.8
1	B	355	MET	2.6
1	B	194	GLY	2.6
1	A	324	ILE	2.4
1	B	361	MET	2.3
1	A	388	MET	2.2
1	B	91	TYR	2.2
1	B	407	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PEG	B	804	7/7	0.70	0.15	54,60,61,62	0
4	PEG	B	806	7/7	0.80	0.24	45,51,61,62	0
4	PEG	A	804	7/7	0.81	0.14	56,62,66,67	0
4	PEG	A	805	7/7	0.83	0.24	47,61,63,67	0
5	NO3	B	805	4/4	0.86	0.10	59,63,63,67	0
2	CU	B	802	1/1	0.97	0.08	40,40,40,40	0
3	C2O	B	803	3/3	0.99	0.07	35,35,45,46	0
2	CU	B	801	1/1	0.99	0.12	53,53,53,53	0
3	C2O	A	803	3/3	0.99	0.05	29,29,37,39	0
2	CU	A	802	1/1	1.00	0.09	27,27,27,27	0
2	CU	A	801	1/1	1.00	0.10	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.