



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 19, 2023 – 12:34 PM EST

PDB ID : 4FA4
Title : Crystal Structure of WT MauG in Complex with Pre-Methylamine Dehydrogenase Aged 10 Days
Authors : Yukl, E.T.; Wilmot, C.M.
Deposited on : 2012-05-21
Resolution : 2.14 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	 5% 75% 20% 5%
1	B	373	 5% 81% 14% 2%
2	C	137	 5% 72% 22% 3%
2	E	137	 3% 72% 19% 8%
3	D	385	 5% 80% 17% 2%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	385	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into two segments: a green segment on the left labeled '82%' and a yellow segment on the right labeled '15%'. A small red square is at the start of the bar, and a small grey square is at the end. A '%' symbol is positioned above the start of the bar, and a '.' symbol is positioned below the end of the bar.</p>

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 14775 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylamine utilization protein MauG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	354	2813	1752	514	536	11	0	7	0
1	B	359	2781	1733	502	535	11	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	368	HIS	-	expression tag	UNP Q51658
A	369	HIS	-	expression tag	UNP Q51658
A	370	HIS	-	expression tag	UNP Q51658
A	371	HIS	-	expression tag	UNP Q51658
A	372	HIS	-	expression tag	UNP Q51658
A	373	HIS	-	expression tag	UNP Q51658
B	368	HIS	-	expression tag	UNP Q51658
B	369	HIS	-	expression tag	UNP Q51658
B	370	HIS	-	expression tag	UNP Q51658
B	371	HIS	-	expression tag	UNP Q51658
B	372	HIS	-	expression tag	UNP Q51658
B	373	HIS	-	expression tag	UNP Q51658

- Molecule 2 is a protein called Methylamine dehydrogenase light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	131	1017	628	179	196	14	0	1	0
2	E	126	973	601	163	194	15	0	3	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	132	HIS	-	expression tag	UNP P22619
C	133	HIS	-	expression tag	UNP P22619
C	134	HIS	-	expression tag	UNP P22619
C	135	HIS	-	expression tag	UNP P22619
C	136	HIS	-	expression tag	UNP P22619
C	137	HIS	-	expression tag	UNP P22619
E	132	HIS	-	expression tag	UNP P22619
E	133	HIS	-	expression tag	UNP P22619
E	134	HIS	-	expression tag	UNP P22619
E	135	HIS	-	expression tag	UNP P22619
E	136	HIS	-	expression tag	UNP P22619
E	137	HIS	-	expression tag	UNP P22619

- Molecule 3 is a protein called Methylamine dehydrogenase heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	376	Total	C	N	O	S	0	5	0
			2959	1874	506	570	9			
3	F	376	Total	C	N	O	S	0	4	0
			2957	1872	511	565	9			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is HEME C (three-letter code: HEC) (formula: C₃₄H₃₄FeN₄O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
5	A	1	43	34	1	4	4	0	0
5	A	1	43	34	1	4	4	0	0
5	B	1	43	34	1	4	4	0	0
5	B	1	43	34	1	4	4	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Na	0	0
			1	1		
8	B	2	Total	Na	0	0
			2	2		

- Molecule 9 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
9	B	1	5	4	1	0	0

- Molecule 10 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
10	D	1	4	2	2	0	0
10	F	1	4	2	2	0	0


- Molecule 11 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	179	Total O 181 181	0	2
11	B	258	Total O 260 260	0	2
11	C	61	Total O 61 61	0	0
11	D	184	Total O 184 184	0	0
11	E	87	Total O 87 87	0	0
11	F	293	Total O 294 294	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methylamine utilization protein MauG



- Molecule 1: Methylamine utilization protein MauG




- Molecule 2: Methylamine dehydrogenase light chain





- Molecule 2: Methylamine dehydrogenase light chain







- Molecule 3: Methylamine dehydrogenase heavy chain



- Molecule 3: Methylamine dehydrogenase heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	55.53Å 83.52Å 107.78Å 109.92° 91.52° 105.78°	Depositor
Resolution (Å)	43.46 – 2.14 37.17 – 2.14	Depositor EDS
% Data completeness (in resolution range)	97.4 (43.46-2.14) 97.5 (37.17-2.14)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 2.14Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.163 , 0.227 0.172 , 0.235	Depositor DCC
R_{free} test set	4717 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtrriage
Anisotropy	0.043	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14775	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 0AF, HEC, PO4, CA, PGE, ACT, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.90	0/2880	0.88	2/3902 (0.1%)
1	B	1.04	0/2845	0.95	5/3859 (0.1%)
2	C	0.94	1/1035 (0.1%)	0.90	1/1413 (0.1%)
2	E	1.00	0/988	0.96	1/1348 (0.1%)
3	D	0.87	0/3040	0.88	1/4142 (0.0%)
3	F	1.02	1/3037 (0.0%)	0.95	0/4136
All	All	0.96	2/13825 (0.0%)	0.92	10/18800 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	77	CYS	CB-SG	-5.79	1.72	1.81
3	F	365	GLU	CG-CD	5.39	1.60	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	39	ARG	NE-CZ-NH2	-9.98	115.31	120.30
1	B	39	ARG	NE-CZ-NH1	8.53	124.57	120.30
1	B	65	ARG	NE-CZ-NH1	-5.91	117.35	120.30
3	D	35	ARG	NE-CZ-NH2	-5.75	117.42	120.30
1	B	353	ARG	NE-CZ-NH1	5.56	123.08	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2813	0	2695	66	0
1	B	2781	0	2655	38	0
2	C	1017	0	906	30	0
2	E	973	0	876	17	0
3	D	2959	0	2831	43	0
3	F	2957	0	2841	34	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	86	0	61	8	0
5	B	86	0	62	13	0
6	A	10	0	14	1	0
7	A	4	0	6	0	0
7	B	4	0	6	0	0
8	A	1	0	0	0	0
8	B	2	0	0	0	0
9	B	5	0	0	0	0
10	D	4	0	3	0	0
10	F	4	0	3	0	0
11	A	181	0	0	6	0
11	B	260	0	0	3	0
11	C	61	0	0	2	0
11	D	184	0	0	8	0
11	E	87	0	0	1	0
11	F	294	0	0	2	0
All	All	14775	0	12959	212	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 212 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:34:CYS:SG	5:B:402:HEC:HAC	1.71	1.23
1:A:204:CYS:SG	5:A:403:HEC:HAC	1.81	1.16
1:A:133[B]:ARG:HB3	1:A:133[B]:ARG:NH1	1.64	1.12

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:CYS:SG	5:B:403:HEC:HAC	1.90	1.07
1:A:202[A]:ARG:HH22	2:C:127:VAL:HG11	1.31	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/373 (96%)	342 (95%)	18 (5%)	0	100	100
1	B	358/373 (96%)	344 (96%)	13 (4%)	1 (0%)	41	36
2	C	129/137 (94%)	123 (95%)	5 (4%)	1 (1%)	19	11
2	E	126/137 (92%)	123 (98%)	3 (2%)	0	100	100
3	D	379/385 (98%)	360 (95%)	16 (4%)	3 (1%)	19	11
3	F	378/385 (98%)	363 (96%)	13 (3%)	2 (0%)	29	22
All	All	1730/1790 (97%)	1655 (96%)	68 (4%)	7 (0%)	34	29

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	362	ARG
3	D	102[A]	ILE
3	D	102[B]	ILE
3	F	55	PHE
2	C	67	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/292 (97%)	273 (96%)	11 (4%)	32	29
1	B	280/292 (96%)	270 (96%)	10 (4%)	35	32
2	C	111/112 (99%)	108 (97%)	3 (3%)	44	43
2	E	107/112 (96%)	100 (94%)	7 (6%)	17	11
3	D	309/310 (100%)	299 (97%)	10 (3%)	39	37
3	F	308/310 (99%)	301 (98%)	7 (2%)	50	51
All	All	1399/1428 (98%)	1351 (97%)	48 (3%)	38	34

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	173	LYS
2	E	58	VAL
3	D	209	GLU
3	D	316	LEU
2	E	71	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	235	GLN
3	D	14	GLN
1	B	84	ASN
2	C	134	HIS
1	B	29	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	0AF	C	57	2	13,16,17	1.12	0	11,22,24	1.96	2 (18%)
2	0AF	E	57	2	13,16,17	1.09	0	11,22,24	1.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0AF	C	57	2	-	0/4/6/8	0/2/2/2
2	0AF	E	57	2	-	0/4/6/8	0/2/2/2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	57	0AF	CB-CA-C	4.87	120.60	111.47
2	C	57	0AF	CG-CB-CA	-2.03	111.39	114.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	57	0AF	5	0
2	E	57	0AF	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 5 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	B	405	-	3,3,3	0.47	0	2,2,2	0.14	0
9	PO4	B	404	-	4,4,4	0.72	0	6,6,6	0.71	0
10	ACT	F	401	-	3,3,3	0.94	0	3,3,3	1.45	0
5	HEC	A	403	1	32,50,50	1.76	9 (28%)	24,82,82	2.77	10 (41%)
7	EDO	A	405	-	3,3,3	0.47	0	2,2,2	0.38	0
6	PGE	A	404	-	9,9,9	0.64	0	8,8,8	0.47	0
10	ACT	D	401	-	3,3,3	0.78	0	3,3,3	1.04	0
5	HEC	A	402	1,11	32,50,50	1.59	9 (28%)	24,82,82	2.80	12 (50%)
5	HEC	B	402	1,11	32,50,50	1.62	8 (25%)	24,82,82	2.84	12 (50%)
5	HEC	B	403	1	32,50,50	1.50	5 (15%)	24,82,82	2.94	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	B	405	-	-	1/1/1/1	-
5	HEC	A	403	1	-	2/10/54/54	-
7	EDO	A	405	-	-	1/1/1/1	-
6	PGE	A	404	-	-	3/7/7/7	-
5	HEC	A	402	1,11	-	2/10/54/54	-
5	HEC	B	402	1,11	-	1/10/54/54	-
5	HEC	B	403	1	-	2/10/54/54	-

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	403	HEC	C3A-C4A	4.17	1.52	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	403	HEC	C3C-C4C	3.97	1.50	1.43
5	A	403	HEC	C3C-C4C	3.65	1.49	1.43
5	B	403	HEC	C3A-C4A	3.54	1.50	1.42
5	B	402	HEC	C1D-CHD	3.46	1.50	1.41

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	402	HEC	CMB-C2B-C3B	7.79	134.99	125.82
5	B	403	HEC	CMC-C2C-C3C	6.95	133.99	125.82
5	A	403	HEC	C1D-C2D-C3D	-6.44	102.52	107.00
5	A	403	HEC	CBD-CAD-C3D	-6.42	101.67	112.62
5	B	403	HEC	C1D-C2D-C3D	-6.41	102.54	107.00

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	404	PGE	O1-C1-C2-O2
7	B	405	EDO	O1-C1-C2-O2
7	A	405	EDO	O1-C1-C2-O2
5	A	402	HEC	CAD-CBD-CGD-O2D
6	A	404	PGE	C1-C2-O2-C3


There are no ring outliers.


5 monomers are involved in 22 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	403	HEC	5	0
6	A	404	PGE	1	0
5	A	402	HEC	3	0
5	B	402	HEC	7	0
5	B	403	HEC	6	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/373 (94%)	-0.14	2 (0%) 89 91	29, 39, 55, 71	0
1	B	359/373 (96%)	-0.24	4 (1%) 80 84	22, 33, 51, 87	0
2	C	130/137 (94%)	0.25	7 (5%) 25 32	23, 36, 65, 82	0
2	E	125/137 (91%)	-0.11	4 (3%) 47 55	23, 29, 42, 69	0
3	D	376/385 (97%)	0.03	19 (5%) 28 34	25, 42, 67, 77	0
3	F	376/385 (97%)	-0.21	3 (0%) 86 89	21, 30, 47, 61	0
All	All	1720/1790 (96%)	-0.11	39 (2%) 60 67	21, 35, 58, 87	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	ALA	5.4
1	B	6	ALA	5.4
2	C	106	ILE	4.6
3	D	135	VAL	3.6
3	D	207	GLY	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	0AF	C	57	15/16	0.94	0.23	39,43,47,49	0
2	0AF	E	57	15/16	0.95	0.19	31,34,38,40	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	PGE	A	404	10/10	0.70	0.39	78,86,90,90	0
7	EDO	A	405	4/4	0.92	0.17	59,59,60,60	0
7	EDO	B	405	4/4	0.95	0.08	44,46,47,48	0
8	NA	A	406	1/1	0.95	0.13	61,61,61,61	0
9	PO4	B	404	5/5	0.96	0.20	83,84,84,85	0
10	ACT	D	401	4/4	0.96	0.13	46,47,47,48	0
5	HEC	A	402	43/43	0.97	0.13	27,35,38,40	0
8	NA	B	406	1/1	0.97	0.07	38,38,38,38	0
5	HEC	A	403	43/43	0.98	0.13	27,31,35,38	0
8	NA	B	407	1/1	0.98	0.08	42,42,42,42	0
5	HEC	B	402	43/43	0.98	0.12	20,27,31,31	0
4	CA	A	401	1/1	0.98	0.12	33,33,33,33	0
10	ACT	F	401	4/4	0.98	0.13	42,44,46,46	0
5	HEC	B	403	43/43	0.99	0.17	16,22,26,29	0
4	CA	B	401	1/1	1.00	0.13	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEC A 402:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.