

# wwPDB X-ray Structure Validation Summary Report (i)

#### Feb 18, 2024 – 07:38 AM EST

PDB ID : 4FAA

Title : Structure of Recombinant Cytochrome ba3 Oxidase mutant A120F+A204F

from Thermus thermophilus

Authors: Li, Y.; Chen, Y.; Stout, C.D.

Deposited on : 2012-05-21

Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{\rm A})}) \end{array}$
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	568	69%	23%	• 6%
2	В	168	74%	22%	
3	С	34	53% 35%	•	9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	HAS	A	603	X	-	_	_



# 2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 6278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Cytochrome c oxidase subunit 1.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	٨	535	Total	С	N	О	S	0	0	0
1	A	999	4203	2863	663	661	16	0	0	

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	expression tag	UNP Q5SJ79
A	-4	HIS	-	expression tag	UNP Q5SJ79
A	-3	HIS	-	expression tag	UNP Q5SJ79
A	-2	HIS	-	expression tag	UNP Q5SJ79
A	-1	HIS	-	expression tag	UNP Q5SJ79
A	0	HIS	-	expression tag	UNP Q5SJ79
A	1	HIS	-	expression tag	UNP Q5SJ79
A	120	PHE	ALA	engineered mutation	UNP Q5SJ79
A	204	PHE	ALA	engineered mutation	UNP Q5SJ79

• Molecule 2 is a protein called Cytochrome c oxidase subunit 2.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	В	166	Total 1281	C 833	N 212	O 232	S 4	0	0	0

• Molecule 3 is a protein called Cytochrome c oxidase polypeptide 2A.


Mol	Chain	Residues		Aton	ıs		ZeroOcc	AltConf	Trace
3	С	31	Total 241	C 169	N 37	O 35	0	0	0

• Molecule 4 is COPPER (II) ION (three-letter code: CU) (formula: Cu).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cu 1 1	0	0

 $\bullet$  Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $\rm C_{34}H_{32}FeN_4O_4).$ 



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
5	A	1	Total	~ .	Fe	N	O	0	0
			43	34	1	4	4		


 $\bullet$  Molecule 6 is HEME-AS (three-letter code: HAS) (formula:  $\mathrm{C}_{54}\mathrm{H}_{64}\mathrm{FeN_4O_6}).$ 






Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
6	A	1	Total	С	Fe	N	O	0	0
			65	54	1	4	6		

 $\bullet$  Molecule 7 is HYDROGEN PEROXIDE (three-letter code: PEO) (formula:  $\mathrm{H}_2\mathrm{O}_2).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O 2 2	0	0

• Molecule 8 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula:  $C_{21}H_{40}O_4$ ).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O	0	0
			25 21 4		
8	A	1	Total C O 25 21 4	0	0
			25 21 4 Total C O		
8	A	1	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	0	0
0	٨	1	Total C O	0	0
8	A	1	21 19 2	0	0
8	A	1	Total C O	0	0
8	А	1	18 14 4	0	0
8	A	1	Total C O	0	0
	Λ	1	17 13 4	U	U
8	A	1	Total C O	0	0
		_	8 4 4		Ü
8	A	1	Total C O	0	0
			15 11 4		
8	A	1	Total C O	0	0
			20 16 4 Total C O		
8	A	1	$\begin{vmatrix} 10ta1 & C & O \\ 25 & 21 & 4 \end{vmatrix}$	0	0
			Total C O		
8	A	1	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	0	0
	Δ.		Total C O		0
8	A	1	25 21 4	0	0
8	В	1	Total C O	0	0
0	D	1	25 21 4	0	U
8	В	1	Total C O	0	0
	ט	1	25 21 4	U	U
8	В	1	Total C O	0	0
		1	25 21 4		
8	С	1	Total C O	0	0
	Ý		25 21 4		

 $\bullet$  Molecule 9 is DINUCLEAR COPPER ION (three-letter code: CUA) (formula: Cu2).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	В	1	Total Cu 2 2	0	0

#### • Molecule 10 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	51	Total O 51 51	0	0
10	В	44	Total O 44 44	0	0
10	С	2	Total O 2 2	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytochrome c oxidase subunit 1





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	143.28Å 98.29Å 95.05Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $127.50^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	56.84 - 2.80	Depositor
Resolution (A)	56.84 - 2.80	EDS
% Data completeness	97.2 (56.84-2.80)	Depositor
(in resolution range)	97.3 (56.84-2.80)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.65 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 2011_01_25_1225)	Depositor
$R, R_{free}$	0.194 , 0.254	Depositor
it, it free	0.190 , 0.249	DCC
$R_{free}$ test set	1289  reflections  (5.12%)	wwPDB-VP
Wilson B-factor ( $\mathring{A}^2$ )	37.1	Xtriage
Anisotropy	0.192	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37, 61.7	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.47, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6278	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.54% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, PEO, CUA, HEM, OLC, HAS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.50	0/4356	0.57	$2/5982 \ (0.0\%)$
2	В	0.53	0/1317	0.58	0/1801
3	С	0.48	0/247	0.52	0/335
All	All	0.51	0/5920	0.57	2/8118 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
1	A	131	VAL	N-CA-C	-5.83	95.26	111.00
1	A	369	PHE	N-CA-CB	-5.34	100.98	110.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	В	48	GLY	Peptide



# 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4203	0	4278	108	0
2	В	1281	0	1249	27	0
3	С	241	0	267	9	0
4	A	1	0	0	0	0
5	A	43	0	30	0	0
6	A	65	0	62	2	0
7	A	2	0	0	0	0
8	A	243	0	353	28	0
8	В	75	0	120	10	0
8	С	25	0	40	3	0
9	В	2	0	0	0	0
10	A	51	0	0	4	0
10	В	44	0	0	2	2
10	С	2	0	0	0	0
All	All	6278	0	6399	150	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 150 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} & & & & & & & & & & & & & & & & & & &$	Clash overlap (Å)
1:A:233:HIS:NE2	1:A:237:TYR:HE2	1.26	1.30
1:A:233:HIS:NE2	1:A:237:TYR:CE2	2.10	1.19
1:A:233:HIS:CD2	1:A:237:TYR:HE2	1.76	1.04
1:A:174:ASN:HB3	1:A:177:LYS:HD2	1.49	0.94
2:B:51:GLU:OE2	10:B:309:HOH:O	1.89	0.90

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
10:B:330:HOH:O	10:B:330:HOH:O[2_554]	1.94	0.26
10:B:308:HOH:O	10:B:324:HOH:O[2_554]	2.08	0.12



# 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	529/568 (93%)	489 (92%)	40 (8%)	0	100	100
2	В	164/168 (98%)	154 (94%)	10 (6%)	0	100	100
3	С	29/34 (85%)	24 (83%)	5 (17%)	0	100	100
All	All	722/770 (94%)	667 (92%)	55 (8%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	428/464~(92%)	407 (95%)	21 (5%)	25 57
2	В	132/138 (96%)	123 (93%)	9 (7%)	16 42
3	С	24/27 (89%)	19 (79%)	5 (21%)	1 3
All	All	584/629 (93%)	549 (94%)	35 (6%)	19 48

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	$\mathbf{Type}$	
2	В	167	LYS	
2	В	168	GLU	
3	С	13	LEU	
1	A 332		LEU	
1	A	326	LEU	



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	В	122	ASN
2	В	157	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

# 5.6 Ligand geometry (i)

Of 21 ligands modelled in this entry, 1 is monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	${ m gths}$	Во	ond angl	les
MIOI	Type	Chain	nes	tes Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
8	OLC	A	608	-	20,20,24	0.57	0	20,20,25	0.62	0
8	OLC	A	611	-	7,7,24	0.38	0	6,7,25	0.21	0
9	CUA	В	201	2	0,1,1	-	-	-		
8	OLC	A	605	-	24,24,24	0.48	0	25,25,25	0.77	0
8	OLC	В	203	-	24,24,24	0.57	1 (4%)	25,25,25	0.83	2 (8%)
8	OLC	A	612	-	14,14,24	0.56	0	15,15,25	0.73	0
8	OLC	С	101	-	24,24,24	0.53	0	25,25,25	0.57	0
8	OLC	A	616	-	24,24,24	0.55	1 (4%)	25,25,25	0.61	0
8	OLC	В	202	-	24,24,24	0.62	1 (4%)	25,25,25	0.68	0



Mol	Type	Chain	Res	Link	В	ond leng	gths	Во	ond angl	les
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
8	OLC	В	204	-	24,24,24	0.49	0	25,25,25	0.65	0
8	OLC	A	607	-	22,22,24	0.53	0	23,23,25	0.50	0
8	OLC	A	610	-	16,16,24	0.77	1 (6%)	17,17,25	0.86	0
7	PEO	A	604	4,6	1,1,1	0.52	0	-		
8	OLC	A	606	-	24,24,24	0.49	0	25,25,25	0.81	0
6	HAS	A	603	7,1	69,72,72	2.43	23 (33%)	73,109,109	1.42	9 (12%)
8	OLC	A	609	-	17,17,24	0.65	1 (5%)	18,18,25	0.76	0
8	OLC	A	615	-	20,20,24	0.51	0	21,21,25	0.70	0
8	OLC	A	614	-	24,24,24	0.57	1 (4%)	25,25,25	0.76	0
5	HEM	A	602	1	41,50,50	3.04	18 (43%)	45,82,82	2.13	11 (24%)
8	OLC	A	613	-	19,19,24	0.67	1 (5%)	20,20,25	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	OLC	A	612	-	-	5/14/14/24	-
8	OLC	A	615	-	-	8/20/20/24	-
8	OLC	A	608	-	-	8/19/19/24	-
8	OLC	A	611	-	-	1/6/6/24	-
8	OLC	С	101	-	-	19/24/24/24	-
8	OLC	A	605	-	-	10/24/24/24	-
8	OLC	A	606	-	-	13/24/24/24	-
8	OLC	A	616	-	-	11/24/24/24	-
8	OLC	В	202	-	-	9/24/24/24	-
8	OLC	В	203	-	-	9/24/24/24	-
6	HAS	A	603	7,1	1/1/8/18	15/40/82/82	-
8	OLC	В	204	-	-	12/24/24/24	-
8	OLC	A	607	-	-	7/22/22/24	-
8	OLC	A	610	-	-	9/16/16/24	-
8	OLC	A	614	-	-	12/24/24/24	-
5	HEM	A	602	1	-	3/12/54/54	-
8	OLC	A	609	-	-	7/17/17/24	-
8	OLC	A	613	-	-	10/19/19/24	-



The worst	5	of	48	bond	length	outliers	are	listed	below:
<b>110</b> 110100	$\overline{}$	01	10	OIIG	10115011	Cathern	COL C	IID CCC	CIC III.

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
6	A	603	HAS	C3C-C2C	7.84	1.51	1.40
6	A	603	HAS	C3C-CAC	-7.44	1.32	1.47
5	A	602	HEM	FE-NB	7.28	2.32	1.96
5	A	602	HEM	C3B-C2B	6.40	1.50	1.37
5	A	602	HEM	C3C-C2C	6.14	1.48	1.40

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
5	A	602	HEM	C1B-NB-C4B	7.56	112.88	105.07
5	A	602	HEM	CHC-C4B-NB	5.38	130.27	124.43
6	A	603	HAS	C1D-ND-C4D	-4.93	99.99	105.07
5	A	602	HEM	C4A-C3A-C2A	3.79	109.63	107.00
5	A	602	HEM	C4D-ND-C1D	3.78	108.98	105.07

#### All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	603	HAS	NA

#### 5 of 168 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	602	HEM	C2B-C3B-CAB-CBB
6	A	603	HAS	C1D-C2D-CMD-OMD
6	A	603	HAS	C3D-C2D-CMD-OMD
8	A	606	OLC	C21-C22-C24-O25
8	A	608	OLC	C2-C1-O20-C21

There are no ring outliers.

14 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	608	OLC	1	0
8	A	611	OLC	3	0
8	A	605	OLC	8	0
8	В	203	OLC	8	0
8	A	612	OLC	2	0
8	С	101	OLC	3	0
8	A	616	OLC	4	0
8	В	202	OLC	2	0


Continued on next page...




Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	607	OLC	1	0
8	A	606	OLC	4	0
6	A	603	HAS	2	0
8	A	609	OLC	1	0
8	A	614	OLC	3	0
8	A	613	OLC	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2	$OWAB(A^2)$	Q<0.9
1	A	535/568~(94%)	-0.32	14 (2%) 56 46	21, 29, 49, 77	1 (0%)
2	В	166/168 (98%)	-0.38	2 (1%) 79 73	22, 30, 46, 68	0
3	С	31/34 (91%)	-0.59	0 100 100	23, 28, 37, 49	0
All	All	732/770 (95%)	-0.34	16 (2%) 62 52	21, 29, 48, 77	1 (0%)

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	513	SER	4.7
1	A	322	PHE	3.6
1	A	60	PRO	3.1
1	A	14	ALA	3.1
1	A	13	GLU	2.9

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

# 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathrm{A}}^2)$	Q < 0.9
8	OLC	A	610	17/25	0.64	0.31	39,50,74,74	0
8	OLC	В	203	25/25	0.69	0.31	41,53,74,85	0
8	OLC	A	613	20/25	0.73	0.32	39,49,65,66	0
8	OLC	A	608	21/25	0.76	0.30	38,50,58,66	0
8	OLC	A	606	25/25	0.76	0.26	34,43,82,92	0
8	OLC	В	204	25/25	0.76	0.30	35,46,73,78	0
8	OLC	A	612	15/25	0.80	0.52	47,59,73,74	0
8	OLC	С	101	25/25	0.80	0.24	39,61,77,81	0
8	OLC	В	202	25/25	0.81	0.22	31,39,56,62	0
8	OLC	A	616	25/25	0.81	0.29	38,49,56,58	0
8	OLC	A	615	21/25	0.82	0.25	39,51,65,73	0
8	OLC	A	605	25/25	0.82	0.26	27,44,64,71	0
8	OLC	A	614	25/25	0.82	0.26	38,45,54,61	0
8	OLC	A	609	18/25	0.83	0.23	30,47,69,69	0
8	OLC	A	611	8/25	0.88	0.27	45,47,54,57	0
8	OLC	A	607	23/25	0.88	0.21	34,40,53,56	0
5	HEM	A	602	43/43	0.96	0.14	18,23,26,31	0
6	HAS	A	603	65/65	0.96	0.16	17,23,34,39	0
4	CU	A	601	1/1	0.99	0.12	25,25,25,25	0
7	PEO	A	604	2/2	0.99	0.12	19,19,19,25	0
9	CUA	В	201	2/2	0.99	0.11	21,21,21,24	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around OLC A 610:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around OLC B 203:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










# Electron density around OLC A 613: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)


# 







# Electron density around OLC B 204:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












# Electron density around OLC A 612:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around OLC C 101:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)
















# Electron density around OLC A 615:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around OLC A 605:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












# Electron density around OLC A 614: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around OLC A 609:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $mF_o$ -DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)












#### Electron density around OLC A 611:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around OLC A 607:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)
















# 6.5 Other polymers (i)

There are no such residues in this entry.

