



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 27, 2020 – 05:16 pm BST

PDB ID : 1FBA  
Title : THE CRYSTAL STRUCTURE OF FRUCTOSE-1,6-BISPHOSPHATE ALDOLASE FROM DROSOPHILA MELANOGASTER AT 2.5 ANGSTROMS RESOLUTION  
Authors : Piontek, K.; Hester, G.; Brenner-Holzach, O.  
Deposited on : 1992-06-08  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	 4% 85% 14% .
1	B	361	 4% 87% 12% ..
1	C	361	 9% 79% 19% ..
1	D	361	 5% 85% 12% ..

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12796 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	361	Total 2745	C 1728	N 480	O 531	S 6	0	0	0
1	B	361	Total 2745	C 1728	N 480	O 531	S 6	0	0	0
1	C	361	Total 2745	C 1728	N 480	O 531	S 6	0	0	0
1	D	361	Total 2745	C 1728	N 480	O 531	S 6	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	456	Total 456	O 456	0	0
2	B	477	Total 477	O 477	0	0
2	C	404	Total 404	O 404	0	0
2	D	479	Total 479	O 479	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE




- Molecule 1: FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE



- Molecule 1: FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE



- Molecule 1: FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.60Å 116.80Å 151.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.90 7.99 – 1.96	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-1.90) 95.4 (7.99-1.96)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.72 (at 1.97Å)	Xtrriage
Refinement program	PROLSQ, X-PLOR	Depositor
R, $R_{free}$	0.179 , (Not available) 0.162 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.2	Xtrriage
Anisotropy	0.661	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 72.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12796	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.79	0/2789	1.11	11/3779 (0.3%)
1	B	0.80	0/2789	1.10	8/3779 (0.2%)
1	C	0.76	0/2789	1.18	18/3779 (0.5%)
1	D	0.79	0/2789	1.15	12/3779 (0.3%)
All	All	0.78	0/11156	1.14	49/15116 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
1	D	0	3
All	All	0	5

There are no bond length outliers.

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	16	ARG	NE-CZ-NH1	12.18	126.39	120.30
1	C	330	ARG	NE-CZ-NH1	11.86	126.23	120.30
1	D	16	ARG	CD-NE-CZ	11.45	139.63	123.60
1	C	133	ARG	NE-CZ-NH1	10.76	125.68	120.30
1	C	16	ARG	NE-CZ-NH1	10.09	125.34	120.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	42	ARG	Sidechain
1	C	133	ARG	Sidechain
1	D	293	ARG	Sidechain
1	D	330	ARG	Sidechain
1	D	56	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2745	0	2769	39	0
1	B	2745	0	2769	36	0
1	C	2745	0	2769	59	0
1	D	2745	0	2769	46	0
2	A	456	0	0	10	0
2	B	477	0	0	7	0
2	C	404	0	0	9	0
2	D	479	0	0	13	0
All	All	12796	0	11076	179	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 179 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:VAL:HG21	1:B:362:ALA:HB1	1.37	1.02
1:B:330:ARG:HD2	2:B:615:HOH:O	1.64	0.98
1:D:359:ALA:O	1:D:360:ASN:HB2	1.69	0.92
1:A:325:ASN:HB3	2:A:550:HOH:O	1.68	0.92
1:C:152:LYS:HE2	1:C:362:ALA:HB3	1.55	0.88

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/361 (99%)	348 (97%)	9 (2%)	2 (1%)	25	15
1	B	359/361 (99%)	352 (98%)	5 (1%)	2 (1%)	25	15
1	C	359/361 (99%)	344 (96%)	13 (4%)	2 (1%)	25	15
1	D	359/361 (99%)	347 (97%)	9 (2%)	3 (1%)	19	9
All	All	1436/1444 (99%)	1391 (97%)	36 (2%)	9 (1%)	25	15

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	362	ALA
1	C	360	ASN
1	D	360	ASN
1	B	360	ASN
1	C	353	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/287 (100%)	281 (98%)	6 (2%)	53	48
1	B	287/287 (100%)	276 (96%)	11 (4%)	33	24
1	C	287/287 (100%)	273 (95%)	14 (5%)	25	15
1	D	287/287 (100%)	272 (95%)	15 (5%)	23	14
All	All	1148/1148 (100%)	1102 (96%)	46 (4%)	31	22

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	133	ARG
1	C	230	PRO
1	D	284	SER
1	C	164	LEU
1	C	193	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	254	GLN
1	B	319	ASN
1	D	72	ASN
1	B	161	GLN
1	B	237	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	360/361 (99%)	0.06	13 (3%) 42 45	2, 11, 42, 79	0
1	B	360/361 (99%)	0.02	15 (4%) 36 39	3, 12, 36, 89	0
1	C	360/361 (99%)	0.37	34 (9%) 8 9	3, 16, 55, 86	0
1	D	360/361 (99%)	0.05	18 (5%) 28 32	2, 11, 45, 86	0
All	All	1440/1444 (99%)	0.13	80 (5%) 24 27	2, 13, 45, 89	0

The worst 5 of 80 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	363	TYR	11.1
1	D	363	TYR	9.9
1	D	361	HIS	8.8
1	A	363	TYR	7.4
1	C	361	HIS	6.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.