



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 09:00 pm BST

PDB ID : 6FDO
Title : Rio2 structure
Authors : Fribourg, S.
Deposited on : 2017-12-26
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	 3% 64% 14% •• 19%
1	B	353	 5% 65% 13% • 20%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4675 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase RIO2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	2318	1476	397	430	15	0	0	0
1	B	284	2296	1464	390	427	15	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	33	Total	O	0	0
			33	33		
2	B	28	Total	O	0	0
			28	28		

4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	67.56 Å 67.56 Å 312.19 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.69 – 2.60 46.81 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.69-2.60) 100.0 (46.81-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.61 Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.195 , 0.250 0.162 , 0.199	Depositor DCC
R_{free} test set	1194 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	67.2	Xtrriage
Anisotropy	0.134	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 63.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtrriage
Estimated twinning fraction	0.324 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4675	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2369	0.76	3/3195 (0.1%)
1	B	0.51	0/2345	0.74	0/3162
All	All	0.51	0/4714	0.75	3/6357 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	70	VAL	C-N-CA	6.70	138.45	121.70
1	A	44	LEU	C-N-CA	5.56	135.61	121.70
1	A	45	LYS	C-N-CA	5.08	134.41	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	295	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2318	0	2275	26	2
1	B	2296	0	2257	24	0
2	A	33	0	0	1	0
2	B	28	0	0	0	0
All	All	4675	0	4532	48	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:LEU:O	1:A:298:VAL:N	2.16	0.76
1:B:167:ALA:HB1	1:B:298:VAL:HG23	1.71	0.70
1:A:111:ILE:HD11	1:A:119:GLN:HB3	1.80	0.64
1:A:296:LEU:N	1:A:296:LEU:HD23	2.13	0.63
1:B:67:THR:HG1	1:B:70:VAL:N	2.00	0.59
1:A:208:VAL:HG23	1:A:242:ILE:HD11	1.85	0.58
1:A:310:GLN:OE1	1:B:130:THR:HG21	2.04	0.57
1:A:293:GLU:HG3	1:A:294:ASP:H	1.70	0.56
1:B:65:GLU:O	1:B:71:GLN:HA	2.07	0.55
1:B:33:VAL:HG23	1:B:37:LEU:HD23	1.88	0.54
1:B:32:ILE:HD11	1:B:70:VAL:HG23	1.91	0.51
1:A:235:ILE:HG13	1:A:245:ILE:HG13	1.95	0.49
1:A:45:LYS:H	1:A:47:GLY:H	1.61	0.49
1:A:230:ASN:HB3	1:B:260:TRP:CZ2	2.48	0.48
1:A:262:PHE:CE1	1:A:290:ILE:HD11	2.48	0.48
1:B:208:VAL:HG23	1:B:242:ILE:HD11	1.96	0.48
1:A:33:VAL:HG23	1:A:37:LEU:HD23	1.96	0.47
1:A:26:GLY:O	1:A:29:ASN:O	2.33	0.47
1:B:244:MET:HG2	1:B:247:PHE:HD2	1.79	0.47
1:A:56:LEU:HB3	1:A:62:ILE:HG12	1.96	0.46
1:B:235:ILE:O	1:B:242:ILE:HA	2.16	0.46
1:B:168:LEU:O	1:B:173:PHE:HB2	2.15	0.46
1:B:167:ALA:CB	1:B:298:VAL:HG23	2.41	0.46
1:B:208:VAL:HG22	1:B:273:PHE:CZ	2.51	0.45
1:B:68:LYS:HB2	1:B:70:VAL:N	2.31	0.45
1:B:167:ALA:O	1:B:171:ARG:HD2	2.16	0.45
1:B:221:ASN:O	1:B:292:ARG:HA	2.17	0.45
1:B:159:MET:CE	1:B:310:GLN:HG3	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:MET:HA	1:A:309:MET:HE1	1.99	0.44
1:A:44:LEU:HB3	1:A:47:GLY:HA2	2.00	0.43
1:A:85:LEU:HD13	1:A:97:VAL:HG21	2.00	0.43
1:B:274:MET:SD	1:B:280:GLU:HB2	2.59	0.43
1:B:110:TYR:CE1	1:B:124:LEU:HD12	2.54	0.43
1:A:293:GLU:H	1:A:295:THR:HG23	1.84	0.42
1:B:159:MET:HG3	1:B:181:TYR:CZ	2.55	0.42
1:A:175:VAL:HG12	1:A:244:MET:HB3	2.01	0.42
1:A:200:HIS:HA	1:A:276:ARG:HD2	2.02	0.42
1:A:31:GLU:HG2	2:A:413:HOH:O	2.18	0.42
1:A:44:LEU:HB3	1:A:45:LYS:H	1.50	0.42
1:B:31:GLU:OE2	1:B:130:THR:HG23	2.19	0.42
1:A:274:MET:O	1:A:278:SER:HA	2.20	0.41
1:B:205:PRO:HA	1:B:208:VAL:HG13	2.03	0.41
1:A:205:PRO:HA	1:A:208:VAL:CG1	2.51	0.41
1:B:159:MET:HA	1:B:309:MET:HE1	2.02	0.41
1:A:27:MET:CE	1:A:147:ASN:HB3	2.51	0.41
1:B:27:MET:HB3	1:B:151:LEU:HD22	2.02	0.41
1:A:244:MET:HG2	1:A:247:PHE:HD2	1.85	0.40
1:A:296:LEU:CD2	1:A:296:LEU:N	2.84	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:ARG:CZ	1:A:294:ASP:OD1[5_665]	2.14	0.06
1:A:66:ARG:NE	1:A:294:ASP:OD1[5_665]	2.15	0.05

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/353 (80%)	261 (93%)	13 (5%)	8 (3%)	5	7
1	B	278/353 (79%)	252 (91%)	21 (8%)	5 (2%)	8	16
All	All	560/706 (79%)	513 (92%)	34 (6%)	13 (2%)	6	11

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	45	LYS
1	A	71	GLN
1	A	297	ASP
1	B	71	GLN
1	B	292	ARG
1	A	30	HIS
1	A	44	LEU
1	A	295	THR
1	A	48	GLY
1	B	67	THR
1	B	50	ASN
1	B	43	SER
1	A	47	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/312 (81%)	227 (90%)	25 (10%)	8	15
1	B	250/312 (80%)	230 (92%)	20 (8%)	12	24
All	All	502/624 (80%)	457 (91%)	45 (9%)	9	18

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	HIS
1	A	44	LEU
1	A	45	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	46	HIS
1	A	66	ARG
1	A	70	VAL
1	A	80	TYR
1	A	99	ASN
1	A	101	MET
1	A	115	GLU
1	A	118	GLN
1	A	159	MET
1	A	160	LYS
1	A	165	MET
1	A	204	ASP
1	A	238	GLU
1	A	247	PHE
1	A	274	MET
1	A	288	LYS
1	A	291	ARG
1	A	295	THR
1	A	297	ASP
1	A	301	SER
1	A	312	ASP
1	A	313	ASP
1	B	16	ASP
1	B	44	LEU
1	B	60	LYS
1	B	70	VAL
1	B	74	ARG
1	B	80	TYR
1	B	89	SER
1	B	95	GLU
1	B	115	GLU
1	B	130	THR
1	B	152	TYR
1	B	208	VAL
1	B	211	GLU
1	B	247	PHE
1	B	257	ASN
1	B	282	GLU
1	B	288	LYS
1	B	293	GLU
1	B	295	THR
1	B	297	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	147	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/353 (81%)	0.35	12 (4%) 36 29	51, 75, 114, 135	0
1	B	284/353 (80%)	0.43	16 (5%) 24 19	51, 79, 122, 150	0
All	All	570/706 (80%)	0.39	28 (4%) 29 23	51, 77, 120, 150	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	46	HIS	10.1
1	B	67	THR	5.3
1	B	295	THR	5.1
1	B	44	LEU	4.5
1	B	86	LYS	4.4
1	B	11	TYR	4.3
1	B	19	VAL	4.2
1	B	17	PHE	3.7
1	B	296	LEU	3.6
1	B	294	ASP	3.3
1	A	152	TYR	3.3
1	A	283	LEU	3.1
1	B	82	TYR	3.1
1	B	42	ALA	3.0
1	A	47	GLY	2.8
1	B	14	ARG	2.6
1	B	18	ARG	2.5
1	B	73	TYR	2.5
1	A	44	LEU	2.4
1	A	295	THR	2.4
1	A	18	ARG	2.4
1	A	66	ARG	2.4
1	B	152	TYR	2.3
1	A	14	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	82	TYR	2.1
1	B	15	ASP	2.1
1	A	11	TYR	2.1
1	A	17	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.