



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 6, 2024 – 08:50 pm GMT

PDB ID : 6FEA
Title : A. vinelandii vanadium nitrogenase, turnover state
Authors : Sippel, D.; Einsle, O.
Deposited on : 2017-12-31
Resolution : 1.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	 84% 12% ..
1	D	474	 84% 11% ..
2	B	475	 85% 12% ..
2	E	475	 85% 10% ..
3	C	113	 79% 17% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	113	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into segments: a small red segment at the beginning labeled '4%', a large green segment labeled '73%', and a yellow segment at the end labeled '20%'. There are three small black dots at the far right end of the bar.</p>

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 19493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitrogenase protein alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	461	3764	2403	643	692	26	0	8	0
1	D	461	3764	2403	643	692	26	0	8	0


- Molecule 2 is a protein called Vanadium nitrogenase beta subunit, vnfK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	464	3644	2315	626	682	21	0	2	0
2	E	464	3659	2323	628	687	21	0	4	0

- Molecule 3 is a protein called Vanadium nitrogenase, delta subunit, VnfG.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	113	953	598	171	181	3	0	1	0
3	F	111	947	594	172	179	2	0	2	0

- Molecule 4 is 3-HYDROXY-3-CARBOXY-ADIPIC ACID (three-letter code: HCA) (formula: C₇H₁₀O₇).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	O				
			14	7	7		0	0	
4	D	1	Total	C	O				
			14	7	7		0	0	

- Molecule 5 is FeV (three-letter code: D6N) (formula: CFe₇NS₇V).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
5	A	1	Total	C	Fe	N	S	V		
			17	1	7	1	7	1	0	0
5	D	1	Total	C	Fe	N	S	V		
			17	1	7	1	7	1	0	0

- Molecule 6 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 1 3	0	0
6	D	1	Total C O 4 1 3	0	0

- Molecule 7 is HYDROSULFURIC ACID (three-letter code: H2S) (formula: H₂S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total S 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	1	Total S 1 1	0	0

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Zn 1 1	0	0
8	B	1	Total Zn 1 1	0	0

- Molecule 9 is FE(8)-S(7) CLUSTER (three-letter code: CLF) (formula: Fe₈S₇).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Fe S 16 9 7	0	1
9	D	1	Total Fe S 16 9 7	0	1

- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	B	2	Total Mg 2 2	0	0
10	C	1	Total Mg 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	F	1	Total	Mg	0	0
			1	1		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	580	Total	O	0	0
			580	580		
11	B	562	Total	O	0	0
			562	562		
11	C	178	Total	O	0	0
			178	178		
11	D	570	Total	O	0	0
			570	570		
11	E	607	Total	O	0	0
			607	607		
11	F	155	Total	O	0	0
			155	155		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nitrogenase protein alpha chain



- Molecule 1: Nitrogenase protein alpha chain





- Molecule 2: Vanadium nitrogenase beta subunit, vnfK








- Molecule 2: Vanadium nitrogenase beta subunit, vnfK



- Molecule 3: Vanadium nitrogenase, delta subunit, VnfG



- Molecule 3: Vanadium nitrogenase, delta subunit, VnfG



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	75.61Å 79.75Å 107.16Å 84.05° 72.44° 75.25°	Depositor
Resolution (Å)	102.12 – 1.20 48.38 – 1.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (102.12-1.20) 99.6 (48.38-1.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 1.20Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.117 , 0.145 0.119 , 0.147	Depositor DCC
R_{free} test set	35693 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	10.7	Xtrriage
Anisotropy	0.050	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	19493	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, H2S, ZN, MG, D6N, HCA, CLF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.45	34/3859 (0.9%)	1.31	31/5213 (0.6%)
1	D	1.40	33/3859 (0.9%)	1.30	40/5213 (0.8%)
2	B	1.50	43/3722 (1.2%)	1.41	42/5039 (0.8%)
2	E	1.50	42/3737 (1.1%)	1.33	35/5059 (0.7%)
3	C	1.59	14/973 (1.4%)	1.33	12/1315 (0.9%)
3	F	1.92	21/967 (2.2%)	1.66	27/1307 (2.1%)
All	All	1.50	187/17117 (1.1%)	1.35	187/23146 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	1
2	B	0	1
2	E	0	2
3	F	0	1
All	All	0	6

The worst 5 of 187 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	82	GLU	CD-OE2	24.46	1.52	1.25
1	D	172	SER	CB-OG	-22.46	1.13	1.42
2	E	418	GLU	CD-OE1	17.38	1.44	1.25
1	D	366	GLU	CB-CG	-13.29	1.26	1.52
1	A	14	GLU	CD-OE2	12.82	1.39	1.25

The worst 5 of 187 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	466	ARG	NE-CZ-NH2	-23.73	108.43	120.30
3	F	56	ARG	NE-CZ-NH2	-16.57	112.01	120.30
3	F	56	ARG	NE-CZ-NH1	16.05	128.32	120.30
2	B	466	ARG	NE-CZ-NH1	15.96	128.28	120.30
2	B	418	GLU	OE1-CD-OE2	15.34	141.71	123.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	461	ASP	Sidechain
2	B	445	GLN	Sidechain
1	D	441	ARG	Sidechain
2	E	445	GLN	Sidechain
2	E	449	GLU	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3764	0	3673	28	0
1	D	3764	0	3673	20	0
2	B	3644	0	3616	7	0
2	E	3659	0	3625	15	0
3	C	953	0	916	13	0
3	F	947	0	909	7	0
4	A	14	0	6	1	0
4	D	14	0	6	1	0
5	A	17	0	0	1	0
5	D	17	0	0	1	0
6	A	4	0	0	0	0
6	D	4	0	0	0	0
7	A	1	0	0	0	0
7	D	1	0	0	0	0
8	A	1	0	0	0	0
8	B	1	0	0	1	0
9	A	16	0	0	0	0
9	D	16	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	B	2	0	0	0	0
10	C	1	0	0	0	0
10	F	1	0	0	0	0
11	A	580	0	0	5	0
11	B	562	0	0	6	0
11	C	178	0	0	1	1
11	D	570	0	0	4	1
11	E	607	0	0	6	1
11	F	155	0	0	2	1
All	All	19493	0	16424	89	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 89 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:12:ALA:N	2:E:12:ALA:CA	1.67	1.56
1:A:73:LEU:CD1	1:A:73:LEU:CG	1.76	1.55
1:A:195:MET:CG	1:A:195:MET:SD	2.02	1.48
2:E:13:GLU:CA	2:E:13:GLU:C	1.83	1.47
1:D:190:GLU:HG2	11:D:737:HOH:O	1.69	0.91

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:C:440:HOH:O	11:D:1015:HOH:O[1_644]	1.92	0.28
11:E:1091:HOH:O	11:F:434:HOH:O[1_545]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	467/474 (98%)	457 (98%)	10 (2%)	0	100	100
1	D	467/474 (98%)	457 (98%)	10 (2%)	0	100	100
2	B	464/475 (98%)	454 (98%)	10 (2%)	0	100	100
2	E	466/475 (98%)	457 (98%)	9 (2%)	0	100	100
3	C	112/113 (99%)	107 (96%)	5 (4%)	0	100	100
3	F	111/113 (98%)	108 (97%)	3 (3%)	0	100	100
All	All	2087/2124 (98%)	2040 (98%)	47 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	402/404 (100%)	397 (99%)	5 (1%)	71	37
1	D	402/404 (100%)	394 (98%)	8 (2%)	55	17
2	B	389/398 (98%)	387 (100%)	2 (0%)	88	67
2	E	391/398 (98%)	387 (99%)	4 (1%)	76	47
3	C	103/102 (101%)	100 (97%)	3 (3%)	42	7
3	F	102/102 (100%)	98 (96%)	4 (4%)	32	4
All	All	1789/1808 (99%)	1763 (98%)	26 (2%)	67	29

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	156	LYS
1	D	462	LYS
3	F	94[A]	ARG
1	D	259	ARG
2	E	16	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	54	GLN
2	E	402	GLN
3	F	46	GLN
3	C	46	GLN
3	C	80	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 2 are modelled with single atom and 6 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	CLF	A	506[A]	2	0,24,24	-	-	-		
9	CLF	A	506[B]	2	0,24,24	-	-	-		
6	CO3	A	503	5	2,3,3	0.63	0	2,3,3	1.30	0
6	CO3	D	503	5	2,3,3	0.60	0	2,3,3	0.82	0
9	CLF	D	505[B]	2	0,24,24	-	-	-		
4	HCA	A	501	5	13,13,13	1.93	3 (23%)	14,18,18	1.52	3 (21%)
4	HCA	D	502	5	13,13,13	1.34	2 (15%)	14,18,18	1.45	2 (14%)
5	D6N	A	502	1,6,4	6,28,28	1.56	1 (16%)	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	CLF	D	505[A]	2	0,24,24	-	-	-	-	-
5	D6N	D	501	1,6,4	6,28,28	1.46	1 (16%)	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	CLF	A	506[A]	2	-	-	0/12/10/10
9	CLF	A	506[B]	2	-	-	0/12/10/10
9	CLF	D	505[B]	2	-	-	0/12/10/10
4	HCA	A	501	5	-	2/17/17/17	-
4	HCA	D	502	5	-	2/17/17/17	-
9	CLF	D	505[A]	2	-	-	0/12/10/10

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	HCA	C3-C7	-4.98	1.48	1.53
4	D	502	HCA	C2-C3	-2.98	1.50	1.53
4	A	501	HCA	O1-C1	2.79	1.31	1.22
5	A	502	D6N	S4B-FE7	-2.61	2.22	2.31
5	D	501	D6N	S4B-FE7	-2.45	2.23	2.31

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	502	HCA	O5-C7-C3	-3.52	117.26	122.25
4	D	502	HCA	O6-C7-C3	3.39	118.94	113.05
4	A	501	HCA	O2-C1-C2	2.79	123.30	114.35
4	A	501	HCA	O6-C7-C3	2.78	117.87	113.05
4	A	501	HCA	O1-C1-C2	-2.77	114.83	122.94

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	502	HCA	C4-C5-C6-O4
4	D	502	HCA	C4-C5-C6-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	501	HCA	C4-C5-C6-O3
4	A	501	HCA	C4-C5-C6-O4

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	501	HCA	1	0
4	D	502	HCA	1	0
5	A	502	D6N	1	0
5	D	501	D6N	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	461/474 (97%)	-0.69	2 (0%) 92 92	9, 13, 24, 60	0
1	D	461/474 (97%)	-0.68	1 (0%) 95 94	7, 12, 24, 67	0
2	B	464/475 (97%)	-0.58	3 (0%) 89 90	8, 13, 27, 56	0
2	E	464/475 (97%)	-0.54	1 (0%) 95 94	7, 11, 24, 48	0
3	C	113/113 (100%)	-0.33	2 (1%) 68 68	12, 17, 36, 74	0
3	F	111/113 (98%)	-0.34	4 (3%) 42 41	11, 17, 52, 82	0
All	All	2074/2124 (97%)	-0.59	13 (0%) 89 90	7, 12, 28, 82	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	1	MET	4.4
2	B	12	ALA	4.2
3	F	79	VAL	3.9
3	F	84	ILE	3.9
2	B	376	ASN	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	HCA	A	501	14/14	0.99	0.04	7,9,13,14	0
4	HCA	D	502	14/14	0.99	0.07	6,7,12,13	0
5	D6N	A	502	17/17	1.00	0.04	6,9,9,9	0
5	D6N	D	501	17/17	1.00	0.05	5,7,8,8	0
6	CO3	A	503	4/4	1.00	0.05	9,9,9,9	0
6	CO3	D	503	4/4	1.00	0.06	7,7,8,8	0
7	H2S	A	504	1/1	1.00	0.04	17,17,17,17	0
7	H2S	D	504	1/1	1.00	0.03	16,16,16,16	0
8	ZN	A	505	1/1	1.00	0.04	16,16,16,16	1
8	ZN	B	503	1/1	1.00	0.06	22,22,22,22	1
9	CLF	A	506[A]	15/15	1.00	0.04	7,9,10,10	1
9	CLF	A	506[B]	15/15	1.00	0.04	9,9,10,18	1
9	CLF	D	505[A]	15/15	1.00	0.04	5,7,8,8	1
9	CLF	D	505[B]	15/15	1.00	0.04	7,7,8,13	1
10	MG	B	501	1/1	1.00	0.03	10,10,10,10	0
10	MG	B	502	1/1	1.00	0.03	9,9,9,9	0
10	MG	C	201	1/1	1.00	0.08	17,17,17,17	0
10	MG	F	201	1/1	1.00	0.07	15,15,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around D6N A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.