



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2024 – 11:44 PM EDT

PDB ID : 2FHD  
Title : Crystal structure of Crb2 tandem tudor domains  
Authors : Lee, J.; Botuyan, M.V.; Thompson, J.R.; Mer, G.  
Deposited on : 2005-12-23  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	 4% 61% 27% 8% . .
1	B	153	 13% 50% 24% 9% 8% 8%
1	C	153	 % 60% 20% 10% . 7%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA repair protein rhp9/CRB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	150	1230	788	213	223	6	0	4	0
1	B	140	1124	721	196	204	3	0	1	0
1	C	142	1137	733	193	207	4	0	1	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	355	GLY	-	insertion	GB 1449177
A	356	HIS	-	insertion	GB 1449177
A	357	MSE	-	insertion	GB 1449177
B	355	GLY	-	insertion	GB 1449177
B	356	HIS	-	insertion	GB 1449177
B	357	MSE	-	insertion	GB 1449177
C	355	GLY	-	insertion	GB 1449177
C	356	HIS	-	insertion	GB 1449177
C	357	MSE	-	insertion	GB 1449177

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 10 8 2	0	1
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 10 8 2	0	1
2	C	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	69	Total O 78 78	0	11
3	B	47	Total O 49 49	0	6

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	C	40	Total	O	0	4
			44	44		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA repair protein rhp9/CRB2



- Molecule 1: DNA repair protein rhp9/CRB2



- Molecule 1: DNA repair protein rhp9/CRB2



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.92Å 116.92Å 87.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.00 – 2.40 33.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.00-2.40) 99.9 (33.00-2.40)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	45.76 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.192 , 0.256 0.196 , 0.184	Depositor DCC
$R_{free}$ test set	1366 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.0	Xtrriage
Anisotropy	0.059	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 69.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3722	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.37	7/1249 (0.6%)	1.30	11/1676 (0.7%)
1	B	1.22	2/1143 (0.2%)	1.28	10/1541 (0.6%)
1	C	1.21	1/1156 (0.1%)	1.25	10/1556 (0.6%)
All	All	1.27	10/3548 (0.3%)	1.28	31/4773 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	12
1	C	0	4
All	All	0	18

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	458	ASP	CB-CG	-9.28	1.32	1.51
1	B	459	MSE	N-CA	9.25	1.64	1.46
1	A	496	LEU	N-CA	8.14	1.62	1.46
1	A	469	VAL	CB-CG2	-6.08	1.40	1.52
1	A	417	PHE	CD2-CE2	5.79	1.50	1.39

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	489	ARG	NE-CZ-NH2	-11.86	114.37	120.30
1	A	458	ASP	CB-CG-OD1	-11.44	108.01	118.30

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	484	ARG	NE-CZ-NH1	-9.19	115.71	120.30
1	B	455	LEU	N-CA-C	9.18	135.80	111.00
1	A	359	ARG	N-CA-C	-8.96	86.80	111.00

There are no chirality outliers.

5 of 18 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	358	SER	Peptide
1	A	373	TYR	Peptide
1	B	387	HIS	Peptide
1	B	390	VAL	Peptide
1	B	391	THR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1230	0	1258	51	0
1	B	1124	0	1150	77	0
1	C	1137	0	1162	46	0
2	A	25	0	0	0	0
2	B	10	0	0	1	0
2	C	25	0	0	0	0
3	A	78	0	0	6	1
3	B	49	0	0	3	0
3	C	44	0	0	2	1
All	All	3722	0	3570	166	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 166 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:405:MSE:SE	1:A:405:MSE:CE	2.15	1.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:357:MSE:SE	1:A:357:MSE:CE	2.15	1.43
1:A:499[A]:MSE:SE	1:A:499[A]:MSE:CE	2.16	1.43
1:B:395:MSE:CE	1:B:395:MSE:SE	2.26	1.33
1:A:388:SER:HB2	1:A:395:MSE:CE	1.67	1.23

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:121:HOH:O	3:C:122:HOH:O[3_665]	1.63	0.57

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	152/153 (99%)	140 (92%)	8 (5%)	4 (3%)	4	4
1	B	139/153 (91%)	116 (84%)	11 (8%)	12 (9%)	0	0
1	C	141/153 (92%)	129 (92%)	6 (4%)	6 (4%)	2	1
All	All	432/459 (94%)	385 (89%)	25 (6%)	22 (5%)	1	1

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	496	LEU
1	A	497	LYS
1	B	388	SER
1	B	391	THR
1	B	393	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/133 (105%)	117 (84%)	23 (16%)	2	2
1	B	127/133 (96%)	100 (79%)	27 (21%)	1	1
1	C	129/133 (97%)	110 (85%)	19 (15%)	2	3
All	All	396/399 (99%)	327 (83%)	69 (17%)	1	2

5 of 69 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	448	SER
1	C	456	ASN
1	C	489	ARG
1	B	386	VAL
1	B	382	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	464	HIS
1	C	456	ASN
1	B	456	ASN
1	C	434	HIS
1	B	444	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	A	307	-	4,4,4	0.82	0	6,6,6	0.81	0
2	PO4	C	305	-	4,4,4	1.06	0	6,6,6	0.45	0
2	PO4	C	302	-	4,4,4	0.78	0	6,6,6	0.68	0
2	PO4	A	301[B]	-	4,4,4	0.61	0	6,6,6	1.79	2 (33%)
2	PO4	A	312	-	4,4,4	1.36	1 (25%)	6,6,6	0.85	0
2	PO4	A	301[A]	-	4,4,4	0.93	0	6,6,6	1.32	0
2	PO4	B	311[B]	-	4,4,4	1.44	0	6,6,6	0.92	0
2	PO4	C	309	-	4,4,4	0.96	0	6,6,6	0.63	0
2	PO4	B	311[A]	-	4,4,4	0.72	0	6,6,6	0.70	0
2	PO4	C	303	-	4,4,4	1.38	1 (25%)	6,6,6	0.35	0
2	PO4	C	304	-	4,4,4	0.27	0	6,6,6	0.92	0
2	PO4	A	306	-	4,4,4	0.84	0	6,6,6	0.65	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	312	PO4	P-O1	2.30	1.56	1.50
2	C	303	PO4	P-O1	2.17	1.55	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301[B]	PO4	O4-P-O3	2.84	116.75	107.91
2	A	301[B]	PO4	O3-P-O1	-2.21	103.14	110.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	311[B]	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	145/153 (94%)	-0.32	6 (4%) 42 39	16, 30, 54, 66	3 (2%)
1	B	137/153 (89%)	0.35	20 (14%) 7 6	25, 40, 79, 88	1 (0%)
1	C	138/153 (90%)	-0.30	1 (0%) 84 81	22, 35, 58, 65	1 (0%)
All	All	420/459 (91%)	-0.10	27 (6%) 27 24	16, 35, 63, 88	5 (1%)

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	390	VAL	6.3
1	A	496	LEU	5.7
1	B	498	ASP	5.2
1	A	373	TYR	4.8
1	B	403	ALA	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	C	302	5/5	0.59	0.19	125,125,125,126	0
2	PO4	A	312	5/5	0.68	0.36	120,120,122,122	0
2	PO4	C	304	5/5	0.70	0.18	91,92,94,95	0
2	PO4	A	306	5/5	0.71	0.13	121,122,122,123	0
2	PO4	B	311[A]	5/5	0.80	0.23	57,60,61,62	5
2	PO4	B	311[B]	5/5	0.80	0.23	27,30,35,37	5
2	PO4	C	303	5/5	0.84	0.15	66,67,67,69	0
2	PO4	C	309	5/5	0.85	0.15	90,91,92,92	0
2	PO4	A	301[A]	5/5	0.86	0.23	54,56,57,59	5
2	PO4	C	305	5/5	0.86	0.14	90,91,92,93	0
2	PO4	A	301[B]	5/5	0.86	0.23	50,51,52,54	5
2	PO4	A	307	5/5	0.88	0.14	76,76,78,80	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.