

Full wwPDB X-ray Structure Validation Report (i)

Aug 20, 2023 - 03:49 AM EDT

PDB ID	:	2FHL
Title	:	avidin related protein (AVR4)-BNA complex
Authors	:	Livnah, O.; Prizant, M.
Deposited on		
Resolution	:	1.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
Clashscore	141614	1252 (1.10-1.02)
Ramachandran outliers	138981	1204 (1.10-1.02)
Sidechain outliers	138945	1202 (1.10-1.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	122	84%	13%	•••
1	В	122	84%	12%	•••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BNI	В	502	-	-	Х	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Avidin-related protein 4/5.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	120	Total	С	Ν	0	S	0	Б	0
	Л	120	986	621	179	181	5	0	5	0
1	В	120	Total	С	Ν	Ο	\mathbf{S}	0	Б	0
	D	120	985	622	176	182	5	0	5	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	122	SER	-	cloning artifact	UNP P56734
В	322	SER	-	cloning artifact	UNP P56734

• Molecule 2 is 5-(2-OXO-HEXAHYDRO-THIENO[3,4-D]IMIDAZOL-6-YL)-PENTANOIC ACID (4-NITRO-PHENYL)-AMIDE (three-letter code: BNI) (formula: C₁₆H₂₀N₄O₄S).




Mol	Chain	Residues		Atc	\mathbf{ms}			ZeroOcc	AltConf
2	А	1	Total	С	Ν	0	S	0	0
_		-	25	16	4	4	1	Ŭ	Ű



Mol	Chain	Residues		Ato	\mathbf{ms}			ZeroOcc	AltConf
2	Р	1	Total	С	Ν	0	S	0	0
	D	1	25	16	4	4	1	0	0

• Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 3 & 1 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 3 & 1 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 3 & 1 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 3 & 1 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 3 & 1 & 2 \end{array}$	0	0



• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	168	Total O 168 168	0	0
4	В	127	Total O 127 127	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

 \bullet Molecule 1: Avidin-related protein 4/5





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	78.04Å 78.04Å 110.03Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 - 1.05	Depositor
% Data completeness	100.0 (40.00-1.05)	Depositor
(in resolution range)	100.0 (40.00-1.05)	Depositor
R_{merge}	(Not available)	Depositor
R _{sym}	0.05	Depositor
Refinement program	REFMAC $5.1.24$	Depositor
R, R_{free}	0.167 , 0.177	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2346	wwPDB-VP
Average B, all atoms $(Å^2)$	14.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BNI, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.45	0/1012	0.75	3/1369~(0.2%)
1	В	0.44	0/1011	0.78	3/1368~(0.2%)
All	All	0.44	0/2023	0.77	6/2737~(0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	1	0
1	В	2	0
All	All	3	0

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	312[A]	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	В	312[B]	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	А	112[A]	ARG	NE-CZ-NH2	-5.62	117.49	120.30
1	А	112[B]	ARG	NE-CZ-NH2	-5.62	117.49	120.30
1	А	4	CYS	CA-CB-SG	5.54	123.98	114.00
1	В	239	ASP	CB-CG-OD2	5.54	123.29	118.30

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	А	61	THR	CB
1	В	261	THR	CB



Mol	Chain	Res	Type	Atom
1	В	278	THR	CB

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	986	0	964	23	0
1	В	985	0	970	24	0
2	А	25	0	20	8	0
2	В	25	0	20	9	0
3	А	12	0	4	1	0
3	В	18	0	6	1	0
4	А	168	0	0	1	0
4	В	127	0	0	1	0
All	All	2346	0	1984	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112[A]:ARG:HH22	2:A:501:BNI:C23	1.94	0.80
1:A:39:ASP:HB2	2:A:501:BNI:C23	2.14	0.78
1:B:312[A]:ARG:HH22	2:B:502:BNI:C22	1.98	0.76
1:A:80:GLN:HE22	1:B:301:VAL:H	1.32	0.75
1:A:39:ASP:HB2	2:A:501:BNI:C24	2.16	0.74
1:A:101:VAL:H	1:B:280:GLN:HE22	1.38	0.70
1:B:312[A]:ARG:HH22	2:B:502:BNI:C23	2.05	0.68
1:A:80:GLN:HE21	1:B:298:ARG:HH21	1.40	0.68
1:A:61:THR:HG23	1:B:274:THR:OG1	1.95	0.66
1:A:98:ARG:HH21	1:B:280:GLN:HE21	1.43	0.66
1:A:74:THR:OG1	1:B:261:THR:HG23	1.96	0.65
1:A:112[A]:ARG:HH22	2:A:501:BNI:C24	2.12	0.63



		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:B:239:ASP:HB2	2:B:502:BNI:C23	2.28	0.63
1:B:239:ASP:HB2	2:B:502:BNI:C24	2.31	0.60
1:B:312[A]:ARG:HH12	2:B:502:BNI:C23	2.15	0.58
1:A:54:HIS:HD2	4:B:522:HOH:O	1.88	0.56
4:A:532:HOH:O	1:B:254:HIS:HD2	1.88	0.56
3:A:405:FMT:O2	1:B:267:HIS:HE1	1.89	0.56
1:B:312[A]:ARG:HH12	2:B:502:BNI:H23	1.73	0.53
1:A:67:HIS:HE1	3:B:401:FMT:O1	1.91	0.52
1:A:112[A]:ARG:HH22	2:A:501:BNI:C22	2.22	0.52
1:B:262:PHE:O	1:B:278:THR:HG23	2.11	0.50
1:B:312[A]:ARG:NH2	2:B:502:BNI:C23	2.74	0.50
1:A:21:ARG:HH11	1:A:30:THR:HG23	1.77	0.48
1:B:204:CYS:N	1:B:283:ILE:HD11	2.28	0.48
1:A:98:ARG:HH21	1:B:280:GLN:NE2	2.13	0.47
1:A:6:LEU:H	1:A:53:GLN:HE22	1.63	0.45
1:A:54:HIS:HE1	1:B:273:SER:O	2.00	0.44
1:B:312[A]:ARG:HH22	2:B:502:BNI:C21	2.30	0.44
1:A:112[A]:ARG:HH12	2:A:501:BNI:H23	1.82	0.44
1:B:312[A]:ARG:NH1	2:B:502:BNI:C23	2.81	0.43
1:A:39:ASP:CB	2:A:501:BNI:C23	2.93	0.43
1:A:112[A]:ARG:HH12	2:A:501:BNI:C23	2.32	0.43
1:A:80:GLN:NE2	1:B:298:ARG:HH21	2.10	0.42
1:A:98:ARG:NH2	1:B:280:GLN:HE21	2.16	0.41
1:A:73:SER:O	1:B:254:HIS:HE1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentil	es
1	А	123/122~(101%)	122~(99%)	1 (1%)	0	100 100)



	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	В	123/122~(101%)	123 (100%)	0	0	100	100
All	All	246/244~(101%)	245 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percent	tiles
1	А	109/106~(103%)	104~(95%)	5(5%)	27	3
1	В	110/106 (104%)	106 (96%)	4 (4%)	35	4
All	All	219/212~(103%)	210~(96%)	9 (4%)	38	3

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	3	LYS
1	А	4	CYS
1	А	55	LYS
1	А	85[A]	ARG
1	А	85[B]	ARG
1	В	255	LYS
1	В	294[A]	MET
1	В	294[B]	MET
1	В	321	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such side chains are listed below:

Mol	Chain	Res	Type
1	А	53	GLN
1	А	54	HIS
1	А	67	HIS
1	А	80	GLN



Continued from previous page...

Mol	Chain	Res	Type
1	В	253	GLN
1	В	254	HIS
1	В	267	HIS
1	В	280	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain		Res	Link	Bo	ond leng	ths	Bond angles		
	Type	Ullain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	FMT	А	405	-	2,2,2	0.80	0	$1,\!1,\!1$	0.59	0
3	FMT	В	401	-	2,2,2	0.71	0	1,1,1	0.55	0
3	FMT	А	403	-	2,2,2	0.64	0	$1,\!1,\!1$	0.27	0
2	BNI	А	501	-	26,27,27	1.94	8 (30%)	$35,\!37,\!37$	1.29	4 (11%)
3	FMT	А	402	-	2,2,2	0.64	0	1,1,1	0.27	0
3	FMT	В	404	-	2,2,2	0.70	0	1,1,1	0.64	0
3	FMT	В	409	-	2,2,2	0.65	0	1,1,1	0.36	0
3	FMT	В	410	-	2,2,2	0.70	0	1,1,1	0.46	0
3	FMT	В	406	-	2,2,2	0.66	0	1,1,1	0.41	0
3	FMT	В	408	-	2,2,2	0.68	0	1,1,1	0.44	0



Mal	Mol Type Chain		Res	Link	Bond lengths			Bond angles		
IVIOI	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	BNI	В	502	-	26,27,27	2.00	8 (30%)	35,37,37	1.33	4 (11%)
3	FMT	А	407	-	2,2,2	0.69	0	1,1,1	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BNI	В	502	-	-	0/13/36/36	0/3/3/3
2	BNI	А	501	-	-	0/13/36/36	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
2	А	501	BNI	C24-C18	4.43	1.46	1.39
2	В	502	BNI	C21-C22	4.41	1.47	1.38
2	В	502	BNI	C24-C18	4.34	1.46	1.39
2	А	501	BNI	C21-C22	4.13	1.46	1.38
2	А	501	BNI	C21-C20	3.79	1.45	1.38
2	В	502	BNI	C21-C20	3.74	1.45	1.38
2	В	502	BNI	C22-N25	3.04	1.52	1.45
2	А	501	BNI	C22-N25	2.92	1.52	1.45
2	В	502	BNI	C23-C22	2.91	1.44	1.38
2	А	501	BNI	C20-C18	2.86	1.44	1.39
2	В	502	BNI	C20-C18	2.78	1.43	1.39
2	А	501	BNI	C23-C22	2.34	1.43	1.38
2	В	502	BNI	C18-N17	-2.18	1.37	1.41
2	А	501	BNI	C24-C23	2.17	1.42	1.38
2	В	502	BNI	C7-C2	2.15	1.57	1.52
2	А	501	BNI	C18-N17	-2.04	1.37	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	502	BNI	C10-C1-N17	3.50	120.75	114.59
2	А	501	BNI	C21-C22-N25	3.39	121.92	119.38
2	А	501	BNI	C10-C1-N17	3.16	120.17	114.59
2	А	501	BNI	C18-N17-C1	2.55	131.95	127.50
2	В	502	BNI	C6-C5-C4	2.26	110.62	108.66
2	В	502	BNI	C5-C6-S1	-2.09	104.52	106.31
2	В	502	BNI	C18-N17-C1	2.07	131.12	127.50



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	501	BNI	C2-C4-N2	2.05	114.96	113.13

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

4 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	405	FMT	1	0
3	В	401	FMT	1	0
2	А	501	BNI	8	0
2	В	502	BNI	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

