



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2024 – 04:08 pm GMT

PDB ID : 5FJO  
Title : N-acyl amino acid racemase from *Amycolatopsis* sp. Ts-1-60: G291D- F323Y mutant in complex with N-acetyl naphthylalanine  
Authors : Sanchez-Carron, G.; Campopiano, D.; Grogan, G.  
Deposited on : 2015-10-12  
Resolution : 2.08 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	 91% 7%
1	B	368	 93% 5%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6463 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called N-succinylamino acid racemase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	367	2770	1760	480	518	12	0	0	0
1	B	368	2776	1763	481	520	12	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	291	ASP	GLY	engineered mutation	UNP Q44244
A	323	TYR	PHE	engineered mutation	UNP Q44244
B	291	ASP	GLY	engineered mutation	UNP Q44244
B	323	TYR	PHE	engineered mutation	UNP Q44244

- Molecule 2 is N-acetyl naphthylalanine (three-letter code: NPQ) (formula: C<sub>15</sub>H<sub>15</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	15	1	3		
2	B	1	Total	C	N	O	0	0
			19	15	1	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	445	Total	O	0	0
			445	445		
4	B	432	Total	O	0	0
			432	432		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-succinylamino acid racemase



- Molecule 1: N-succinylamino acid racemase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	164.71Å 164.71Å 169.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.55 – 2.08 68.55 – 2.08	Depositor EDS
% Data completeness (in resolution range)	100.0 (68.55-2.08) 100.0 (68.55-2.08)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.99 (at 2.08Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.145 , 0.175 0.158 , 0.185	Depositor DCC
$R_{free}$ test set	3520 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.7	Xtrriage
Anisotropy	0.151	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 54.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.014 for -l,-k,-h 0.014 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6463	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NPQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.00	3/2826 (0.1%)	1.18	21/3850 (0.5%)
1	B	1.01	3/2832 (0.1%)	1.16	19/3858 (0.5%)
All	All	1.01	6/5658 (0.1%)	1.17	40/7708 (0.5%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	344	GLY	N-CA	7.17	1.56	1.46
1	A	122	GLU	CD-OE1	5.81	1.32	1.25
1	A	159	ARG	CD-NE	-5.21	1.37	1.46
1	A	30	GLU	CD-OE2	5.07	1.31	1.25
1	B	214	GLU	CD-OE1	5.03	1.31	1.25
1	B	122	GLU	CD-OE2	-5.01	1.20	1.25

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	159	ARG	NE-CZ-NH1	19.48	130.04	120.30
1	B	113	ARG	NE-CZ-NH2	-19.38	110.61	120.30
1	A	159	ARG	NE-CZ-NH1	18.45	129.52	120.30
1	B	113	ARG	NE-CZ-NH1	18.26	129.43	120.30
1	A	113	ARG	NE-CZ-NH1	18.11	129.35	120.30
1	A	113	ARG	NE-CZ-NH2	-17.42	111.59	120.30
1	A	159	ARG	NE-CZ-NH2	-16.07	112.27	120.30
1	B	159	ARG	NE-CZ-NH2	-15.93	112.33	120.30
1	A	10	ARG	NE-CZ-NH2	-12.32	114.14	120.30
1	A	10	ARG	NE-CZ-NH1	9.75	125.17	120.30
1	A	274	ARG	NE-CZ-NH2	-9.67	115.47	120.30
1	B	9	ARG	NE-CZ-NH2	-9.60	115.50	120.30
1	B	230	ARG	NE-CZ-NH1	8.82	124.71	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	274	ARG	NE-CZ-NH2	-8.63	115.98	120.30
1	A	9	ARG	NE-CZ-NH2	-8.57	116.01	120.30
1	A	113	ARG	CD-NE-CZ	7.93	134.71	123.60
1	B	343	THR	C-N-CA	-7.87	105.78	122.30
1	A	230	ARG	NE-CZ-NH1	7.75	124.17	120.30
1	B	113	ARG	CD-NE-CZ	7.37	133.91	123.60
1	A	159	ARG	CD-NE-CZ	6.52	132.73	123.60
1	A	127	ARG	NE-CZ-NH2	-6.41	117.09	120.30
1	A	343	THR	C-N-CA	-6.26	109.16	122.30
1	B	127	ARG	NE-CZ-NH2	-6.22	117.19	120.30
1	B	359	VAL	N-CA-CB	-6.16	97.95	111.50
1	B	97	ARG	NE-CZ-NH2	-6.03	117.28	120.30
1	B	159	ARG	CD-NE-CZ	5.97	131.95	123.60
1	B	231	ARG	NE-CZ-NH2	-5.74	117.43	120.30
1	B	274	ARG	NE-CZ-NH1	5.72	123.16	120.30
1	A	10	ARG	CG-CD-NE	-5.70	99.84	111.80
1	A	113	ARG	CG-CD-NE	-5.48	100.30	111.80
1	B	9	ARG	NE-CZ-NH1	5.45	123.03	120.30
1	B	321	ASP	CB-CG-OD1	5.42	123.18	118.30
1	B	97	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	128	ASP	CB-CG-OD1	5.25	123.02	118.30
1	B	113	ARG	CG-CD-NE	-5.21	100.87	111.80
1	A	231	ARG	NE-CZ-NH2	-5.17	117.72	120.30
1	A	8	LEU	CA-CB-CG	5.09	127.02	115.30
1	A	239	ASP	CB-CG-OD1	5.05	122.84	118.30
1	A	321	ASP	CB-CG-OD1	5.01	122.81	118.30
1	A	97	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2770	0	2806	20	0
1	B	2776	0	2811	16	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	19	0	0	0	0
2	B	19	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	445	0	0	5	1
4	B	432	0	0	12	2
All	All	6463	0	5617	36	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:ARG:HD2	1:A:32:LEU:HD23	1.54	0.89
1:B:274:ARG:NH2	4:B:2216:HOH:O	1.98	0.83
1:B:26:GLN:OE1	4:B:2043:HOH:O	1.98	0.82
1:B:11:VAL:HG13	1:B:359:VAL:HG13	1.64	0.78
1:A:8:LEU:HD12	1:A:366:ILE:CD1	2.15	0.76
1:A:10:ARG:HD2	1:A:32:LEU:CD2	2.17	0.75
1:A:8:LEU:HD12	1:A:366:ILE:HD11	1.70	0.72
1:B:233:GLN:NE2	4:B:2322:HOH:O	2.26	0.68
1:B:178:GLU:OE2	4:B:2284:HOH:O	2.13	0.67
1:A:7:GLU:CD	1:A:35:ARG:HH12	2.05	0.60
1:B:9:ARG:HD2	4:B:2013:HOH:O	2.01	0.59
1:A:7:GLU:OE2	1:A:35:ARG:NH1	2.35	0.59
1:A:7:GLU:CD	1:A:35:ARG:NH1	2.56	0.58
1:B:21:THR:OG1	4:B:2043:HOH:O	2.17	0.57
1:B:274:ARG:NE	4:B:2216:HOH:O	2.35	0.56
1:A:10:ARG:CD	1:A:32:LEU:CD2	2.84	0.55
1:B:274:ARG:CZ	4:B:2216:HOH:O	2.50	0.53
1:A:50:MET:HG3	4:A:2085:HOH:O	2.08	0.52
1:A:231:ARG:NH2	4:A:2352:HOH:O	2.42	0.51
1:A:8:LEU:HD12	1:A:366:ILE:HD13	1.94	0.49
1:B:231:ARG:NH2	4:B:2344:HOH:O	2.46	0.46
1:A:68:ARG:NH1	4:A:2118:HOH:O	2.27	0.46
1:A:127:ARG:CZ	4:A:2222:HOH:O	2.64	0.46
1:A:2:LYS:HE3	1:A:78:ALA:O	2.15	0.46
1:A:35:ARG:NE	1:A:42:GLU:OE2	2.44	0.45
1:B:50:MET:HG3	4:B:2099:HOH:O	2.16	0.44
1:B:8:LEU:HD23	1:B:8:LEU:C	2.38	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:ARG:HD2	1:B:187:GLN:OE1	2.18	0.44
1:A:127:ARG:NH2	4:A:2222:HOH:O	2.52	0.42
1:B:274:ARG:HD2	4:B:2365:HOH:O	2.20	0.41
1:A:159:ARG:HD2	1:A:187:GLN:OE1	2.20	0.41
1:A:134:VAL:HG23	1:A:322:ARG:HD3	2.02	0.41
1:A:237:CYS:HA	1:A:259:ILE:O	2.21	0.40
1:B:237:CYS:HA	1:B:259:ILE:O	2.21	0.40
1:A:171:GLU:OE2	1:A:174:ARG:NH1	2.55	0.40
1:B:113:ARG:NH2	4:B:2189:HOH:O	2.54	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2214:HOH:O	4:B:2217:HOH:O[3_555]	2.08	0.12
4:B:2027:HOH:O	4:B:2086:HOH:O[15_545]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	365/368 (99%)	357 (98%)	8 (2%)	0	100	100
1	B	366/368 (100%)	358 (98%)	8 (2%)	0	100	100
All	All	731/736 (99%)	715 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/292 (100%)	289 (99%)	2 (1%)	84	87
1	B	292/292 (100%)	290 (99%)	2 (1%)	84	87
All	All	583/584 (100%)	579 (99%)	4 (1%)	84	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	322	ARG
1	B	322	ARG
1	B	359	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NPQ	A	1368	3	20,20,20	1.78	5 (25%)	27,27,27	3.18	14 (51%)
2	NPQ	B	1369	3	20,20,20	1.97	6 (30%)	27,27,27	2.88	13 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NPQ	A	1368	3	-	3/12/12/12	0/2/2/2
2	NPQ	B	1369	3	-	2/12/12/12	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1369	NPQ	CAA-CAN	4.05	1.59	1.50
2	B	1369	NPQ	CB-CA	-3.77	1.44	1.54
2	A	1368	NPQ	CAF-CAI	3.36	1.44	1.36
2	A	1368	NPQ	CB-CA	-3.24	1.46	1.54
2	B	1369	NPQ	CAF-CAI	3.09	1.43	1.36
2	B	1369	NPQ	CAJ-CAQ	-3.08	1.34	1.41
2	A	1368	NPQ	CAJ-CAQ	-2.97	1.34	1.41
2	B	1369	NPQ	OXT-C	-2.91	1.21	1.30
2	A	1368	NPQ	CAA-CAN	2.84	1.56	1.50
2	B	1369	NPQ	CA-C	2.67	1.59	1.52
2	A	1368	NPQ	CA-C	2.38	1.58	1.52

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1368	NPQ	CB-CG-CD2	7.44	136.94	120.57
2	B	1369	NPQ	CB-CG-CD2	6.84	135.62	120.57
2	A	1368	NPQ	CB-CG-CD1	-6.65	107.70	120.91
2	B	1369	NPQ	CB-CG-CD1	-6.01	108.97	120.91
2	A	1368	NPQ	CAH-CAQ-CAR	5.59	128.94	118.92
2	B	1369	NPQ	CAH-CAQ-CAR	5.32	128.46	118.92
2	A	1368	NPQ	CAF-CAE-CAH	-4.09	114.71	120.44
2	A	1368	NPQ	CAI-CAR-CAQ	-4.06	111.64	118.92

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1368	NPQ	CAE-CAF-CAI	3.72	125.65	120.44
2	A	1368	NPQ	CG-CB-CA	3.60	123.32	113.39
2	B	1369	NPQ	CAJ-CAQ-CAH	-3.53	114.94	123.19
2	A	1368	NPQ	CAA-CAN-N	-3.40	110.34	116.10
2	B	1369	NPQ	CAI-CAR-CAQ	-3.32	112.97	118.92
2	A	1368	NPQ	CAJ-CAQ-CAH	-3.30	115.48	123.19
2	B	1369	NPQ	CD2-CAR-CAQ	3.21	123.54	118.96
2	B	1369	NPQ	CAE-CAF-CAI	3.21	124.93	120.44
2	A	1368	NPQ	OAB-CAN-N	3.11	127.67	121.95
2	B	1369	NPQ	CAJ-CD1-CG	2.98	127.28	121.14
2	A	1368	NPQ	CB-CA-N	2.97	117.06	110.79
2	A	1368	NPQ	CD2-CAR-CAQ	2.92	123.13	118.96
2	B	1369	NPQ	CAF-CAE-CAH	-2.84	116.45	120.44
2	B	1369	NPQ	OAB-CAN-N	2.80	127.09	121.95
2	B	1369	NPQ	CA-N-CAN	2.78	129.73	121.48
2	B	1369	NPQ	CG-CB-CA	2.76	120.99	113.39
2	A	1368	NPQ	CA-N-CAN	2.54	129.04	121.48
2	A	1368	NPQ	CAJ-CD1-CG	2.47	126.22	121.14
2	B	1369	NPQ	CB-CA-N	2.24	115.50	110.79

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1368	NPQ	C-CA-N-CAN
2	B	1369	NPQ	O-C-CA-CB
2	A	1368	NPQ	O-C-CA-CB
2	B	1369	NPQ	OXT-C-CA-CB
2	A	1368	NPQ	OXT-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	367/368 (99%)	-0.64	2 (0%) 91   92	11, 18, 34, 71	0
1	B	368/368 (100%)	-0.68	0 100   100	11, 17, 34, 62	0
All	All	735/736 (99%)	-0.66	2 (0%) 94   94	11, 18, 34, 71	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	40	ALA	4.3
1	A	79	GLU	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NPQ	A	1368	19/19	0.95	0.14	18,33,54,59	0
2	NPQ	B	1369	19/19	0.95	0.13	17,32,65,69	0
3	MG	A	1369	1/1	0.99	0.05	17,17,17,17	0


*Continued on next page...*




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	B	1370	1/1	0.99	0.10	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.