



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 7, 2020 – 02:02 am BST

PDB ID : 6FNH
Title : Crystal Structure of Ephrin A2 (EphA2) Receptor Protein Kinase with a pyr
azolo[3,4-d]pyrimidine fragment of NVP-BHG712
Authors : Kudlinzki, D.; Troester, A.; Witt, K.; Linhard, V.L.; Gande, S.L.; Saxena, K.;
Schwalbe, H.
Deposited on : 2018-02-04
Resolution : 1.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2907 (1.40-1.36)
Clashscore	141614	3037 (1.40-1.36)
Ramachandran outliers	138981	2970 (1.40-1.36)
Sidechain outliers	138945	2969 (1.40-1.36)
RSRZ outliers	127900	2846 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	 6% 85% 6% 9%
1	B	306	 5% 80% 8% 11%
1	C	306	 14% 84% 5% 12%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7371 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ephrin type-A receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	277	Total 2248	C 1441	N 386	O 402	S 19	0	5	0
1	B	271	Total 2187	C 1402	N 374	O 393	S 18	0	3	0
1	C	270	Total 2171	C 1394	N 372	O 387	S 18	0	2	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	595	GLY	-	expression tag	UNP P29317
B	595	GLY	-	expression tag	UNP P29317
C	595	GLY	-	expression tag	UNP P29317

- Molecule 2 is 1-methyl-6-pyridin-3-yl-pyrazolo[3,4-d]pyrimidin-4-amine (three-letter code: DXK) (formula: C₁₁H₁₀N₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			17	11	6		
2	B	1	Total	C	N	0	0
			17	11	6		
2	C	1	Total	C	N	0	0
			17	11	6		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	249	Total O 249 249	0	0
4	B	259	Total O 259 259	0	0
4	C	138	Total O 138 138	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ephrin type-A receptor 2



- Molecule 1: Ephrin type-A receptor 2



- Molecule 1: Ephrin type-A receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.50Å 90.67Å 200.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.95 – 1.38 49.21 – 1.38	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.95-1.38) 99.4 (49.21-1.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.95 (at 1.38Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, R_{free}	0.206 , 0.221 0.206 , 0.221	Depositor DCC
R_{free} test set	2216 reflections (1.05%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtrriage
Anisotropy	0.263	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7371	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DXK, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/2294	0.59	0/3087
1	B	0.40	0/2232	0.59	0/3006
1	C	0.38	0/2218	0.58	0/2988
All	All	0.38	0/6744	0.59	0/9081

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2248	0	2280	11	0
1	B	2187	0	2216	15	0
1	C	2171	0	2200	11	0
2	A	17	0	0	0	0
2	B	17	0	0	0	0
2	C	17	0	0	0	0
3	A	44	0	66	2	0
3	B	16	0	24	0	0
3	C	8	0	12	0	0
4	A	249	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	259	0	0	0	0
4	C	138	0	0	0	0
All	All	7371	0	6798	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:760:LEU:HD13	1:C:760:LEU:C	1.96	0.85
1:A:879:ASP:HA	1:A:882:LYS:HD2	1.59	0.84
1:A:661:LEU:HD11	1:A:688:MET:HE1	1.72	0.71
1:C:696[A]:GLU:HG3	1:C:749:SER:HB3	1.74	0.69
1:C:760:LEU:C	1:C:760:LEU:CD1	2.65	0.65
1:C:879:ASP:HA	1:C:882:LYS:HD2	1.81	0.62
1:C:782:ARG:NH1	1:C:821:LEU:O	2.30	0.61
1:B:879:ASP:OD1	1:B:882:LYS:NZ	2.37	0.58
1:B:661:LEU:HD11	1:B:688:MET:HE1	1.87	0.57
1:B:738:ARG:HD3	1:B:764:LEU:HB3	1.87	0.57
1:A:765:GLU:HB3	1:A:781:ILE:HD13	1.87	0.56
1:B:633:LYS:HA	1:B:640:GLU:HG2	1.89	0.55
1:A:683:SER:HA	1:A:688:MET:HE2	1.88	0.54
1:C:781:ILE:HG21	1:C:827:MET:SD	2.49	0.53
1:A:765:GLU:HG3	1:A:778:LYS:HD3	1.89	0.53
1:A:793:LYS:NZ	3:A:1005:EDO:H21	2.24	0.52
1:B:666:ILE:O	1:B:669:GLN:HG2	2.11	0.51
1:C:683:SER:HA	1:C:688:MET:HE3	1.93	0.49
1:C:818:TYR:O	1:C:821:LEU:HB2	2.13	0.49
1:B:661:LEU:HD21	1:B:688:MET:HE2	1.95	0.49
1:B:769:GLU:HG3	1:B:792:ARG:HG3	1.96	0.48
1:A:750:ASN:O	3:A:1012:EDO:H12	2.13	0.48
1:B:737:HIS:O	1:B:738:ARG:HB2	2.14	0.47
1:B:781:ILE:HD13	1:B:827:MET:HG3	1.95	0.47
1:B:872:ASP:O	1:B:876:ARG:HG3	2.14	0.47
1:B:617:LYS:HE3	1:B:617:LYS:HB3	1.73	0.45
1:A:666[B]:ILE:HD12	1:A:760:LEU:HD11	1.99	0.45
1:A:608:ILE:HD12	1:A:691:ILE:HD12	1.98	0.44
1:C:696[A]:GLU:H	1:C:696[A]:GLU:CD	2.21	0.43
1:B:791:TYR:HB2	1:B:793:LYS:HE2	2.00	0.43
1:C:696[A]:GLU:HG3	1:C:749:SER:CB	2.46	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:667:MET:HE2	1:A:678:LEU:HB2	2.02	0.42
1:B:723:ILE:HD13	1:B:745[B]:ILE:HD13	2.02	0.41
1:B:874:LEU:HB3	1:B:881:LEU:HD21	2.03	0.41
1:C:646:LYS:O	1:C:689:MET:HA	2.21	0.40
1:A:638:LYS:HG2	1:A:639:LYS:N	2.36	0.40
1:B:646:LYS:NZ	1:B:663:GLU:OE2	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/306 (90%)	273 (99%)	3 (1%)	0	100	100
1	B	268/306 (88%)	263 (98%)	5 (2%)	0	100	100
1	C	268/306 (88%)	259 (97%)	9 (3%)	0	100	100
All	All	812/918 (88%)	795 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/262 (93%)	243 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	237/262 (90%)	236 (100%)	1 (0%)	91	80
1	C	233/262 (89%)	233 (100%)	0	100	100
All	All	713/786 (91%)	712 (100%)	1 (0%)	93	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	764	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DXK	A	1001	-	17,19,19	2.99	6 (35%)	20,27,27	2.29	3 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DXK	B	1001	-	17,19,19	3.00	6 (35%)	20,27,27	2.44	4 (20%)
3	EDO	A	1010	-	3,3,3	0.46	0	2,2,2	0.62	0
3	EDO	B	1005	-	3,3,3	0.48	0	2,2,2	0.41	0
3	EDO	A	1005	-	3,3,3	0.37	0	2,2,2	0.27	0
3	EDO	C	1002	-	3,3,3	0.45	0	2,2,2	0.33	0
3	EDO	A	1012	-	3,3,3	0.46	0	2,2,2	0.34	0
3	EDO	A	1002	-	3,3,3	0.51	0	2,2,2	0.13	0
2	DXK	C	1001	-	17,19,19	3.24	6 (35%)	20,27,27	2.55	4 (20%)
3	EDO	A	1003	-	3,3,3	0.55	0	2,2,2	0.29	0
3	EDO	A	1006	-	3,3,3	0.49	0	2,2,2	0.27	0
3	EDO	A	1008	-	3,3,3	0.47	0	2,2,2	0.38	0
3	EDO	A	1007	-	3,3,3	0.47	0	2,2,2	0.34	0
3	EDO	A	1011	-	3,3,3	0.45	0	2,2,2	0.34	0
3	EDO	B	1004	-	3,3,3	0.49	0	2,2,2	0.24	0
3	EDO	B	1003	-	3,3,3	0.50	0	2,2,2	0.25	0
3	EDO	B	1002	-	3,3,3	0.51	0	2,2,2	0.39	0
3	EDO	C	1003	-	3,3,3	0.46	0	2,2,2	0.31	0
3	EDO	A	1009	-	3,3,3	0.46	0	2,2,2	0.34	0
3	EDO	A	1004	-	3,3,3	0.47	0	2,2,2	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DXK	A	1001	-	-	0/4/4/4	0/3/3/3
2	DXK	B	1001	-	-	0/4/4/4	0/3/3/3
3	EDO	A	1010	-	-	0/1/1/1	-
3	EDO	B	1005	-	-	0/1/1/1	-
3	EDO	A	1005	-	-	1/1/1/1	-
3	EDO	C	1002	-	-	0/1/1/1	-
3	EDO	A	1012	-	-	0/1/1/1	-
3	EDO	A	1002	-	-	1/1/1/1	-
2	DXK	C	1001	-	-	0/4/4/4	0/3/3/3
3	EDO	A	1003	-	-	0/1/1/1	-
3	EDO	A	1006	-	-	0/1/1/1	-
3	EDO	A	1008	-	-	0/1/1/1	-
3	EDO	A	1007	-	-	1/1/1/1	-
3	EDO	A	1011	-	-	1/1/1/1	-
3	EDO	B	1004	-	-	1/1/1/1	-
3	EDO	B	1003	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	1002	-	-	1/1/1/1	-
3	EDO	C	1003	-	-	1/1/1/1	-
3	EDO	A	1009	-	-	1/1/1/1	-
3	EDO	A	1004	-	-	0/1/1/1	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1001	DXK	C6-C5	-10.95	1.34	1.45
2	A	1001	DXK	C6-C5	-10.02	1.35	1.45
2	B	1001	DXK	C6-C5	-9.84	1.35	1.45
2	C	1001	DXK	C5-C4	-4.25	1.32	1.43
2	B	1001	DXK	C5-C4	-4.10	1.33	1.43
2	A	1001	DXK	C5-C4	-3.74	1.34	1.43
2	A	1001	DXK	CAB-NAD	3.51	1.41	1.34
2	C	1001	DXK	CAA-C2	-3.51	1.39	1.48
2	B	1001	DXK	CAA-C2	-3.43	1.39	1.48
2	C	1001	DXK	CAO-C5	-3.42	1.33	1.40
2	A	1001	DXK	CAA-C2	-3.21	1.40	1.48
2	B	1001	DXK	CAB-NAD	3.08	1.40	1.34
2	B	1001	DXK	CAO-C5	-2.99	1.34	1.40
2	C	1001	DXK	CAB-NAD	2.86	1.40	1.34
2	A	1001	DXK	CAO-C5	-2.61	1.35	1.40
2	B	1001	DXK	CAF-NAD	2.59	1.41	1.33
2	A	1001	DXK	CAF-NAD	2.41	1.40	1.33
2	C	1001	DXK	CAF-NAD	2.39	1.40	1.33

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1001	DXK	N3-C2-N1	-7.95	119.86	126.11
2	A	1001	DXK	N3-C2-N1	-7.47	120.24	126.11
2	C	1001	DXK	N3-C2-N1	-7.29	120.38	126.11
2	C	1001	DXK	CAO-NAN-NAM	5.38	109.83	104.23
2	C	1001	DXK	C2-N3-C4	4.21	120.33	115.08
2	A	1001	DXK	C2-N3-C4	3.95	120.00	115.08
2	B	1001	DXK	C2-N3-C4	3.90	119.95	115.08
2	B	1001	DXK	CAO-NAN-NAM	3.36	107.73	104.23
2	A	1001	DXK	CAO-NAN-NAM	3.13	107.49	104.23
2	C	1001	DXK	C5-C6-NAP	-2.43	117.77	121.54
2	B	1001	DXK	C6-N1-C2	2.11	119.57	116.63

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1005	EDO	O1-C1-C2-O2
3	A	1009	EDO	O1-C1-C2-O2
3	B	1002	EDO	O1-C1-C2-O2
3	C	1003	EDO	O1-C1-C2-O2
3	A	1002	EDO	O1-C1-C2-O2
3	B	1004	EDO	O1-C1-C2-O2
3	A	1007	EDO	O1-C1-C2-O2
3	A	1011	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1005	EDO	1	0
3	A	1012	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	277/306 (90%)	0.45	17 (6%) 21 21	25, 36, 73, 91	0
1	B	271/306 (88%)	0.45	15 (5%) 25 25	25, 35, 69, 92	0
1	C	270/306 (88%)	0.72	42 (15%) 2 1	31, 45, 83, 93	0
All	All	818/918 (89%)	0.54	74 (9%) 9 10	25, 39, 76, 93	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	634	THR	8.3
1	C	760	LEU	7.6
1	B	602	LEU	6.6
1	C	638	LYS	6.5
1	A	604	PHE	6.0
1	A	763	VAL	5.8
1	C	828	LYS	5.7
1	C	781	ILE	5.7
1	B	766	ASP	5.7
1	C	639	LYS	5.5
1	A	819	TRP	5.5
1	C	780	PRO	5.4
1	C	637	GLY	5.3
1	C	624	PHE	5.3
1	B	765	GLU	5.3
1	A	602	LEU	5.2
1	C	889	PRO	5.0
1	B	634	THR	4.8
1	C	820	GLU	4.8
1	C	821	LEU	4.6
1	C	819	TRP	4.5
1	A	766	ASP	4.2
1	C	827	MET	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	762	ARG	4.1
1	C	686	LYS	4.1
1	C	830	ILE	4.0
1	A	760	LEU	4.0
1	C	791	TYR	4.0
1	C	823	ASN	3.9
1	C	759	GLY	3.9
1	C	824	HIS	3.9
1	B	635	SER	3.9
1	A	888	ASP	3.8
1	C	826	VAL	3.8
1	C	658	VAL	3.7
1	C	685	TYR	3.7
1	B	763	VAL	3.6
1	C	822	SER	3.4
1	C	655	LYS	3.3
1	C	831	ASN	3.3
1	C	789	ILE	3.3
1	C	688	MET	3.2
1	C	834	PHE	3.1
1	C	735	TYR	3.1
1	C	623	GLU	3.0
1	C	758	PHE	3.0
1	A	889	PRO	2.9
1	C	654	GLU	2.9
1	A	781	ILE	2.9
1	A	637	GLY	2.9
1	B	778	LYS	2.9
1	A	820	GLU	2.8
1	B	876	ARG	2.8
1	A	761	SER	2.8
1	A	638	LYS	2.7
1	C	687	PRO	2.7
1	A	764	LEU	2.7
1	B	760	LEU	2.6
1	C	661	LEU	2.5
1	C	651	GLY	2.5
1	A	779	ILE	2.3
1	B	768	PRO	2.3
1	B	764	LEU	2.3
1	B	780	PRO	2.2
1	C	794	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	779	ILE	2.2
1	C	890	ARG	2.2
1	A	605	THR	2.1
1	C	860	ARG	2.1
1	C	786	PRO	2.1
1	C	684	LYS	2.1
1	A	762	ARG	2.1
1	B	628	TYR	2.1
1	C	609	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	B	1003	4/4	0.55	0.12	69,70,72,74	0
3	EDO	A	1007	4/4	0.62	0.21	66,66,68,69	0
3	EDO	B	1005	4/4	0.64	0.16	67,71,71,72	0
3	EDO	A	1011	4/4	0.66	0.22	50,51,57,60	0
3	EDO	A	1003	4/4	0.70	0.18	49,52,53,53	0
3	EDO	A	1002	4/4	0.72	0.12	51,55,59,63	0
3	EDO	A	1012	4/4	0.74	0.17	34,47,49,50	0
3	EDO	B	1004	4/4	0.77	0.16	47,52,56,62	0
3	EDO	C	1002	4/4	0.79	0.15	58,58,60,65	0
3	EDO	A	1010	4/4	0.81	0.12	39,44,45,49	0
3	EDO	A	1005	4/4	0.81	0.17	56,58,61,62	0
3	EDO	A	1006	4/4	0.84	0.09	52,57,57,61	0
3	EDO	A	1008	4/4	0.84	0.09	60,61,61,62	0
3	EDO	A	1004	4/4	0.85	0.24	56,57,59,61	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	B	1002	4/4	0.86	0.16	41,47,51,56	0
3	EDO	A	1009	4/4	0.88	0.12	61,65,66,69	0
2	DXK	A	1001	17/17	0.92	0.07	28,35,43,47	0
2	DXK	B	1001	17/17	0.93	0.07	29,36,44,49	0
2	DXK	C	1001	17/17	0.93	0.16	39,52,59,65	0
3	EDO	C	1003	4/4	0.94	0.05	57,64,68,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.