



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 5, 2021 – 03:18 pm BST

PDB ID : 5FPH
Title : The GTPase domains of the immunity-related Irga6 dimerize in a parallel head-to-head fashion
Authors : Schulte, K.; Pawlowski, N.; Faelber, K.; Froehlich, C.; Howard, J.; Daumke, O.
Deposited on : 2015-11-30
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	
1	B	423	
1	C	423	
1	D	423	
1	E	423	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	423	<p>3% 56% 36% 7%</p>
1	G	423	<p>2% 52% 40% 6%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GNP	A	501	-	-	X	-
2	GNP	C	501	-	-	X	-
2	GNP	E	501	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 22647 atoms, of which 24 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTERFERON-INDUCIBLE GTPASE 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	395	3204	2062	522	606	4	10	0	0	0
1	B	396	3204	2066	518	606	4	10	0	0	0
1	C	389	3153	2033	514	592	4	10	0	0	0
1	D	396	3193	2051	521	607	4	10	0	0	0
1	E	396	3205	2065	521	605	4	10	0	0	0
1	F	394	3194	2055	521	604	4	10	0	0	0
1	G	399	3241	2086	527	614	4	10	0	0	0

There are 98 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	GLY	-	expression tag	UNP Q9QZ85
A	-8	SER	-	expression tag	UNP Q9QZ85
A	-7	PRO	-	expression tag	UNP Q9QZ85
A	-6	GLY	-	expression tag	UNP Q9QZ85
A	-5	ILE	-	expression tag	UNP Q9QZ85
A	-4	PRO	-	expression tag	UNP Q9QZ85
A	-3	GLY	-	expression tag	UNP Q9QZ85
A	-2	SER	-	expression tag	UNP Q9QZ85
A	-1	THR	-	expression tag	UNP Q9QZ85
A	0	THR	-	expression tag	UNP Q9QZ85
A	31	GLU	ARG	engineered mutation	UNP Q9QZ85
A	32	GLU	LYS	engineered mutation	UNP Q9QZ85
A	176	GLU	LYS	engineered mutation	UNP Q9QZ85
A	246	GLU	LYS	engineered mutation	UNP Q9QZ85
B	-9	GLY	-	expression tag	UNP Q9QZ85

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-8	SER	-	expression tag	UNP Q9QZ85
B	-7	PRO	-	expression tag	UNP Q9QZ85
B	-6	GLY	-	expression tag	UNP Q9QZ85
B	-5	ILE	-	expression tag	UNP Q9QZ85
B	-4	PRO	-	expression tag	UNP Q9QZ85
B	-3	GLY	-	expression tag	UNP Q9QZ85
B	-2	SER	-	expression tag	UNP Q9QZ85
B	-1	THR	-	expression tag	UNP Q9QZ85
B	0	THR	-	expression tag	UNP Q9QZ85
B	31	GLU	ARG	engineered mutation	UNP Q9QZ85
B	32	GLU	LYS	engineered mutation	UNP Q9QZ85
B	176	GLU	LYS	engineered mutation	UNP Q9QZ85
B	246	GLU	LYS	engineered mutation	UNP Q9QZ85
C	-9	GLY	-	expression tag	UNP Q9QZ85
C	-8	SER	-	expression tag	UNP Q9QZ85
C	-7	PRO	-	expression tag	UNP Q9QZ85
C	-6	GLY	-	expression tag	UNP Q9QZ85
C	-5	ILE	-	expression tag	UNP Q9QZ85
C	-4	PRO	-	expression tag	UNP Q9QZ85
C	-3	GLY	-	expression tag	UNP Q9QZ85
C	-2	SER	-	expression tag	UNP Q9QZ85
C	-1	THR	-	expression tag	UNP Q9QZ85
C	0	THR	-	expression tag	UNP Q9QZ85
C	31	GLU	ARG	engineered mutation	UNP Q9QZ85
C	32	GLU	LYS	engineered mutation	UNP Q9QZ85
C	176	GLU	LYS	engineered mutation	UNP Q9QZ85
C	246	GLU	LYS	engineered mutation	UNP Q9QZ85
D	-9	GLY	-	expression tag	UNP Q9QZ85
D	-8	SER	-	expression tag	UNP Q9QZ85
D	-7	PRO	-	expression tag	UNP Q9QZ85
D	-6	GLY	-	expression tag	UNP Q9QZ85
D	-5	ILE	-	expression tag	UNP Q9QZ85
D	-4	PRO	-	expression tag	UNP Q9QZ85
D	-3	GLY	-	expression tag	UNP Q9QZ85
D	-2	SER	-	expression tag	UNP Q9QZ85
D	-1	THR	-	expression tag	UNP Q9QZ85
D	0	THR	-	expression tag	UNP Q9QZ85
D	31	GLU	ARG	engineered mutation	UNP Q9QZ85
D	32	GLU	LYS	engineered mutation	UNP Q9QZ85
D	176	GLU	LYS	engineered mutation	UNP Q9QZ85
D	246	GLU	LYS	engineered mutation	UNP Q9QZ85
E	-9	GLY	-	expression tag	UNP Q9QZ85


Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-8	SER	-	expression tag	UNP Q9QZ85
E	-7	PRO	-	expression tag	UNP Q9QZ85
E	-6	GLY	-	expression tag	UNP Q9QZ85
E	-5	ILE	-	expression tag	UNP Q9QZ85
E	-4	PRO	-	expression tag	UNP Q9QZ85
E	-3	GLY	-	expression tag	UNP Q9QZ85
E	-2	SER	-	expression tag	UNP Q9QZ85
E	-1	THR	-	expression tag	UNP Q9QZ85
E	0	THR	-	expression tag	UNP Q9QZ85
E	31	GLU	ARG	engineered mutation	UNP Q9QZ85
E	32	GLU	LYS	engineered mutation	UNP Q9QZ85
E	176	GLU	LYS	engineered mutation	UNP Q9QZ85
E	246	GLU	LYS	engineered mutation	UNP Q9QZ85
F	-9	GLY	-	expression tag	UNP Q9QZ85
F	-8	SER	-	expression tag	UNP Q9QZ85
F	-7	PRO	-	expression tag	UNP Q9QZ85
F	-6	GLY	-	expression tag	UNP Q9QZ85
F	-5	ILE	-	expression tag	UNP Q9QZ85
F	-4	PRO	-	expression tag	UNP Q9QZ85
F	-3	GLY	-	expression tag	UNP Q9QZ85
F	-2	SER	-	expression tag	UNP Q9QZ85
F	-1	THR	-	expression tag	UNP Q9QZ85
F	0	THR	-	expression tag	UNP Q9QZ85
F	31	GLU	ARG	engineered mutation	UNP Q9QZ85
F	32	GLU	LYS	engineered mutation	UNP Q9QZ85
F	176	GLU	LYS	engineered mutation	UNP Q9QZ85
F	246	GLU	LYS	engineered mutation	UNP Q9QZ85
G	-9	GLY	-	expression tag	UNP Q9QZ85
G	-8	SER	-	expression tag	UNP Q9QZ85
G	-7	PRO	-	expression tag	UNP Q9QZ85
G	-6	GLY	-	expression tag	UNP Q9QZ85
G	-5	ILE	-	expression tag	UNP Q9QZ85
G	-4	PRO	-	expression tag	UNP Q9QZ85
G	-3	GLY	-	expression tag	UNP Q9QZ85
G	-2	SER	-	expression tag	UNP Q9QZ85
G	-1	THR	-	expression tag	UNP Q9QZ85
G	0	THR	-	expression tag	UNP Q9QZ85
G	31	GLU	ARG	engineered mutation	UNP Q9QZ85
G	32	GLU	LYS	engineered mutation	UNP Q9QZ85
G	176	GLU	LYS	engineered mutation	UNP Q9QZ85
G	246	GLU	LYS	engineered mutation	UNP Q9QZ85

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter

code: GNP) (formula: C₁₀H₁₇N₆O₁₃P₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
2	B	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
2	C	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
2	E	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
2	F	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
2	G	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	2	Total	Mg	0	0
			2	2		
3	C	2	Total	Mg	0	0
			2	2		
3	D	1	Total	Mg	0	0
			1	1		
3	E	2	Total	Mg	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	2	Total	Mg	0	0
			2	2		
3	G	2	Total	Mg	0	0
			2	2		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	H	N	O	0	0
			20	4	12	1	3		
4	E	1	Total	C	H	N	O	0	0
			20	4	12	1	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			1	1		
5	B	2	Total	O	0	0
			2	2		
5	C	1	Total	O	0	0
			1	1		
5	D	1	Total	O	0	0
			1	1		
5	F	2	Total	O	0	0
			2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	1	Total	O	0	0
			1	1		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: INTERFERON-INDUCIBLE GTPASE 1




• Molecule 1: INTERFERON-INDUCIBLE GTPASE 1







• Molecule 1: INTERFERON-INDUCIBLE GTPASE 1




• Molecule 1: INTERFERON-INDUCIBLE GTPASE 1






• Molecule 1: INTERFERON-INDUCIBLE GTPASE 1




• Molecule 1: INTERFERON-INDUCIBLE GTPASE 1





● Molecule 1: INTERFERON-INDUCIBLE GTPASE 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	98.51Å 98.51Å 1289.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	68.98 – 3.20 68.98 – 3.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (68.98-3.20) 100.0 (68.98-3.20)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 3.19Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.297 , 0.317 0.297 , 0.316	Depositor DCC
R_{free} test set	3215 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	75.4	Xtrriage
Anisotropy	0.039	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 29.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.36$, $\langle L^2 \rangle = 0.19$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	22647	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, MG, GNP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3258	0.43	0/4383
1	B	0.24	0/3259	0.45	0/4386
1	C	0.24	0/3204	0.42	0/4306
1	D	0.24	0/3245	0.43	0/4366
1	E	0.23	0/3259	0.41	0/4383
1	F	0.22	0/3247	0.41	0/4367
1	G	0.23	0/3298	0.41	0/4439
All	All	0.23	0/22770	0.42	0/30630

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3204	0	3214	218	0
1	B	3204	0	3213	267	0
1	C	3153	0	3172	208	0
1	D	3193	0	3195	238	0
1	E	3205	0	3204	176	0
1	F	3194	0	3191	176	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	3241	0	3247	158	0
2	A	32	0	13	18	0
2	B	32	0	13	8	0
2	C	32	0	13	11	0
2	E	32	0	13	9	0
2	F	32	0	13	5	0
2	G	32	0	13	8	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	1	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
3	G	2	0	0	0	0
4	C	8	12	12	1	0
4	E	8	12	12	1	0
5	A	1	0	0	1	0
5	B	2	0	0	1	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	F	2	0	0	0	0
5	G	1	0	0	0	0
All	All	22623	24	22538	1372	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 1372 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:ALA:HB1	1:A:127:LEU:HD11	1.23	1.11
1:C:173:MSE:HE2	1:D:141:LEU:HB3	1.26	1.10
1:B:78:THR:HG21	1:B:106:GLU:HG2	1.29	1.08
1:G:186:ASP:HA	1:G:189:ILE:HD11	1.36	1.06
1:C:181:VAL:HG13	1:C:230:LEU:HD21	1.40	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	389/423 (92%)	373 (96%)	16 (4%)	0	100	100
1	B	392/423 (93%)	378 (96%)	12 (3%)	2 (0%)	29	67
1	C	381/423 (90%)	369 (97%)	12 (3%)	0	100	100
1	D	392/423 (93%)	371 (95%)	18 (5%)	3 (1%)	19	58
1	E	390/423 (92%)	377 (97%)	12 (3%)	1 (0%)	41	74
1	F	386/423 (91%)	369 (96%)	17 (4%)	0	100	100
1	G	397/423 (94%)	376 (95%)	19 (5%)	2 (0%)	29	67
All	All	2727/2961 (92%)	2613 (96%)	106 (4%)	8 (0%)	41	74

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	336	ILE
1	D	335	GLU
1	G	222	ILE
1	D	104	VAL
1	D	107	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/374 (96%)	350 (97%)	11 (3%)	41	73
1	B	360/374 (96%)	343 (95%)	17 (5%)	26	62

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	355/374 (95%)	345 (97%)	10 (3%)	43	74
1	D	358/374 (96%)	351 (98%)	7 (2%)	55	80
1	E	359/374 (96%)	350 (98%)	9 (2%)	47	77
1	F	359/374 (96%)	348 (97%)	11 (3%)	40	72
1	G	365/374 (98%)	355 (97%)	10 (3%)	44	75
All	All	2517/2618 (96%)	2442 (97%)	75 (3%)	41	73

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	300	PHE
1	G	204	LYS
1	F	305	ASP
1	G	108	THR
1	B	334	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	ASN
1	C	40	ASN
1	C	87	ASN
1	D	360	GLN
1	F	212	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 21 ligands modelled in this entry, 13 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GNP	A	501	3	28,34,34	2.58	11 (39%)	30,54,54	2.71	11 (36%)
2	GNP	E	501	3	28,34,34	2.58	8 (28%)	30,54,54	2.24	8 (26%)
4	TRS	E	504	-	7,7,7	0.38	0	9,9,9	0.31	0
2	GNP	G	501	3	28,34,34	2.61	10 (35%)	30,54,54	2.21	8 (26%)
2	GNP	B	501	3	28,34,34	2.61	8 (28%)	30,54,54	2.26	9 (30%)
2	GNP	F	501	3	28,34,34	2.61	8 (28%)	30,54,54	2.22	9 (30%)
2	GNP	C	501	3	28,34,34	2.67	10 (35%)	30,54,54	2.36	7 (23%)
4	TRS	C	504	-	7,7,7	0.33	0	9,9,9	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GNP	A	501	3	-	11/17/38/38	0/3/3/3
2	GNP	E	501	3	-	8/17/38/38	0/3/3/3
4	TRS	E	504	-	-	0/9/9/9	-
2	GNP	G	501	3	-	10/17/38/38	0/3/3/3
2	GNP	B	501	3	-	4/17/38/38	0/3/3/3
2	GNP	F	501	3	-	9/17/38/38	0/3/3/3
2	GNP	C	501	3	-	6/17/38/38	0/3/3/3
4	TRS	C	504	-	-	3/9/9/9	-

The worst 5 of 55 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	GNP	C4-N9	-7.45	1.37	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	501	GNP	C4-N9	-7.35	1.37	1.47
2	B	501	GNP	C4-N9	-7.32	1.38	1.47
2	A	501	GNP	C4-N9	-7.26	1.38	1.47
2	F	501	GNP	C4-N9	-7.26	1.38	1.47

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	GNP	C4-C5-N7	6.38	110.92	102.46
2	E	501	GNP	C4-C5-N7	6.36	110.89	102.46
2	F	501	GNP	C4-C5-N7	6.23	110.72	102.46
2	G	501	GNP	C4-C5-N7	6.20	110.68	102.46
2	B	501	GNP	C4-C5-N7	6.20	110.67	102.46

There are no chirality outliers.

5 of 51 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	GNP	PA-O3A-PB-O1B
2	A	501	GNP	PA-O3A-PB-O2B
2	A	501	GNP	C5'-O5'-PA-O1A
2	A	501	GNP	C5'-O5'-PA-O2A
2	A	501	GNP	O4'-C1'-N9-C4


There are no ring outliers.


8 monomers are involved in 59 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	GNP	18	0
2	E	501	GNP	9	0
4	E	504	TRS	1	0
2	G	501	GNP	8	0
2	B	501	GNP	8	0
2	F	501	GNP	5	0
2	C	501	GNP	11	0
4	C	504	TRS	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	385/423 (91%)	0.22	17 (4%) 34 21	39, 68, 108, 144	0
1	B	386/423 (91%)	0.30	15 (3%) 39 25	38, 67, 104, 126	0
1	C	379/423 (89%)	0.28	13 (3%) 45 29	38, 64, 103, 126	0
1	D	386/423 (91%)	0.36	19 (4%) 29 17	38, 67, 112, 137	0
1	E	386/423 (91%)	0.20	14 (3%) 42 27	36, 62, 99, 135	0
1	F	384/423 (90%)	0.31	14 (3%) 42 27	37, 65, 98, 111	0
1	G	389/423 (91%)	0.17	7 (1%) 68 55	33, 59, 95, 147	0
All	All	2695/2961 (91%)	0.26	99 (3%) 41 26	33, 65, 103, 147	0

The worst 5 of 99 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	98	GLY	6.2
1	D	131	GLY	6.0
1	C	297	SER	5.2
1	D	134	ASN	4.9
1	C	194	ASP	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	TRS	E	504	8/8	0.55	0.28	79,109,138,138	0
2	GNP	A	501	32/32	0.69	0.24	60,80,100,116	32
3	MG	A	503	1/1	0.73	0.23	38,38,38,38	0
4	TRS	C	504	8/8	0.81	0.20	81,99,142,142	0
3	MG	B	502	1/1	0.81	0.50	110,110,110,110	0
2	GNP	E	501	32/32	0.87	0.25	34,67,79,88	32
2	GNP	B	501	32/32	0.88	0.23	56,69,88,95	32
2	GNP	C	501	32/32	0.89	0.21	46,63,84,91	32
3	MG	F	503	1/1	0.89	0.20	36,36,36,36	0
3	MG	E	502	1/1	0.90	0.14	44,44,44,44	0
2	GNP	F	501	32/32	0.90	0.18	49,66,77,80	32
2	GNP	G	501	32/32	0.90	0.21	44,59,74,79	32
3	MG	B	503	1/1	0.90	0.23	44,44,44,44	0
3	MG	G	502	1/1	0.95	0.08	43,43,43,43	0
3	MG	F	502	1/1	0.95	0.17	71,71,71,71	0
3	MG	A	502	1/1	0.95	0.50	91,91,91,91	0
3	MG	G	503	1/1	0.96	0.19	43,43,43,43	0
3	MG	E	503	1/1	0.96	0.21	20,20,20,20	0
3	MG	D	501	1/1	0.96	0.28	32,32,32,32	0
3	MG	C	503	1/1	0.97	0.20	30,30,30,30	0
3	MG	C	502	1/1	0.99	0.11	68,68,68,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.