



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2024 – 06:50 pm GMT

PDB ID : 6FPI
Title : Structure of fully reduced Hydrogenase (Hyd-1) variant E28Q
Authors : Carr, S.B.; Armstrong, F.A.; Evans, R.M.
Deposited on : 2018-02-09
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	S	327	 2% 76% 5% 18%
1	T	327	 1% 77% 18%
2	L	582	 95% 5%
2	M	582	 96% 1%

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 28141 atoms, of which 13309 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogenase-1 small chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	S	268	4174	1339	2070	360	384	21	86	6	0
1	T	268	4202	1346	2084	366	385	21	89	8	0

- Molecule 2 is a protein called Hydrogenase-1 large chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	L	581	9182	2943	4541	815	856	27	246	16	0
2	M	581	9202	2951	4560	807	857	27	237	18	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
L	28	GLN	GLU	conflict	UNP P0ACD8
M	28	GLN	GLU	conflict	UNP P0ACD8

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	S	1	Total	Fe S	0	0
			8	4 4		
3	T	1	Total	Fe S	0	0
			8	4 4		

- Molecule 4 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	S	1	Total	Fe S	0	0
			7	3 4		
4	T	1	Total	Fe S	0	0
			7	3 4		

- Molecule 5 is FE4-S3 CLUSTER (three-letter code: SF3) (formula: Fe₄S₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
5	S	1	Total	Fe	S	0	0
			7	4	3		
5	T	1	Total	Fe	S	0	0
			7	4	3		

- Molecule 6 is DODECYL-BETA-D-MALTOSE (three-letter code: LMT) (formula: C₂₄H₄₆O₁₁).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
6	S	1	40	13	26	1	0	
6	T	1	40	13	26	1	0	

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
7	S	3	3	3	0	0
7	T	3	3	3	0	0

- Molecule 8 is NI-FE REDUCED ACTIVE CENTER (three-letter code: EJ2) (formula: C₃HFeN₂NiO).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	Fe	H	N	Ni			O
8	L	1	9	3	1	1	2	1	1	0	0
8	M	1	9	3	1	1	2	1	1	0	0

- Molecule 9 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
9	L	1	1	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	M	1	Total	Mg	0	0
			1	1		

- Molecule 10 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	L	1	Total	O	S	0	0
			5	4	1		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	S	193	Total	O	0	0
			193	193		
11	L	405	Total	O	0	0
			405	405		
11	T	183	Total	O	0	0
			183	183		
11	M	445	Total	O	0	0
			445	445		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hydrogenase-1 small chain




- Molecule 1: Hydrogenase-1 small chain



- Molecule 2: Hydrogenase-1 large chain



- Molecule 2: Hydrogenase-1 large chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.82Å 97.73Å 183.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.51 – 1.50 83.51 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.0 (83.51-1.50) 98.0 (83.51-1.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.145 , 0.167 0.150 , 0.173	Depositor DCC
R_{free} test set	13349 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtrriage
Anisotropy	0.128	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.013 for k,h,-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	28141	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SF3, LMT, SF4, F3S, EJ2, SO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	S	0.84	1/2175 (0.0%)	0.98	11/2953 (0.4%)
1	T	0.83	0/2195	0.94	5/2980 (0.2%)
2	L	0.77	1/4803 (0.0%)	0.84	7/6534 (0.1%)
2	M	0.76	0/4812	0.86	7/6546 (0.1%)
All	All	0.79	2/13985 (0.0%)	0.89	30/19013 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	S	262	SER	CB-OG	6.42	1.50	1.42
2	L	379	VAL	C-O	5.14	1.33	1.23

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	T	185	ARG	NE-CZ-NH1	10.47	125.54	120.30
1	S	193	ARG	NE-CZ-NH2	-9.82	115.39	120.30
1	S	193	ARG	NE-CZ-NH1	9.78	125.19	120.30
1	T	260	ARG	NE-CZ-NH2	-8.01	116.30	120.30
2	L	363	ARG	NE-CZ-NH2	7.81	124.21	120.30
2	L	363	ARG	NE-CZ-NH1	-7.74	116.43	120.30
1	S	234	ARG	NE-CZ-NH1	-7.50	116.55	120.30
1	S	260	ARG	CG-CD-NE	-7.41	96.25	111.80
2	M	363	ARG	NE-CZ-NH2	7.21	123.91	120.30
1	S	193	ARG	CG-CD-NE	-7.21	96.66	111.80
1	S	185[A]	ARG	NE-CZ-NH1	6.89	123.74	120.30
1	S	185[B]	ARG	NE-CZ-NH1	6.89	123.74	120.30
1	S	260	ARG	NE-CZ-NH2	-6.58	117.01	120.30
2	M	26	ARG	NE-CZ-NH1	6.52	123.56	120.30
2	L	26	ARG	NE-CZ-NH1	6.09	123.34	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	M	574	ASP	CB-CG-OD2	-5.99	112.91	118.30
2	L	54	ARG	NE-CZ-NH2	-5.88	117.36	120.30
2	M	32	ARG	NE-CZ-NH1	5.76	123.18	120.30
2	L	130	ASP	CB-CG-OD2	-5.69	113.18	118.30
2	M	169	ARG	NE-CZ-NH2	-5.65	117.48	120.30
1	T	93	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	T	234	ARG	NE-CZ-NH1	5.50	123.05	120.30
2	M	338	ASP	CB-CG-OD1	5.44	123.20	118.30
2	M	363	ARG	NE-CZ-NH1	-5.38	117.61	120.30
2	L	574	ASP	CB-CG-OD2	-5.36	113.48	118.30
1	S	185[A]	ARG	NE-CZ-NH2	-5.23	117.68	120.30
1	S	185[B]	ARG	NE-CZ-NH2	-5.23	117.68	120.30
1	S	168	ARG	NE-CZ-NH2	-5.23	117.69	120.30
1	T	177	ARG	NE-CZ-NH2	-5.08	117.76	120.30
2	L	101	ASP	CB-CG-OD1	5.02	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2104	2070	2062	11	0
1	T	2118	2084	2076	13	0
2	L	4641	4541	4535	17	0
2	M	4642	4560	4559	11	0
3	S	8	0	0	0	0
3	T	8	0	0	0	0
4	S	7	0	0	0	0
4	T	7	0	0	0	0
5	S	7	0	0	0	0
5	T	7	0	0	0	0
6	S	14	26	25	1	0
6	T	14	26	25	0	0
7	S	3	0	0	0	0
7	T	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	L	8	1	0	0	0
8	M	8	1	0	0	0
9	L	1	0	0	0	0
9	M	1	0	0	0	0
10	L	5	0	0	0	0
11	L	405	0	0	3	0
11	M	445	0	0	0	0
11	S	193	0	0	2	0
11	T	183	0	0	3	0
All	All	14832	13309	13282	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:457[A]:HIS:CE1	11:L:701:HOH:O	1.87	1.25
11:S:503:HOH:O	2:L:183:ARG:HD2	1.10	1.23
2:L:457[A]:HIS:ND1	11:L:701:HOH:O	1.77	1.07
2:L:457[A]:HIS:ND1	11:L:702:HOH:O	2.14	0.81
1:T:168:ARG:CZ	11:T:502:HOH:O	2.30	0.79
1:T:93:ARG:CZ	1:T:97:GLU:OE1	2.45	0.64
2:L:135:LEU:HD22	2:L:187:TRP:CD1	2.36	0.60
1:S:13:HIS:ND1	1:S:16[B]:GLU:OE2	2.29	0.59
2:L:73[B]:GLU:HA	2:L:73[B]:GLU:OE1	2.02	0.59
1:S:197:ASP:OD2	1:T:197:ASP:OD2	2.20	0.58
11:S:503:HOH:O	2:L:183:ARG:CD	1.95	0.54
2:M:530:VAL:CG1	2:M:531:PRO:HD2	2.40	0.51
1:T:261:GLY:O	11:T:501:HOH:O	2.19	0.51
1:S:39[B]:LEU:HD21	1:S:161:THR:HG22	1.94	0.49
1:T:186:ILE:HD11	1:T:228:ASN:HB3	1.94	0.49
1:S:234:ARG:CZ	1:T:234:ARG:CZ	2.91	0.48
2:L:530:VAL:CG1	2:L:531:PRO:HD2	2.44	0.48
1:S:186:ILE:HD11	1:S:228:ASN:HB3	1.96	0.47
1:S:7:ILE:HG21	6:S:404:LMT:C6	2.45	0.47
2:L:445:ILE:O	2:L:450:GLY:HA3	2.16	0.46
1:T:168:ARG:NH1	11:T:502:HOH:O	2.47	0.45
2:M:60[B]:LEU:HD11	2:M:72:VAL:CG1	2.47	0.45
2:L:530:VAL:HG11	2:L:579:CYS:HB3	1.98	0.44
2:L:535:ASN:HB3	2:L:548:TYR:CE1	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:540:ASP:HB2	2:M:541:PRO:CD	2.48	0.43
1:S:61:GLU:OE1	1:S:101:ARG:NH1	2.52	0.43
2:L:56:LEU:O	2:L:60:LEU:HD23	2.19	0.43
2:M:261:GLN:HE21	2:M:265[B]:THR:HG23	1.82	0.43
2:M:530:VAL:HG13	2:M:531:PRO:HD2	2.01	0.43
2:M:73[B]:GLU:HA	2:M:73[B]:GLU:OE1	2.19	0.43
1:S:234:ARG:NH1	1:T:234:ARG:NE	2.67	0.43
2:L:536:ALA:HB2	2:L:548:TYR:CE2	2.54	0.42
2:L:4:GLN:HA	2:L:12:ILE:O	2.18	0.42
2:M:488[B]:GLU:HG3	2:M:489:LYS:HG3	2.01	0.42
1:T:93:ARG:NH2	1:T:97:GLU:OE1	2.54	0.41
1:T:260:ARG:HH11	1:T:260:ARG:HG3	1.85	0.41
2:M:535:ASN:HB3	2:M:548:TYR:CE1	2.56	0.41
1:S:234:ARG:NH1	1:T:234:ARG:NH2	2.68	0.41
1:S:234:ARG:NH1	1:T:234:ARG:CZ	2.83	0.41
2:M:272:ASN:OD1	2:M:460:GLN:HG3	2.21	0.41
2:M:73[B]:GLU:OE2	2:M:81:GLY:N	2.54	0.41
2:L:519[B]:ARG:HG2	2:L:520:ASP:OD2	2.20	0.40
1:S:241:PHE:CE2	1:S:243:ILE:HB	2.56	0.40
2:L:540:ASP:HB2	2:L:541:PRO:CD	2.51	0.40
2:L:144:LYS:HB3	2:L:197[A]:GLU:HG2	2.02	0.40
1:T:89:ILE:HB	2:M:51:THR:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	272/327 (83%)	260 (96%)	12 (4%)	0	100	100
1	T	274/327 (84%)	266 (97%)	8 (3%)	0	100	100
2	L	595/582 (102%)	580 (98%)	15 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	M	597/582 (103%)	581 (97%)	16 (3%)	0	100	100
All	All	1738/1818 (96%)	1687 (97%)	51 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	227/266 (85%)	224 (99%)	3 (1%)	69	44
1	T	228/266 (86%)	225 (99%)	3 (1%)	69	44
2	L	496/481 (103%)	495 (100%)	1 (0%)	93	86
2	M	498/481 (104%)	498 (100%)	0	100	100
All	All	1449/1494 (97%)	1442 (100%)	7 (0%)	91	78

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	S	16[A]	GLU
1	S	16[B]	GLU
1	S	191	TYR
2	L	524	ASP
1	T	16	GLU
1	T	191	TYR
1	T	242	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	T	244	GLN
2	M	261	GLN
2	M	467	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 8 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SF4	T	401	1	0,12,12	-	-	-		
6	LMT	T	404	-	13,13,36	0.35	0	12,12,47	0.39	0
6	LMT	S	404	-	13,13,36	0.27	0	12,12,47	0.42	0
8	EJ2	M	601	2	1,6,6	2.92	1 (100%)	-		
4	F3S	S	402	1	0,9,9	-	-	-		
8	EJ2	L	601	2	1,6,6	2.06	1 (100%)	-		
4	F3S	T	402	1	0,9,9	-	-	-		
10	SO4	L	603	-	4,4,4	0.32	0	6,6,6	0.29	0
3	SF4	S	401	1	0,12,12	-	-	-		
5	SF3	T	403	1	0,8,8	-	-	-		
5	SF3	S	403	1	0,8,8	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	T	401	1	-	-	0/6/5/5
6	LMT	T	404	-	-	3/11/11/61	-
6	LMT	S	404	-	-	4/11/11/61	-
4	F3S	S	402	1	-	-	0/3/3/3
4	F3S	T	402	1	-	-	0/3/3/3
3	SF4	S	401	1	-	-	0/6/5/5
5	SF3	T	403	1	-	-	0/2/2/2
5	SF3	S	403	1	-	-	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	M	601	EJ2	O3-C3	2.92	1.20	1.16
8	L	601	EJ2	O3-C3	-2.06	1.13	1.16

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	S	404	LMT	O1'-C1-C2-C3
6	T	404	LMT	C11-C10-C9-C8
6	S	404	LMT	C2-C3-C4-C5
6	S	404	LMT	C11-C10-C9-C8
6	T	404	LMT	C1-C2-C3-C4
6	T	404	LMT	C4-C5-C6-C7
6	S	404	LMT	C1-C2-C3-C4


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	S	404	LMT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	S	268/327 (81%)	-0.49	5 (1%) 66 71	12, 18, 31, 64	0
1	T	268/327 (81%)	-0.48	4 (1%) 73 78	12, 19, 31, 58	0
2	L	581/582 (99%)	-0.58	2 (0%) 94 95	13, 21, 34, 58	0
2	M	581/582 (99%)	-0.59	0 100 100	13, 20, 33, 47	0
All	All	1698/1818 (93%)	-0.55	11 (0%) 89 91	12, 20, 34, 64	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	S	268	VAL	5.0
1	T	268	VAL	4.7
1	S	270	ILE	3.7
1	S	269	ASP	3.6
1	T	271	PRO	3.3
1	S	4	LYS	3.3
1	T	270	ILE	3.2
1	T	269	ASP	3.0
1	S	271	PRO	2.6
2	L	380	LYS	2.2
2	L	378	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	LMT	S	404	14/35	0.68	0.33	0,25,32,34	40
6	LMT	T	404	14/35	0.79	0.18	0,26,32,33	40
10	SO4	L	603	5/5	0.98	0.13	20,21,23,27	5
7	CL	S	406	1/1	0.99	0.05	25,25,25,25	0
7	CL	T	405	1/1	0.99	0.06	21,21,21,21	0
7	CL	T	406	1/1	0.99	0.04	27,27,27,27	0
7	CL	T	407	1/1	0.99	0.06	19,19,19,19	0
8	EJ2	L	601	8/8	0.99	0.08	11,13,14,16	0
7	CL	S	405	1/1	0.99	0.07	19,19,19,19	0
4	F3S	T	402	7/7	1.00	0.07	13,14,14,15	0
7	CL	S	407	1/1	1.00	0.06	18,18,18,18	0
5	SF3	S	403	7/7	1.00	0.07	12,13,14,14	0
5	SF3	T	403	7/7	1.00	0.08	13,14,14,15	0
3	SF4	S	401	8/8	1.00	0.08	13,13,14,14	0
3	SF4	T	401	8/8	1.00	0.08	13,13,14,14	0
8	EJ2	M	601	8/8	1.00	0.08	12,13,14,16	0
9	MG	L	602	1/1	1.00	0.11	12,12,12,12	0
9	MG	M	602	1/1	1.00	0.14	11,11,11,11	0
4	F3S	S	402	7/7	1.00	0.07	13,13,14,14	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LMT S 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around LMT T 404:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.