

Full wwPDB X-ray Structure Validation Report (i)

Jun 17, 2025 – 01:06 pm BST

PDB ID : 9FPX / pdb 00009fpx

Title : Crystal structure of DYRK1A in complex with Curcumin

Authors: Grygier, P.; Pustelny, K.; Czarna, A.

Deposited on : 2024-06-14

Resolution : 2.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1 Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 2.0rc1

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	371	75%	16%	•	9%
1	В	371	72%	19%		9%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5258 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dual specificity tyrosine-phosphorylation-regulated kinase 1A.


Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace		
1	A	337	Total 2660	C 1714	N 451	O 477	P 1	S 17	0	0	0
1	В	337	Total 2436	C 1550	N 426	O 443	P 1	S 16	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	125	MET	-	initiating methionine	UNP Q13627
A	490	HIS	-	expression tag	UNP Q13627
A	491	HIS	-	expression tag	UNP Q13627
A	492	HIS	-	expression tag	UNP Q13627
A	493	HIS	-	expression tag	UNP Q13627
A	494	HIS	-	expression tag	UNP Q13627
A	495	HIS	-	expression tag	UNP Q13627
В	125	MET	-	initiating methionine	UNP Q13627
В	490	HIS	-	expression tag	UNP Q13627
В	491	HIS	-	expression tag	UNP Q13627
В	492	HIS	-	expression tag	UNP Q13627
В	493	HIS	-	expression tag	UNP Q13627
В	494	HIS		expression tag	UNP Q13627
В	495	HIS	-	expression tag	UNP Q13627


• Molecule 2 is curcumin (CCD ID: CC9) (formula: C₂₁H₂₀O₆) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 27 21 6	0	0
2	В	1	Total C O 27 21 6	0	0

 \bullet Molecule 3 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $\mathrm{C_8H_{18}O_5}).$



Mo	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 13 8 5	0	0

• Molecule 4 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	65	Total O 65 65	0	0
4	В	30	Total O 30 30	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 63	Depositor
Cell constants	132.05Å 132.05Å 91.79Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.53 - 2.69	Depositor
resolution (A)	48.53 - 2.69	EDS
% Data completeness	92.1 (48.53-2.69)	Depositor
(in resolution range)	92.1 (48.53-2.69)	EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.97 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.240 , 0.278	Depositor
it, it free	0.243 , 0.279	DCC
R_{free} test set	1257 reflections $(4.89%)$	wwPDB-VP
Wilson B-factor (Å ²)	55.0	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.35 \; , 75.3$	EDS
L-test for twinning ²	$< L > = 0.51, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	0.038 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5258	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, CC9, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.20	0/2705	0.37	1/3662 (0.0%)	
1	В	0.16	0/2476	0.37	0/3366	
All	All	0.18	0/5181	0.37	1/7028 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
1	A	386	ASP	N-CA-C	-5.48	106.67	113.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2660	0	2536	38	0
1	В	2436	0	2088	46	0
2	A	27	0	0	1	0
2	В	27	0	0	2	0
3	A	13	0	18	0	0
4	A	65	0	0	1	0
4	В	30	0	0	0	0
All	All	5258	0	4642	84	0



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 9.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:B:175:LYS:HG3	1:B:240:MET:HE1	1.63	0.80
1:A:388:ALA:HB3	1:A:391:ALA:HB2	1.62	0.79
1:B:438:ARG:HD2	1:B:444:HIS:CE1	2.21	0.75
1:B:380:PRO:HB2	1:B:385:LEU:HD11	1.68	0.75
1:A:351:LEU:HD12	1:A:459:MET:HE1	1.75	0.68
1:B:388:ALA:HB3	1:B:391:ALA:HB2	1.78	0.66
1:A:280:GLU:OE1	1:A:280:GLU:N	2.25	0.64
1:B:307:ASP:HA	2:B:501:CC9:C7	2.27	0.64
1:A:348:GLY:HA2	1:A:459:MET:HE3	1.79	0.62
1:A:164:LEU:HD11	1:A:172:GLN:HB3	1.81	0.62
1:A:461:ASP:O	1:A:467:ARG:NH1	2.32	0.61
1:B:261:LEU:HG	1:B:265:PHE:CE2	2.36	0.60
1:B:190:ILE:HG12	1:B:200:ALA:HB2	1.84	0.60
1:B:242:SER:OG	1:B:243:TYR:N	2.30	0.60
1:A:381:PRO:HB2	1:A:383:HIS:ND1	2.17	0.58
1:B:461:ASP:OD2	1:B:466:THR:OG1	2.21	0.58
1:A:147:TYR:HB2	1:A:233:HIS:CD2	2.41	0.56
1:A:209:LEU:HG	1:A:281:LEU:HD23	1.86	0.56
1:B:445:THR:HG23	1:B:447:ALA:H	1.69	0.56
1:B:287:ASP:OD1	1:B:289:LYS:HD2	2.08	0.54
1:B:194:LYS:O	1:B:197:LEU:N	2.41	0.53
1:B:350:ILE:O	1:B:354:MET:HG2	2.08	0.53
1:B:283:ILE:HA	1:B:313:GLN:HA	1.89	0.53
1:B:322:ILE:HD12	1:B:323:GLN:HB2	1.91	0.53
1:A:165:ILE:HD13	1:A:240:MET:HE1	1.91	0.52
1:B:290:PRO:HG3	1:B:350:ILE:HG23	1.92	0.51
1:B:438:ARG:HD2	1:B:444:HIS:HE1	1.71	0.51
1:A:199:GLN:HA	1:A:202:ILE:HD12	1.91	0.51
1:A:307:ASP:HA	2:A:501:CC9:C19	2.41	0.51
1:A:429:VAL:HG12	1:A:446:VAL:HG13	1.92	0.51
1:B:259:LEU:O	1:B:263:ARG:HG3	2.12	0.50
1:B:352:VAL:O	1:B:356:THR:HG23	2.12	0.49
1:B:248:LEU:O	1:B:252:THR:HG23	2.12	0.49
1:A:217:MET:HB3	1:A:275:PHE:HB2	1.95	0.49
1:A:222:VAL:HB	1:A:306:VAL:HG12	1.93	0.49
1:A:336:MET:HE2	1:A:388:ALA:HA	1.96	0.48
1:B:438:ARG:HB3	1:B:444:HIS:HD1	1.78	0.48

Continued on next page...



Continued from previous page...

Continuea from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:149:VAL:HG21	1:A:164:LEU:HD13	1.95	0.48
1:B:263:ARG:HG2	1:B:478:PHE:CZ	2.49	0.48
1:B:445:THR:HG23	1:B:447:ALA:N	2.29	0.48
1:A:210:MET:HE1	1:A:308:PHE:CE2	2.49	0.47
1:B:260:ASN:O	1:B:263:ARG:HB2	2.16	0.46
1:A:377:LEU:HD11	1:A:460:LEU:HB3	1.98	0.46
1:B:339:ASP:N	1:B:339:ASP:OD1	2.48	0.46
1:A:217:MET:SD	1:A:274:LEU:HD12	2.56	0.45
1:B:210:MET:C	1:B:212:LYS:H	2.24	0.45
1:A:331:GLU:CD	1:A:467:ARG:HH22	2.23	0.45
1:A:458:ARG:HA	1:A:458:ARG:HD2	1.83	0.45
1:A:146:ASP:OD1	1:A:191:LYS:HG2	2.17	0.45
1:B:284:ILE:N	1:B:312:CYS:O	2.43	0.45
1:B:346:SER:O	1:B:350:ILE:HG13	2.16	0.45
1:A:334:LEU:HD23	1:A:394:PHE:CE2	2.52	0.44
1:A:280:GLU:H	1:A:280:GLU:CD	2.20	0.44
1:B:347:LEU:O	1:B:351:LEU:HG	2.17	0.44
1:B:292:ASN:ND2	1:B:307:ASP:HB3	2.32	0.44
1:B:190:ILE:HG12	1:B:200:ALA:CB	2.48	0.44
1:A:278:THR:HB	1:A:281:LEU:HD12	2.00	0.44
1:B:203:GLU:HG3	1:B:308:PHE:O	2.17	0.43
1:A:382:ALA:HA	1:A:385:LEU:HB2	2.01	0.43
1:B:263:ARG:NH1	1:B:479:PHE:HA	2.33	0.43
1:B:354:MET:HA	1:B:354:MET:HE3	2.01	0.43
1:B:341:ALA:HB1	1:B:467:ARG:HD2	1.98	0.43
1:A:334:LEU:HB2	1:A:336:MET:HE3	1.99	0.43
1:A:385:LEU:C	1:A:387:GLN:H	2.26	0.43
1:B:210:MET:SD	1:B:221:ILE:HG21	2.59	0.43
1:B:334:LEU:HB3	1:B:388:ALA:HB1	2.00	0.43
1:B:458:ARG:HB3	1:B:468:ILE:HB	2.00	0.43
1:B:241:LEU:N	2:B:501:CC9:O4	2.48	0.42
1:B:288:LEU:HB3	1:B:350:ILE:HD11	2.01	0.42
1:A:164:LEU:HD12	1:A:174:VAL:HG12	2.01	0.42
1:A:320:GLN:O	1:A:321:PTR:C	2.68	0.42
1:B:424:HIS:C	1:B:424:HIS:CD2	2.97	0.42
1:B:170:PHE:CD1	1:B:188:LYS:HE3	2.55	0.41
1:A:330:PRO:O	1:A:334:LEU:HG	2.21	0.41
1:B:366:GLU:OE1	1:B:390:LYS:NZ	2.53	0.41
1:A:322:ILE:HD11	1:A:332:VAL:CG2	2.51	0.41
1:A:174:VAL:HG22	1:A:187:ILE:HB	2.03	0.41
1:A:264:LYS:NZ	4:A:606:HOH:O	2.45	0.41

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:B:249:LEU:HD21	1:B:354:MET:HE3	2.03	0.41
1:A:379:ILE:HA	1:A:380:PRO:HD3	1.95	0.41
1:B:330:PRO:HD3	1:B:345:TRP:CZ2	2.56	0.41
1:A:155:TRP:HB2	1:A:159:TYR:HB2	2.02	0.40
1:A:379:ILE:O	1:A:379:ILE:HG13	2.17	0.40
1:B:275:PHE:O	1:B:278:THR:HG23	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	in Analysed Favoured Allowed		Outliers	Perce	entiles	
1	A	330/371~(89%)	311 (94%)	19 (6%)	0	100	100
1	В	$332/371\ (90\%)$	304 (92%)	26 (8%)	2 (1%)	22	45
All	All	$662/742 \ (89\%)$	615 (93%)	45 (7%)	2 (0%)	37	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	194	LYS
1	В	242	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	267/329 (81%)	263 (98%)	4 (2%)	60	83	
1	В	202/329 (61%)	200 (99%)	2 (1%)	73	89	
All	All	469/658 (71%)	463 (99%)	6 (1%)	65	85	

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	173	VAL
1	A	284	ILE
1	A	305	ILE
1	A	379	ILE
1	В	157	ASP
1	В	400	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	GLN
1	A	387	GLN
1	В	223	HIS
1	В	404	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Γ.	Mol	Tuno	Chain	Dec	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
-	MOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	1	PTR	A	321	1	15,16,17	1.23	1 (6%)	19,22,24	0.63	0



Mol	Type	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
IVIOI	туре	Chain	rtes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	В	321	1	15,16,17	1.24	1 (6%)	19,22,24	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

	Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
	1	PTR	A	321	1	-	0/10/11/13	0/1/1/1
ſ	1	PTR	В	321	1	-	0/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
1	В	321	PTR	OH-CZ	-4.34	1.30	1.40
1	A	321	PTR	OH-CZ	-4.27	1.31	1.40

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	321	PTR	1	0

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Ty	Type	Chain	Res	Link	Во	nd leng	ths	В	ond ang	les
	туре	Chain	rtes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	PG4	A	502	-	12,12,12	0.12	0	11,11,11	0.65	0
2	CC9	A	501	-	26,28,28	0.62	1 (3%)	35,37,37	0.43	0
2	CC9	В	501	-	26,28,28	0.47	0	35,37,37	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	A	502	-	-	3/10/10/10	-
2	CC9	A	501	-	-	3/18/18/18	0/2/2/2
2	CC9	В	501	-	-	6/18/18/18	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}({ ext{ iny A}})$
2	A	501	CC9	C13-C12	-2.24	1.40	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	CC9	C1-C2-C3-C4
2	В	501	CC9	C1-C2-C3-C4
2	В	501	CC9	C7-C8-O3-C9
2	A	501	CC9	C13-C14-C15-C16
2	A	501	CC9	C13-C14-C15-C21
2	В	501	CC9	O4-C12-C13-C14
3	A	502	PG4	C8-C7-O4-C6
3	A	502	PG4	C1-C2-O2-C3
2	В	501	CC9	C10-C8-O3-C9
3	A	502	PG4	C5-C6-O4-C7
2	В	501	CC9	C11-C12-C13-C14
2	В	501	CC9	C2-C1-C11-C12




There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	CC9	1	0
2	В	501	CC9	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	336/371 (90%)	0.23	14 (4%) 41 39	33, 57, 97, 124	0
1	В	336/371 (90%)	0.68	18 (5%) 32 30	54, 86, 122, 144	0
All	All	672/742 (90%)	0.45	32 (4%) 36 35	33, 72, 116, 144	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	320	GLN	4.2
1	В	270	CYS	3.6
1	В	266	ALA	3.6
1	В	478	PHE	3.4
1	A	139	GLY	3.3
1	A	371	ASN	3.0
1	В	237	VAL	2.8
1	A	395	PHE	2.7
1	В	452	PHE	2.7
1	В	480	LYS	2.7
1	В	243	TYR	2.6
1	В	215	THR	2.6
1	В	217	MET	2.6
1	A	138	ASP	2.5
1	В	211	ASN	2.5
1	A	322	ILE	2.5
1	В	411	GLY	2.5
1	В	477	SER	2.5
1	A	409	LYS	2.4
1	В	245	LEU	2.4
1	В	155	TRP	2.4
1	A	394	PHE	2.4
1	A	408	THR	2.3
1	A	170	PHE	2.3

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	403	TRP	2.3
1	A	366	GLU	2.2
1	A	253	ASN	2.2
1	A	396	GLU	2.2
1	В	379	ILE	2.1
1	A	243	TYR	2.1
1	В	214	ASP	2.1
1	В	196	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	PTR	A	321	16/17	0.87	0.12	64,71,79,85	0
1	PTR	В	321	16/17	0.89	0.10	80,89,94,96	0

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	CC9	В	501	27/27	0.80	0.16	66,82,91,96	0
3	PG4	A	502	13/13	0.80	0.15	59,66,76,80	0
2	CC9	A	501	27/27	0.85	0.13	37,51,82,85	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CC9 B 501:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around CC9 A 501:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

