



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 3, 2024 – 09:58 pm GMT

PDB ID : 5FQ7  
Title : Crystal structure of the SusCD complex BT2261-2264 from *Bacteroides thetaiotaomicron*  
Authors : Glenwright, A.J.; Pothula, K.R.; Chorev, D.S.; Basle, A.; Robinson, C.V.; Kleinekathoefer, U.; Bolam, D.N.; van den Berg, B.  
Deposited on : 2015-12-07  
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	 70% 29% .
1	C	480	 71% 28% .
2	B	984	 66% 29% . .
2	D	984	 68% 26% . .
3	E	148	 84% 14% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	148	<p>8% 76% 22%</p>
4	G	212	<p>8% 29% 11% 60%</p>
4	H	212	<p>4% 28% 13% 58%</p>
5	I	10	<p>20% 90% 10%</p>
5	P	10	<p>20% 80% 20%</p>

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 25886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BT\_2263.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	480	3744	2370	616	741	17	0	0	0
1	C	480	3744	2370	616	741	17	0	0	0

- Molecule 2 is a protein called BT\_2264.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	943	7342	4643	1224	1445	30	0	0	0
2	D	943	7337	4642	1224	1441	30	0	0	0

- Molecule 3 is a protein called BT\_2261.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	145	1134	717	178	234	5	0	0	0
3	F	145	1134	717	178	234	5	0	0	0

- Molecule 4 is a protein called BT\_2262.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	G	84	665	422	102	140	1	0	0	0
4	H	89	698	441	108	148	1	0	0	0

- Molecule 5 is a protein called PEPTIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	I	10	Total	C	N	O	0	0	0
			40	20	10	10			
5	P	10	Total	C	N	O	0	0	0
			40	20	10	10			


- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		


- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	3	Total	Na	0	0
			3	3		
7	C	1	Total	Na	0	0
			1	1		
7	D	2	Total	Na	0	0
			2	2		






• Molecule 2: BT\_2264








• Molecule 3: BT\_2261



• Molecule 3: BT\_2261




• Molecule 4: BT\_2262





VAL  
GLY  
LYS  
PHE  
GLU  
LEU  
LEU  
ASN  
ASP  
ASP  
ASN  
THR  
THR  
ILE  
THR  
PRO  
LEU  
GLU  
LEU  
TYR  
VAL  
VAL  
ALA  
GLY  
TRP  
GLY  
GLY  
ASP  
SER  
SER  
MET  
ASP  
GLN  
MET  
THR  
THR  
ASN  
THR  
LEU  
LEU  
ASP  
PRO  
ALA  
THR  
GLY  
LEU  
LYS  
LEU  
TRP  
THR  
VAL  
ALA  
TYR  
GLY  
GLN  
LEU  
SER

● Molecule 4: BT\_2262





THR  
ALA  
SER  
PRO  
LEU  
LYS  
SER  
GLY  
ILE  
TYR  
ASP  
THR  
VAL  
ALA  
GLU  
GLY  
SER  
LEU  
ARG  
THR  
ALA  
PRO  
SER  
VAL  
VAL  
VAL  
VAL  
ALA  
PHE  
SER  
SER  
MET  
TYR  
ASP  
GLU  
ILE  
VAL  
ILE  
PHE  
GLN  
V59  
N60  
MET  
GLU  
PRO  
GLY  
ILE  
PHE  
TYR  
ILE  
SER  
ASP  
PHE  
LEU  
GLY  
ALA  
TYR  
TRP  
TYR  
ALA  
GLY  
GLN  
ARG  
LEU  
SER  
ALA  
TYR  
GLY  
PRO  
ASP  
VAL



TYR  
ALA  
MET  
VAL  
GLY  
PHE  
GLU  
LEU  
ASN  
ASP  
ASN  
THR  
ILE  
THR  
PRO  
LEU  
GLU  
SER  
GLY  
SER  
ASP  
MET  
ASP  
GLN  
MET  
THR  
ASN  
THR  
LEU  
LEU  
ASP  
PRO  
ALA  
THR  
GLY  
TYR  
LEU  
TRP  
VAL  
ALA  
TYR  
ALA  
GLY  
GLN  
SER  
PHE  
ILE  
ASP  
ILE  
VAL

LYS  
GLN

● Molecule 5: PEPTIDE



● Molecule 5: PEPTIDE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.59Å 180.16Å 245.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.43 – 3.40 49.45 – 3.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.43-3.40) 94.7 (49.45-3.40)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 3.40Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.201 , 0.255 0.201 , 0.256	Depositor DCC
$R_{free}$ test set	7451 reflections (9.79%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.4	Xtrriage
Anisotropy	0.394	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	25886	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	1/3835 (0.0%)	0.66	0/5220
1	C	0.68	2/3835 (0.1%)	0.65	0/5220
2	B	0.60	0/7518	0.70	3/10196 (0.0%)
2	D	0.62	1/7513 (0.0%)	0.71	4/10190 (0.0%)
3	E	0.49	0/1162	0.57	0/1578
3	F	0.43	0/1162	0.57	0/1578
4	G	0.36	0/677	0.55	0/918
4	H	0.55	0/711	0.65	0/966
5	I	0.59	0/39	1.00	0/47
5	P	0.55	0/39	1.03	0/47
All	All	0.61	4/26491 (0.0%)	0.67	7/35960 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2
2	D	0	1
All	All	0	3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	225	CYS	CB-SG	-6.56	1.71	1.82
1	A	225	CYS	CB-SG	-6.40	1.71	1.82
2	D	942	THR	C-N	-6.02	1.22	1.34
1	C	166	VAL	CB-CG1	-5.11	1.42	1.52

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	921	LEU	CB-CG-CD1	-7.91	97.56	111.00
2	D	624	GLY	N-CA-C	-7.29	94.87	113.10
2	B	624	GLY	N-CA-C	-6.89	95.88	113.10
2	D	623	LEU	CA-CB-CG	6.38	129.97	115.30
2	B	623	LEU	CA-CB-CG	5.51	127.98	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	623	LEU	Peptide
2	B	949	ILE	Peptide
2	D	623	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3744	0	3562	96	0
1	C	3744	0	3562	95	0
2	B	7342	0	7013	208	0
2	D	7337	0	7014	213	0
3	E	1134	0	1050	12	0
3	F	1134	0	1050	21	0
4	G	665	0	626	16	0
4	H	698	0	656	21	0
5	I	40	0	32	1	0
5	P	40	0	32	1	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	B	3	0	0	0	0
7	C	1	0	0	0	0
7	D	2	0	0	0	0
All	All	25886	0	24597	653	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 653 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:198:PHE:CD1	2:D:851:ARG:NH2	2.23	1.06
2:D:124:TYR:OH	2:D:415:LYS:NZ	1.93	1.02
2:B:172:GLU:HG2	2:B:173:LYS:HG2	1.43	0.98
2:D:198:PHE:HD1	2:D:851:ARG:NH2	1.59	0.97
2:D:579:GLN:HB2	2:D:581:VAL:HG12	1.46	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	478/480 (100%)	425 (89%)	45 (9%)	8 (2%)	9	34
1	C	478/480 (100%)	432 (90%)	40 (8%)	6 (1%)	12	39
2	B	939/984 (95%)	828 (88%)	99 (10%)	12 (1%)	12	39
2	D	939/984 (95%)	848 (90%)	80 (8%)	11 (1%)	13	41
3	E	143/148 (97%)	130 (91%)	11 (8%)	2 (1%)	11	37
3	F	143/148 (97%)	124 (87%)	19 (13%)	0	100	100
4	G	80/212 (38%)	77 (96%)	3 (4%)	0	100	100
4	H	87/212 (41%)	82 (94%)	5 (6%)	0	100	100
5	I	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
5	P	8/10 (80%)	5 (62%)	2 (25%)	1 (12%)	0	2
All	All	3303/3668 (90%)	2958 (90%)	305 (9%)	40 (1%)	13	41

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	411	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	714	LYS
2	B	943	PRO
1	C	57	GLN
1	C	120	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	392/392 (100%)	386 (98%)	6 (2%)	65 82
1	C	392/392 (100%)	387 (99%)	5 (1%)	69 84
2	B	796/836 (95%)	787 (99%)	9 (1%)	73 86
2	D	795/836 (95%)	784 (99%)	11 (1%)	67 83
3	E	121/124 (98%)	121 (100%)	0	100 100
3	F	121/124 (98%)	120 (99%)	1 (1%)	81 91
4	G	73/177 (41%)	71 (97%)	2 (3%)	44 70
4	H	76/177 (43%)	73 (96%)	3 (4%)	32 61
All	All	2766/3058 (90%)	2729 (99%)	37 (1%)	69 84

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	915	ASP
4	H	23	GLU
2	D	920	TRP
4	G	54	LYS
2	B	772	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	800	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	803	GLN
3	E	123	GLN
2	B	490	GLN
1	A	363	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	480/480 (100%)	-0.25	1 (0%) 95 95	52, 70, 91, 112	0
1	C	480/480 (100%)	-0.21	3 (0%) 89 89	56, 72, 93, 120	0
2	B	943/984 (95%)	-0.07	17 (1%) 68 67	55, 75, 109, 147	0
2	D	943/984 (95%)	0.01	13 (1%) 75 74	55, 74, 106, 146	0
3	E	145/148 (97%)	0.24	7 (4%) 30 31	76, 87, 103, 126	0
3	F	145/148 (97%)	0.34	12 (8%) 11 13	80, 99, 126, 151	0
4	G	84/212 (39%)	1.13	18 (21%) 0 1	84, 126, 152, 174	0
4	H	89/212 (41%)	0.70	8 (8%) 9 11	66, 93, 123, 179	0
5	I	10/10 (100%)	1.70	2 (20%) 1 1	53, 62, 75, 78	0
5	P	10/10 (100%)	1.40	2 (20%) 1 1	63, 68, 82, 93	0
All	All	3329/3668 (90%)	-0.00	83 (2%) 57 55	52, 75, 111, 179	0

The worst 5 of 83 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	25	ALA	4.4
4	G	82	THR	4.3
4	H	88	ALA	3.9
4	G	84	THR	3.6
2	B	532	GLY	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	NA	B	1986	1/1	0.84	0.40	27,27,27,27	0
7	NA	C	1481	1/1	0.91	0.16	44,44,44,44	0
7	NA	B	1988	1/1	0.94	0.23	43,43,43,43	0
6	MG	B	1985	1/1	0.96	0.25	23,23,23,23	0
6	MG	D	1985	1/1	0.97	0.21	38,38,38,38	0
7	NA	B	1987	1/1	0.97	0.13	30,30,30,30	0
7	NA	D	1987	1/1	0.97	0.15	34,34,34,34	0
7	NA	D	1986	1/1	0.98	0.21	35,35,35,35	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.