



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 4, 2024 – 11:30 pm GMT

PDB ID : 5FS9
Title : Crystal structure of the G338E mutant of human apoptosis inducing factor
Authors : Sevrioukova, I.
Deposited on : 2015-12-31
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	515	 7% 81% 7% • 11%
1	B	515	 7% 84% 6% 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called APOPTOSIS-INDUCING FACTOR 1, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	459	3597	2284	635	665	13	0	10	0
1	B	465	3640	2308	643	676	13	0	9	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	614	LEU	-	expression tag	UNP O95831
A	615	VAL	-	expression tag	UNP O95831
A	616	PRO	-	expression tag	UNP O95831
A	617	ARG	-	expression tag	UNP O95831
A	338	GLU	GLY	engineered mutation	UNP O95831
B	614	LEU	-	expression tag	UNP O95831
B	615	VAL	-	expression tag	UNP O95831
B	616	PRO	-	expression tag	UNP O95831
B	617	ARG	-	expression tag	UNP O95831
B	338	GLU	GLY	engineered mutation	UNP O95831

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	274	Total 274	O 274	0	0
4	B	267	Total 267	O 267	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: APOPTOSIS-INDUCING FACTOR 1, MITOCHONDRIAL

Chain A: 



- Molecule 1: APOPTOSIS-INDUCING FACTOR 1, MITOCHONDRIAL

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.62Å 114.93Å 121.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	114.93 – 1.75 53.80 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.7 (114.93-1.75) 99.7 (53.80-1.75)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.175 , 0.200 0.183 , 0.208	Depositor DCC
R_{free} test set	6431 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtrriage
Anisotropy	0.038	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7896	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3691	0.69	1/4988 (0.0%)
1	B	0.54	0/3735	0.71	3/5046 (0.1%)
All	All	0.54	0/7426	0.70	4/10034 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	489[A]	ASP	CB-CG-OD1	6.22	123.90	118.30
1	B	489[B]	ASP	CB-CG-OD1	6.22	123.90	118.30
1	B	438	ASP	CB-CG-OD2	-5.69	113.18	118.30
1	A	136	LEU	CB-CG-CD1	5.54	120.41	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	529	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3597	0	3659	23	0
1	B	3640	0	3694	16	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	274	0	0	3	2
4	B	267	0	0	2	2
All	All	7896	0	7431	39	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:454:HIS:ND1	4:A:2211:HOH:O	2.10	0.85
1:B:241:ASN:ND2	1:B:432[A]:ASN:HD21	1.92	0.68
1:A:524:SER:HB3	1:A:526:THR:HG22	1.77	0.67
1:A:153:ARG:HD3	1:A:467:GLU:HG2	1.90	0.54
1:A:131:HIS:HE1	1:A:254:GLU:OE2	1.92	0.52
1:A:527:GLY:HA3	4:A:2041:HOH:O	2.09	0.52
1:A:305:ILE:HD11	1:A:319:LEU:HD12	1.92	0.52
1:A:237:ASP:HB3	1:A:242[B]:MET:HE2	1.92	0.52
1:A:128:ALA:HB2	1:A:228:LEU:HD11	1.93	0.51
1:A:188[A]:THR:HG23	1:A:291:ARG:HG2	1.92	0.51
1:B:131:HIS:HE1	1:B:254:GLU:OE2	1.97	0.48
1:B:347:TYR:HH	1:B:558:GLU:N	2.12	0.47
1:A:182:SER:CB	1:A:187[B]:VAL:HG21	2.44	0.47
1:A:489:ASP:HB2	4:A:2143:HOH:O	2.14	0.47
1:B:228:LEU:HG	1:B:231:LYS:HG3	1.95	0.47
1:B:340:MET:CE	1:B:352:THR:HG21	2.46	0.46
1:A:310:PHE:CE1	1:A:340:MET:HE3	2.51	0.45
1:B:153:ARG:HD3	1:B:467:GLU:HG2	1.97	0.45
1:A:370:GLN:CG	1:A:384:LYS:HA	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:ILE:HG22	1:B:221:GLU:O	2.17	0.44
1:B:131:HIS:HD2	1:B:252:THR:OG1	2.00	0.44
1:B:264:PRO:HD3	1:B:285:ARG:HD3	2.00	0.43
1:B:527:GLY:HA3	4:B:2034:HOH:O	2.18	0.43
1:B:529:ARG:O	1:B:530:SER:HB2	2.18	0.43
1:B:584:ARG:HD2	1:B:608:PHE:HA	1.99	0.43
1:A:370:GLN:HG3	1:A:384:LYS:HA	2.00	0.43
1:A:343:ILE:HG23	1:A:498:VAL:HG12	2.00	0.43
1:B:485:ASP:OD1	1:B:529:ARG:NH1	2.50	0.43
1:A:242[B]:MET:HE2	1:A:242[B]:MET:HB3	1.90	0.42
1:A:340:MET:CE	1:A:352:THR:HG21	2.49	0.42
1:B:405:GLU:HG3	4:B:2164:HOH:O	2.18	0.42
1:A:581:ILE:HD11	1:A:610:ILE:HD11	2.01	0.42
1:A:402:PRO:HG3	1:A:451:ARG:CZ	2.49	0.42
1:A:131:HIS:HD2	1:A:252:THR:OG1	2.02	0.42
1:B:172:ARG:N	1:B:173:PRO:CD	2.83	0.42
1:A:283:LEU:HD23	1:A:396:ALA:HB3	2.01	0.41
1:B:310:PHE:CE1	1:B:340:MET:HE3	2.55	0.41
1:A:182:SER:OG	1:A:187[B]:VAL:HG21	2.19	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2258:HOH:O	4:B:2246:HOH:O[2_454]	1.55	0.65
4:A:2255:HOH:O	4:B:2250:HOH:O[2_454]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	465/515 (90%)	452 (97%)	13 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	470/515 (91%)	456 (97%)	14 (3%)	0	100	100
All	All	935/1030 (91%)	908 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	387/424 (91%)	379 (98%)	8 (2%)	53	31
1	B	392/424 (92%)	387 (99%)	5 (1%)	69	54
All	All	779/848 (92%)	766 (98%)	13 (2%)	60	42

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	136	LEU
1	A	165	ASP
1	A	182	SER
1	A	213	SER
1	A	382	LYS
1	A	524	SER
1	A	529	ARG
1	A	530	SER
1	B	136	LEU
1	B	183	ASP
1	B	324	ARG
1	B	530	SER
1	B	615	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	131	HIS
1	A	250	GLN
1	B	131	HIS
1	B	241	ASN
1	B	250	GLN
1	B	366	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	1000	-	53,58,58	1.30	4 (7%)	68,89,89	1.38	6 (8%)
3	GOL	A	1002	-	5,5,5	0.42	0	5,5,5	0.37	0
2	FAD	B	1000	-	53,58,58	1.30	5 (9%)	68,89,89	1.35	10 (14%)
3	GOL	B	1001	-	5,5,5	0.22	0	5,5,5	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	1000	-	-	3/30/50/50	0/6/6/6
3	GOL	A	1002	-	-	2/4/4/4	-
2	FAD	B	1000	-	-	2/30/50/50	0/6/6/6
3	GOL	B	1001	-	-	0/4/4/4	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1000	FAD	C9A-C5X	4.62	1.48	1.41
2	A	1000	FAD	C9A-C5X	4.51	1.48	1.41
2	A	1000	FAD	C4X-N5	3.44	1.37	1.30
2	B	1000	FAD	C4X-N5	3.16	1.36	1.30
2	B	1000	FAD	C5X-N5	-2.99	1.33	1.39
2	A	1000	FAD	C8-C7	2.41	1.46	1.40
2	B	1000	FAD	C8-C7	2.21	1.46	1.40
2	A	1000	FAD	C5'-C4'	2.16	1.54	1.51
2	B	1000	FAD	C1'-C2'	2.16	1.55	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	FAD	N3A-C2A-N1A	-4.84	121.11	128.68
2	A	1000	FAD	O4B-C1B-C2B	-4.08	100.97	106.93
2	B	1000	FAD	N3A-C2A-N1A	-3.70	122.89	128.68
2	A	1000	FAD	C2A-N1A-C6A	3.21	124.25	118.75
2	B	1000	FAD	O2'-C2'-C1'	3.17	117.47	109.80
2	A	1000	FAD	O2A-PA-O1A	2.80	126.10	112.24
2	B	1000	FAD	O2A-PA-O1A	2.79	126.02	112.24
2	A	1000	FAD	O4-C4-C4X	-2.71	119.41	126.60
2	B	1000	FAD	O4B-C1B-C2B	-2.65	103.05	106.93
2	B	1000	FAD	N6A-C6A-N1A	2.61	124.00	118.57
2	B	1000	FAD	C1B-N9A-C4A	-2.61	122.05	126.64
2	B	1000	FAD	C4-C4X-N5	2.49	121.78	118.23
2	B	1000	FAD	O4-C4-C4X	-2.34	120.39	126.60
2	A	1000	FAD	O2'-C2'-C1'	2.29	115.34	109.80
2	B	1000	FAD	N3-C2-N1	2.07	123.44	119.38
2	B	1000	FAD	C2A-N1A-C6A	2.02	122.21	118.75

There are no chirality outliers.


All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1002	GOL	C1-C2-C3-O3
3	A	1002	GOL	O2-C2-C3-O3
2	A	1000	FAD	PA-O3P-P-O5'
2	B	1000	FAD	PA-O3P-P-O5'
2	B	1000	FAD	O4B-C4B-C5B-O5B
2	A	1000	FAD	O4B-C4B-C5B-O5B
2	A	1000	FAD	C3B-C4B-C5B-O5B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	459/515 (89%)	0.62	37 (8%)	12 16	20, 36, 68, 103	0
1	B	465/515 (90%)	0.39	36 (7%)	13 18	21, 37, 65, 103	0
All	All	924/1030 (89%)	0.51	73 (7%)	12 16	20, 37, 66, 103	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	524	SER	6.6
1	A	528	ILE	6.6
1	A	560	TYR	6.3
1	B	617	ARG	6.3
1	A	612	GLU	5.9
1	A	559	ASP	5.8
1	B	525	GLY	5.6
1	B	524	SER	5.5
1	B	183	ASP	5.3
1	B	127	LYS	5.3
1	A	599	GLU	5.1
1	A	183	ASP	4.7
1	B	526	THR	4.5
1	B	521	THR	4.5
1	A	521	THR	4.3
1	B	126	ASP	4.2
1	A	126	ASP	4.2
1	A	275	ALA	4.0
1	A	525	GLY	3.9
1	B	527	GLY	3.9
1	B	560	TYR	3.8
1	B	528	ILE	3.6
1	A	272	ARG	3.5
1	B	477	TRP	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	186	ASN	3.3
1	A	370	GLN	3.3
1	B	558	GLU	3.3
1	A	274	GLY	3.2
1	A	271	ASP	3.2
1	B	615	VAL	3.2
1	A	128	ALA	3.2
1	A	570	ASP	3.1
1	A	527	GLY	3.0
1	A	459	VAL	3.0
1	A	464	LEU	3.0
1	A	219	HIS	3.0
1	B	570	ASP	2.9
1	B	501[A]	SER	2.9
1	A	368	ILE	2.9
1	B	514	GLN	2.8
1	B	275	ALA	2.8
1	B	559	ASP	2.8
1	B	616	PRO	2.8
1	A	526	THR	2.8
1	B	221	GLU	2.8
1	A	127	LYS	2.8
1	B	199	LYS	2.8
1	B	274	GLY	2.7
1	A	523	GLN	2.7
1	A	247	ASP	2.7
1	B	248	GLY	2.7
1	A	273	ALA	2.6
1	B	614	LEU	2.6
1	A	460	VAL	2.5
1	B	529	ARG	2.5
1	A	408	LYS	2.5
1	A	347	TYR	2.4
1	B	520	ALA	2.4
1	A	276	GLU	2.3
1	B	128	ALA	2.3
1	B	215	GLN	2.2
1	B	500	SER	2.2
1	A	477	TRP	2.2
1	B	276	GLU	2.2
1	A	248	GLY	2.2
1	A	510	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	530	SER	2.1
1	B	189	LYS	2.1
1	B	569	ARG	2.1
1	A	278	LYS	2.1
1	B	571	LYS	2.1
1	A	529	ARG	2.1
1	B	523	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	B	1001	6/6	0.84	0.19	65,69,74,79	0
3	GOL	A	1002	6/6	0.86	0.19	50,63,64,70	0
2	FAD	A	1000	53/53	0.97	0.10	21,25,29,36	0
2	FAD	B	1000	53/53	0.98	0.08	21,25,31,35	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.