



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2024 – 09:36 PM EDT

PDB ID : 4FXW  
Title : Structure of phosphorylated SF1 complex with U2AF65-UHM domain  
Authors : Wang, W.; Bauer, W.J.; Wedekind, J.E.; Kielkopf, C.L.  
Deposited on : 2012-07-03  
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	106	
1	C	106	
2	B	124	
2	D	124	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	C	501	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3409 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Splicing factor U2AF 65 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	105	830	523	143	156	5	3	0	
1	C	104	823	520	135	160	5	0	0	

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	370	GLY	-	EXPRESSION TAG	UNP P26368
A	371	PRO	-	EXPRESSION TAG	UNP P26368
A	372	LEU	-	EXPRESSION TAG	UNP P26368
A	373	GLY	-	EXPRESSION TAG	UNP P26368
A	374	SER	-	EXPRESSION TAG	UNP P26368
C	370	GLY	-	EXPRESSION TAG	UNP P26368
C	371	PRO	-	EXPRESSION TAG	UNP P26368
C	372	LEU	-	EXPRESSION TAG	UNP P26368
C	373	GLY	-	EXPRESSION TAG	UNP P26368
C	374	SER	-	EXPRESSION TAG	UNP P26368

- Molecule 2 is a protein called Splicing factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				Se
2	B	101	813	500	151	158	2	2	0	1	0
2	D	107	872	539	161	168	2	2	0	4	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	9	GLY	-	EXPRESSION TAG	UNP Q15637
B	10	PRO	-	EXPRESSION TAG	UNP Q15637

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	11	LEU	-	EXPRESSION TAG	UNP Q15637
B	12	GLY	-	EXPRESSION TAG	UNP Q15637
B	13	SER	-	EXPRESSION TAG	UNP Q15637
D	9	GLY	-	EXPRESSION TAG	UNP Q15637
D	10	PRO	-	EXPRESSION TAG	UNP Q15637
D	11	LEU	-	EXPRESSION TAG	UNP Q15637
D	12	GLY	-	EXPRESSION TAG	UNP Q15637
D	13	SER	-	EXPRESSION TAG	UNP Q15637

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	8	Total O 8 8	0	0
4	B	9	Total O 9 9	0	0

Continued on next page...

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	C	14	Total	O	0	0
			14	14		
4	D	25	Total	O	0	0
			25	25		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Splicing factor U2AF 65 kDa subunit

Chain A: 




- Molecule 1: Splicing factor U2AF 65 kDa subunit

Chain C: 





- Molecule 2: Splicing factor 1

Chain B: 



- Molecule 2: Splicing factor 1

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.80Å 116.10Å 130.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.56 – 2.29 43.30 – 2.29	Depositor EDS
% Data completeness (in resolution range)	89.4 (34.56-2.29) 89.1 (43.30-2.29)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.72 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, $R_{free}$	0.248 , 0.281 0.246 , 0.275	Depositor DCC
$R_{free}$ test set	2318 reflections (8.77%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.0	Xtrriage
Anisotropy	0.902	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.24 , 57.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3409	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	105.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SEP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/847	0.46	0/1142
1	C	0.22	0/836	0.40	0/1122
2	B	0.22	0/807	0.42	0/1088
2	D	0.22	0/878	0.43	0/1184
All	All	0.22	0/3368	0.43	0/4536

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	409	TYR	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	830	0	788	27	0
1	C	823	0	790	10	0
2	B	813	0	773	34	0
2	D	872	0	824	21	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	2	0
4	A	8	0	0	0	0
4	B	9	0	0	1	0
4	C	14	0	0	0	0
4	D	25	0	0	2	0
All	All	3409	0	3175	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:57:GLN:HA	2:B:57:GLN:NE2	1.55	1.14
2:B:57:GLN:HE21	2:B:57:GLN:CA	1.60	1.12
2:B:24[A]:GLN:HA	2:B:24[A]:GLN:OE1	1.75	0.85
1:C:416:GLU:HG2	1:C:474:PHE:CZ	2.14	0.83
2:B:57:GLN:HA	2:B:57:GLN:HE21	0.71	0.81
2:D:24[B]:GLN:H	2:D:24[B]:GLN:CD	1.83	0.77
1:A:420:PRO:HA	1:A:421:VAL:CG2	2.17	0.75
2:D:24[A]:GLN:NE2	4:D:223:HOH:O	2.20	0.74
2:B:24[A]:GLN:OE1	2:B:24[A]:GLN:CA	2.36	0.73
2:B:21:ARG:HH11	2:B:21:ARG:HB2	1.54	0.73
2:D:55:GLN:CG	2:D:116:MSE:HE2	2.22	0.69
1:A:405:GLU:OE1	1:A:452:ARG:NH2	2.25	0.69
1:A:420:PRO:HA	1:A:421:VAL:HG22	1.73	0.68
2:B:24[A]:GLN:OE1	2:B:25:ASP:N	2.26	0.67
2:D:55:GLN:HG3	2:D:116:MSE:HE2	1.79	0.65
2:D:55:GLN:HG3	2:D:116:MSE:CE	2.27	0.65
2:D:55:GLN:CG	2:D:116:MSE:CE	2.75	0.64
1:A:452:ARG:NH1	2:B:23:ASN:O	2.27	0.64
2:B:23:ASN:OD1	4:B:308:HOH:O	2.15	0.63
1:A:419:ARG:O	1:A:421:VAL:HG22	1.98	0.62
1:A:409:TYR:CD2	1:A:409:TYR:N	2.68	0.61
2:B:116:MSE:O	2:B:120:ASN:ND2	2.34	0.61
1:A:467:ASP:O	1:A:471[B]:ARG:HD3	2.01	0.60

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:420:PRO:CA	1:A:421:VAL:HG22	2.35	0.57
2:B:67:THR:N	2:B:68:GLY:HA2	2.19	0.57
1:A:401:ASP:OD2	2:B:21:ARG:NH1	2.39	0.56
2:D:55:GLN:HG2	2:D:116:MSE:HE2	1.87	0.56
2:D:110:HIS:O	2:D:113:ILE:HG22	2.05	0.55
1:C:381:MSE:HE2	2:D:53:ILE:HD11	1.87	0.55
1:A:395:TYR:CE1	1:A:419:ARG:HD2	2.43	0.54
2:B:57:GLN:NE2	2:B:57:GLN:CA	2.36	0.54
1:C:450:THR:HG22	2:D:38:THR:CG2	2.37	0.53
1:C:450:THR:HG22	2:D:38:THR:HG23	1.90	0.53
1:A:425:GLU:O	1:A:425:GLU:HG2	2.08	0.53
1:C:403:ARG:NH2	3:C:501:SO4:O1	2.42	0.52
1:A:426:VAL:CG2	1:A:427:PRO:HD2	2.40	0.51
1:A:475:TRP:CG	2:B:46:ARG:NH1	2.79	0.51
2:D:126:PRO:HB3	2:D:129:TYR:CD2	2.45	0.50
1:A:381:MSE:HE1	2:B:40:ILE:HG13	1.93	0.50
1:A:468:SER:HA	1:A:471[B]:ARG:NH1	2.25	0.50
1:A:475:TRP:CZ2	2:B:46:ARG:HA	2.47	0.50
2:B:119:LEU:HD12	2:B:119:LEU:O	2.12	0.50
1:A:475:TRP:HZ2	2:B:46:ARG:HA	1.78	0.49
1:A:383:MSE:SE	1:A:459:VAL:HG22	2.62	0.49
1:C:425:GLU:O	1:C:425:GLU:HG3	2.12	0.48
2:D:128:ASP:N	2:D:128:ASP:OD1	2.45	0.48
2:D:24[A]:GLN:OE1	2:D:24[A]:GLN:HA	2.13	0.48
1:A:379:CYS:HB2	1:A:464:CYS:SG	2.53	0.48
1:A:453:LYS:HB2	2:B:23:ASN:CG	2.33	0.48
1:A:407:SER:HA	1:A:410:GLY:HA3	1.96	0.47
2:B:114:THR:O	2:B:117:VAL:HG12	2.14	0.47
1:C:395:TYR:CD1	1:C:419:ARG:NH1	2.84	0.46
2:B:24[B]:GLN:OE1	2:B:24[B]:GLN:N	2.47	0.46
1:C:424:VAL:O	1:C:425:GLU:CG	2.63	0.46
2:B:67:THR:HB	2:B:69:ASP:N	2.31	0.46
1:C:440:PHE:CE2	1:C:444:LYS:HD2	2.50	0.46
2:B:67:THR:HB	2:B:68:GLY:CA	2.46	0.45
1:A:420:PRO:HD3	1:A:429:CYS:SG	2.56	0.45
2:D:24[B]:GLN:NE2	4:D:222:HOH:O	2.50	0.45
2:D:109:ARG:NH2	2:D:129:TYR:HB2	2.32	0.44
2:B:64:LYS:O	2:B:69:ASP:N	2.51	0.44
2:B:24[B]:GLN:O	2:B:24[B]:GLN:HG2	2.18	0.44
2:D:55:GLN:HG3	2:D:116:MSE:HE1	1.98	0.44
2:B:58:ILE:CD1	2:B:112:LEU:HB3	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:123:PHE:CD1	2:B:123:PHE:O	2.71	0.43
2:B:126:PRO:O	2:B:127:ALA:HB3	2.18	0.43
2:D:117:VAL:HG22	2:D:123:PHE:CD1	2.54	0.43
2:B:128:ASP:OD2	2:B:128:ASP:C	2.56	0.43
2:B:80:SEP:O1P	2:B:97:ARG:NH1	2.53	0.42
1:A:453:LYS:HB2	2:B:23:ASN:CB	2.50	0.42
2:B:20:SER:OG	2:B:21:ARG:N	2.52	0.42
1:A:382:ASN:HD22	1:A:458:VAL:HG13	1.83	0.42
2:D:117:VAL:CG2	2:D:123:PHE:CD1	3.03	0.41
1:A:409:TYR:N	1:A:409:TYR:HD2	2.16	0.41
1:A:409:TYR:CD1	1:A:445:ALA:HA	2.55	0.41
1:A:421:VAL:HG23	1:A:421:VAL:O	2.21	0.41
1:C:415:ILE:N	3:C:501:SO4:O4	2.53	0.41
2:D:126:PRO:HB3	2:D:129:TYR:HB3	2.02	0.41
2:B:73:PRO:HA	2:B:74:PRO:HD3	1.95	0.40
2:B:67:THR:HB	2:B:68:GLY:C	2.42	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	106/106 (100%)	102 (96%)	4 (4%)	0	100	100
1	C	100/106 (94%)	97 (97%)	3 (3%)	0	100	100
2	B	96/124 (77%)	88 (92%)	7 (7%)	1 (1%)	15	17
2	D	103/124 (83%)	96 (93%)	6 (6%)	1 (1%)	15	17
All	All	405/460 (88%)	383 (95%)	20 (5%)	2 (0%)	29	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	42	PRO
2	D	43	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/93 (96%)	86 (97%)	3 (3%)	37	51
1	C	92/93 (99%)	88 (96%)	4 (4%)	29	40
2	B	84/109 (77%)	74 (88%)	10 (12%)	5	5
2	D	90/109 (83%)	87 (97%)	3 (3%)	38	53
All	All	355/404 (88%)	335 (94%)	20 (6%)	21	29

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	385	LEU
1	A	401	ASP
1	A	409	TYR
2	B	21	ARG
2	B	23	ASN
2	B	24[A]	GLN
2	B	24[B]	GLN
2	B	57	GLN
2	B	66	ARG
2	B	105	LEU
2	B	117	VAL
2	B	119	LEU
2	B	123	PHE
1	C	390	LEU
1	C	397	GLU
1	C	401	ASP
1	C	446	MSE
2	D	113	ILE
2	D	116	MSE
2	D	123	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	57	GLN
2	B	120	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SEP	D	80	2	8,9,10	1.61	1 (12%)	7,12,14	1.22	1 (14%)
2	SEP	D	82	2	8,9,10	1.61	1 (12%)	7,12,14	1.46	1 (14%)
2	SEP	B	80	2	8,9,10	1.63	1 (12%)	7,12,14	1.45	1 (14%)
2	SEP	B	82	2	8,9,10	1.63	1 (12%)	7,12,14	1.31	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	D	80	2	-	2/6/8/10	-
2	SEP	D	82	2	-	2/6/8/10	-
2	SEP	B	80	2	-	2/6/8/10	-
2	SEP	B	82	2	-	2/6/8/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	82	SEP	P-O1P	3.55	1.61	1.50
2	B	80	SEP	P-O1P	3.53	1.61	1.50
2	D	82	SEP	P-O1P	3.51	1.61	1.50
2	D	80	SEP	P-O1P	3.50	1.61	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	80	SEP	OG-CB-CA	3.28	111.33	108.14
2	D	82	SEP	OG-CB-CA	3.27	111.32	108.14
2	B	82	SEP	OG-CB-CA	2.79	110.86	108.14
2	D	80	SEP	OG-CB-CA	2.55	110.63	108.14

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	80	SEP	N-CA-CB-OG
2	B	80	SEP	C-CA-CB-OG
2	B	82	SEP	N-CA-CB-OG
2	B	82	SEP	C-CA-CB-OG
2	D	80	SEP	N-CA-CB-OG
2	D	80	SEP	C-CA-CB-OG
2	D	82	SEP	N-CA-CB-OG
2	D	82	SEP	C-CA-CB-OG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	80	SEP	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	C	501	-	4,4,4	0.24	0	6,6,6	0.06	0
3	SO4	A	501	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	B	201	-	4,4,4	0.23	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	501	SO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	102/106 (96%)	0.19	5 (4%) 29 36	61, 92, 137, 182	2 (1%)
1	C	101/106 (95%)	0.22	4 (3%) 38 45	62, 93, 156, 214	0
2	B	97/124 (78%)	0.25	3 (3%) 49 56	70, 104, 187, 197	1 (1%)
2	D	103/124 (83%)	0.13	4 (3%) 39 46	63, 107, 175, 203	0
All	All	403/460 (87%)	0.19	16 (3%) 38 45	61, 100, 174, 214	3 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	423	GLY	5.2
1	A	411	LEU	4.2
2	D	32	VAL	3.9
2	B	123	PHE	3.7
1	A	421	VAL	3.7
1	A	422	ASP	2.8
2	D	124	LYS	2.6
2	B	20	SER	2.5
1	C	389	LEU	2.5
1	C	424	VAL	2.4
2	D	128	ASP	2.4
2	D	70	LEU	2.2
1	A	426	VAL	2.1
2	B	125	PRO	2.1
1	A	424	VAL	2.0
1	C	475	TRP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SEP	B	80	10/11	0.91	0.13	111,115,157,174	0
2	SEP	D	80	10/11	0.94	0.10	113,125,144,144	0
2	SEP	D	82	10/11	0.96	0.11	86,100,108,113	0
2	SEP	B	82	10/11	0.97	0.08	88,107,121,121	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	B	201	5/5	0.73	0.12	216,217,217,218	0
3	SO4	A	501	5/5	0.76	0.21	171,172,173,174	0
3	SO4	C	501	5/5	0.93	0.16	167,168,168,169	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.