



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 03:54 PM EDT

PDB ID : 3G3H  
Title : Crystal structure of the GluR6 ligand binding domain dimer K665R I749L Q753K mutant with glutamate and NaCl at 1.5 Angstrom resolution  
Authors : Chaudhry, C.; Mayer, M.L.  
Deposited on : 2009-02-02  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	 4% 86% 10% ..
1	B	259	 2% 86% 11% .

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9422 atoms, of which 4403 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor, ionotropic kainate 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	252	4294	1361	2172	346	400	15	0	19	0
1	B	253	4369	1381	2218	349	406	15	0	23	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P42260
A	118	GLY	-	linker	UNP P42260
A	119	THR	-	linker	UNP P42260
A	149	ARG	LYS	engineered mutation	UNP P42260
A	233	LEU	ILE	engineered mutation	UNP P42260
A	237	LYS	GLN	engineered mutation	UNP P42260
B	1	GLY	-	expression tag	UNP P42260
B	118	GLY	-	linker	UNP P42260
B	119	THR	-	linker	UNP P42260
B	149	ARG	LYS	engineered mutation	UNP P42260
B	233	LEU	ILE	engineered mutation	UNP P42260
B	237	LYS	GLN	engineered mutation	UNP P42260

- Molecule 2 is GLUTAMIC ACID (three-letter code: GLU) (formula: C<sub>5</sub>H<sub>9</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	H	N	O	0	0
			18	5	8	1	4		
2	B	1	Total	C	H	N	O	0	0
			15	5	5	1	4		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Na	0	0
			2	2		
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	317	Total	O	0	11
			323	323		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	B	393	Total 398	O 398	0	16


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate receptor, ionotropic kainate 2



- Molecule 1: Glutamate receptor, ionotropic kainate 2



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.14Å 113.50Å 52.16Å 90.00° 115.14° 90.00°	Depositor
Resolution (Å)	29.52 – 1.50 29.53 – 1.50	Depositor EDS
% Data completeness (in resolution range)	97.8 (29.52-1.50) 97.8 (29.53-1.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.07 (at 1.50Å)	Xtrriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.147 , 0.176 0.150 , 0.177	Depositor DCC
$R_{free}$ test set	4201 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.5	Xtrriage
Anisotropy	0.058	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 51.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.025 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9422	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	0/2217	0.80	1/2983 (0.0%)
1	B	0.71	0/2254	0.86	2/3029 (0.1%)
All	All	0.69	0/4471	0.83	3/6012 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	162	MET	CG-SD-CE	-9.19	85.50	100.20
1	A	4	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	B	96	ARG	NE-CZ-NH2	-5.54	117.53	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2122	2172	2167	26	0
1	B	2151	2218	2212	26	0
2	A	10	8	5	0	0
2	B	10	5	5	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	2	0	0	0	0
4	B	1	0	0	0	0
5	A	323	0	0	6	1
5	B	398	0	0	10	1
All	All	5019	4403	4389	51	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:GLU:HG3	5:A:644:HOH:O	1.67	0.95
1:A:94:TYR:CZ	1:A:98:LYS:HE3	2.14	0.83
1:B:94:TYR:CZ	1:B:98:LYS:HE3	2.17	0.78
1:B:240[A]:GLU:OE1	5:B:707[A]:HOH:O	2.02	0.76
1:B:228[A]:ARG:CZ	5:B:422[A]:HOH:O	2.38	0.71
1:B:65:ASP:OD2	1:B:67:VAL:HG12	1.91	0.70
1:A:228[A]:ARG:CZ	5:A:460[A]:HOH:O	2.46	0.63
1:B:97:GLU:OE2	5:B:685[A]:HOH:O	2.17	0.59
1:B:45:THR:HG21	5:B:443:HOH:O	2.03	0.57
1:A:239[A]:GLN:CG	1:A:244:LEU:HD12	2.36	0.56
1:B:147:PHE:CE2	1:B:189[B]:LEU:HD23	2.41	0.55
1:B:228[A]:ARG:NH2	5:B:422[A]:HOH:O	2.40	0.55
1:A:228[A]:ARG:NH1	5:A:460[A]:HOH:O	2.39	0.55
1:A:239[B]:GLN:OE1	1:A:244:LEU:HD12	2.06	0.54
1:B:94:TYR:OH	1:B:98:LYS:HE3	2.07	0.54
1:A:147:PHE:CE2	1:A:189[B]:LEU:HD13	2.43	0.53
1:A:153:SER:O	1:A:156:ASP:OD2	2.28	0.51
1:B:239[A]:GLN:CG	1:B:244:LEU:HD12	2.42	0.50
1:A:21:LYS:HD2	5:A:463:HOH:O	2.10	0.50
1:B:65:ASP:OD2	1:B:67:VAL:CG1	2.58	0.49
1:B:57[B]:GLU:HG3	5:B:704[B]:HOH:O	2.13	0.49
1:A:147:PHE:CE2	1:A:189[A]:LEU:HD23	2.49	0.48
1:A:239[B]:GLN:NE2	1:B:214[B]:SER:HB3	2.28	0.48
1:A:172:LYS:HG3	5:A:680:HOH:O	2.14	0.47
1:A:153:SER:HA	1:A:156:ASP:OD2	2.15	0.46
1:B:13[B]:GLU:HG3	1:B:16[B]:TYR:HD2	1.80	0.46
1:A:14:GLU:HG3	1:A:18:LEU:HD21	1.96	0.46
1:A:94:TYR:OH	1:A:98:LYS:HE3	2.14	0.46
1:A:136:GLY:HA2	1:A:162[B]:MET:SD	2.56	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:196:GLU:O	1:B:200:GLN:HG2	2.16	0.45
1:A:239[A]:GLN:HG2	1:A:244:LEU:HD12	1.99	0.45
1:B:57[B]:GLU:CG	5:B:704[B]:HOH:O	2.65	0.45
1:A:116[A]:ARG:NH2	1:A:185:ASP:HA	2.32	0.45
1:A:239[A]:GLN:HG3	1:A:244:LEU:HD12	1.98	0.45
1:B:200:GLN:HG2	5:B:534:HOH:O	2.17	0.44
1:A:135:TYR:HA	1:A:187:ALA:O	2.18	0.44
1:B:239[A]:GLN:HG2	1:B:244:LEU:HD12	1.99	0.43
1:A:65:ASP:HB3	1:A:68:ASN:OD1	2.18	0.43
1:B:135:TYR:HA	1:B:187:ALA:O	2.19	0.42
1:B:175:GLU:HG2	5:B:643:HOH:O	2.18	0.42
1:A:65:ASP:OD1	1:A:67:VAL:HG13	2.20	0.42
1:A:88:ALA:HB1	1:A:89:PRO:HD2	2.02	0.42
1:A:113:ILE:HB	1:A:208:ILE:HB	2.03	0.41
1:A:41[B]:ARG:HG3	5:A:627:HOH:O	2.20	0.41
1:A:94:TYR:OH	1:A:98:LYS:CE	2.69	0.41
1:B:47:LEU:HD21	5:B:473:HOH:O	2.21	0.41
1:B:166:ARG:HG3	1:B:167:GLN:N	2.36	0.40
1:B:14:GLU:HG3	1:B:18:LEU:HD21	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:517:HOH:O	5:B:630:HOH:O[2_655]	2.18	0.02

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	267/259 (103%)	261 (98%)	6 (2%)	0	<b>100</b> <b>100</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	274/259 (106%)	271 (99%)	3 (1%)	0	100	100
All	All	541/518 (104%)	532 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/225 (107%)	235 (98%)	5 (2%)	53	23
1	B	245/225 (109%)	242 (99%)	3 (1%)	71	48
All	All	485/450 (108%)	477 (98%)	8 (2%)	67	36

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	VAL
1	A	109	LEU
1	A	156	ASP
1	A	189[A]	LEU
1	A	189[B]	LEU
1	B	109	LEU
1	B	228[A]	ARG
1	B	228[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	204	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLU	A	260	-	2,9,9	1.51	1 (50%)	2,11,11	0.65	0
2	GLU	B	260	-	2,9,9	0.63	0	2,11,11	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLU	A	260	-	-	0/3/9/9	-
2	GLU	B	260	-	-	0/3/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	260	GLU	CA-N	2.13	1.51	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	252/259 (97%)	-0.09	10 (3%) 38 42	7, 14, 32, 65	0
1	B	253/259 (97%)	-0.25	6 (2%) 59 63	6, 12, 28, 54	0
All	All	505/518 (97%)	-0.17	16 (3%) 47 52	6, 13, 30, 65	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	PRO	7.7
1	B	1	GLY	6.4
1	A	257	CYS	6.1
1	B	67	VAL	5.0
1	B	257	CYS	4.8
1	A	67	VAL	4.1
1	A	68	ASN	3.7
1	A	3	ASN	3.5
1	B	2	SER	3.4
1	A	69	GLY	2.7
1	A	203	CYS	2.5
1	A	204[A]	ASN	2.1
1	A	70	GLN	2.1
1	B	66	ASP	2.1
1	A	201	ARG	2.0
1	B	68	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NA	A	263	1/1	0.92	0.13	31,31,31,31	1
4	NA	A	262	1/1	0.98	0.08	14,14,14,14	1
3	CL	B	261	1/1	0.98	0.05	14,14,14,14	1
4	NA	B	262	1/1	0.98	0.06	10,10,10,10	1
2	GLU	A	260	10/10	0.99	0.09	7,9,11,11	0
2	GLU	B	260	10/10	0.99	0.09	5,6,8,8	0
3	CL	A	261	1/1	1.00	0.09	9,9,9,9	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.