

wwPDB X-ray Structure Validation Summary Report (i)

Feb 11, 2024 – 04:28 PM EST

PDB ID : 3G60

Title : Structure of P-glycoprotein Reveals a Molecular Basis for Poly-Specific Drug

Binding

Authors: Aller, S.G.; Yu, J.; Ward, A.; Weng, Y.; Chittaboina, S.; Zhuo, R.; Harrell,

P.M.; Trinh, Y.T.; Zhang, Q.; Urbatsch, I.L.; Chang, G.

Deposited on : 2009-02-05

Resolution : 4.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution		
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$		
R_{free}	130704	1043 (5.00-3.80)		
Clashscore	141614	1111 (5.00-3.80)		
Ramachandran outliers	138981	1059 (5.00-3.80)		
Sidechain outliers	138945	1041 (5.00-3.80)		
RSRZ outliers	127900	1095 (5.08-3.70)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length		Quality of chain			
1	A	1284	16%	59%	15%	•	8%
1	В	1284	16%	59%	16%	•	8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	0JZ	A	6001	_	-	-	X
2	0JZ	В	6002	-	-	-	X



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 18414 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Multidrug resistance protein 1a.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	1182	Total 9171	C 5895	N 1552	O 1686	S 38	0	0	0
1	В	1182	Total 9171	C 5895	N 1552	O 1686	S 38	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1277	TYR	-	expression tag	UNP Q5I1Y5
A	1278	VAL	-	expression tag	UNP Q5I1Y5
A	1279	HIS	-	expression tag	UNP Q5I1Y5
A	1280	HIS	-	expression tag	UNP Q5I1Y5
A	1281	HIS	-	expression tag	UNP Q5I1Y5
A	1282	HIS	-	expression tag	UNP Q5I1Y5
A	1283	HIS	-	expression tag	UNP Q5I1Y5
A	1284	HIS	-	expression tag	UNP Q5I1Y5
В	1277	TYR	-	expression tag	UNP Q5I1Y5
В	1278	VAL	-	expression tag	UNP Q5I1Y5
В	1279	HIS	_	expression tag	UNP Q5I1Y5
В	1280	HIS	-	expression tag	UNP Q5I1Y5
В	1281	HIS	-	expression tag	UNP Q5I1Y5
В	1282	HIS	-	expression tag	UNP Q5I1Y5
В	1283	HIS	-	expression tag	UNP Q5I1Y5
В	1284	HIS	-	expression tag	UNP Q5I1Y5

• Molecule 2 is (4R,11R,18R)-4,11,18-tri(propan-2-yl)-6,13,20-triselena-3,10,17,22,23,24-hexaa zatetracyclo[17.2.1.1 5,8 .1 12,15] tetracosa-1(21),5(24),7,12(23),14,19(22)-hexaene-2,9,16-tri one (three-letter code: 0JZ) (formula: $C_{24}H_{30}N_6O_3Se_3$).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	Se	0	0	
	A	1	36	24	6	3	3	U		
9	D	1	Total	С	N	О	Se	0	0	
2	Б	1	36	24	6	3	3	U	0	




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Multidrug resistance protein 1a









• Molecule 1: Multidrug resistance protein 1a













4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	96.63Å 115.09Å 374.20Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 - 4.40	Depositor
rtesolution (A)	49.02 - 4.31	EDS
% Data completeness	94.7 (19.99-4.40)	Depositor
(in resolution range)	93.3 (49.02-4.31)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) > 1$	5.10 (at 4.29Å)	Xtriage
Refinement program	CNS	Depositor
P. P.	0.314 , 0.365	Depositor
R, R_{free}	0.297 , 0.351	DCC
R_{free} test set	2835 reflections (9.91%)	wwPDB-VP
Wilson B-factor (Å ²)	201.9	Xtriage
Anisotropy	0.359	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.21 , 103.6	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	18414	wwPDB-VP
Average B, all atoms (Å ²)	198.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.39% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 0JZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain			nd lengths	Bond angles		
WIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.41	1/9339~(0.0%)	0.70	13/12626 (0.1%)	
1	В	0.41	0/9339	0.68	8/12626 (0.1%)	
All	All	0.41	$1/18678 \; (0.0\%)$	0.69	21/25252 (0.1%)	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
1	A	851	TRP	CB-CG	5.17	1.59	1.50

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	A	427	CYS	N-CA-C	9.35	136.24	111.00
1	A	1098	LYS	N-CA-C	-7.80	89.94	111.00
1	A	164	VAL	N-CA-C	-7.79	89.98	111.00
1	A	267	LYS	N-CA-C	7.23	130.52	111.00
1	A	165	GLY	N-CA-C	-7.16	95.21	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9171	0	9344	1558	0
1	В	9171	0	9344	1534	0
2	A	36	0	27	12	0
2	В	36	0	27	10	0
All	All	18414	0	18742	3089	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 83.

The worst 5 of 3089 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:797:VAL:O	1:B:801:ASP:HB2	1.45	1.15
1:A:523:ARG:HD3	1:A:524:GLY:H	0.99	1.13
1:B:523:ARG:HD3	1:B:524:GLY:H	0.98	1.12
1:B:858:LEU:O	1:B:862:PRO:HD2	1.51	1.11
1:B:1204:THR:O	1:B:1206:SER:N	1.83	1.10

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1178/1284 (92%)	704 (60%)	302 (26%)	172 (15%)	0 4
1	В	1178/1284 (92%)	707 (60%)	305 (26%)	166 (14%)	0 4
All	All	$2356/2568 \; (92\%)$	1411 (60%)	607 (26%)	338 (14%)	0 4

5 of 338 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	34	SER

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	35	VAL
1	A	52	VAL
1	A	89	THR
1	A	133	CYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	976/1065 (92%)	884 (91%)	92 (9%)	8	8	30
1	В	976/1065 (92%)	880 (90%)	96 (10%)	8	8	29
All	All	1952/2130 (92%)	1764 (90%)	188 (10%)	8	8	29

5 of 188 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	238	LYS
1	В	694	TRP
1	В	254	LEU
1	В	332	PHE
1	В	771	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 83 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	437	GLN
1	В	963	GLN
1	В	515	GLN
1	В	805	ASN
1	В	1099	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	True	Chain	in Res Linl		Link Bond lengths				Bond angles		
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	0JZ	В	6002	-	21,39,39	1.74	4 (19%)	24,57,57	1.91	7 (29%)	
2	0JZ	A	6001	-	21,39,39	2.04	5 (23%)	24,57,57	2.49	9 (37%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0JZ	В	6002	-	-	4/24/48/48	0/3/4/4
2	0JZ	A	6001	-	-	4/24/48/48	0/3/4/4

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(Å)
2	A	6001	0JZ	C16-N17	5.48	1.46	1.34
2	В	6002	0JZ	C09-N10	4.44	1.43	1.34
2	A	6001	0JZ	C02-N03	4.39	1.43	1.34
2	A	6001	0JZ	C09-N10	3.88	1.42	1.34
2	В	6002	0JZ	C02-N03	3.70	1.42	1.34



The worst	5	$\circ f$	16	bond	angle	outliers	are	listed	below:
THE WOLDS	v	$O_{\rm I}$	10	DOM	angic	Outilities	arc	nsucu	DCIOW.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	6001	0JZ	C04-N03-C02	5.49	131.07	122.28
2	A	6001	0JZ	C18-N17-C16	5.23	130.66	122.28
2	A	6001	0JZ	C35-C34-C18	4.30	115.38	111.24
2	В	6002	0JZ	C33-C31-C11	3.77	114.87	111.24
2	В	6002	0JZ	C29-C28-C04	3.75	114.85	111.24

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	6001	0JZ	C05-C04-N03-C02
2	A	6001	0JZ	C28-C04-N03-C02
2	A	6001	0JZ	C19-C18-N17-C16
2	A	6001	0JZ	C34-C18-N17-C16
2	В	6002	0JZ	C05-C04-N03-C02


There are no ring outliers.

2 monomers are involved in 22 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	6002	0JZ	10	0
2	A	6001	0JZ	12	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	$1182/1284\ (92\%)$	-0.12	20 (1%) 70 61	118, 194, 243, 306	0
1	В	1182/1284~(92%)	-0.13	21 (1%) 68 59	123, 200, 243, 301	0
All	All	$2364/2568 \; (92\%)$	-0.12	41 (1%) 70 61	118, 197, 244, 306	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	963	GLN	4.9
1	В	1199	THR	4.5
1	A	1084	ASP	4.0
1	A	574	GLU	3.9
1	В	228	TRP	3.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ m B ext{-}factors(\AA^2)$	Q<0.9
2	0JZ	В	6002	36/36	0.63	0.41	196,196,196,196	0
2	0JZ	A	6001	36/36	0.64	0.43	196,196,196,196	0

Page 19

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

