



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 7, 2024 – 06:58 am GMT

PDB ID : 6G7R
Title : Structure of fully reduced variant E28Q of E. coli hydrogenase-1 at pH 8
Authors : Carr, S.B.; Armstrong, F.A.; Evans, R.M.
Deposited on : 2018-04-06
Resolution : 1.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 130704 | 1223 (1.22-1.18) |
| Clashscore | 141614 | 1286 (1.22-1.18) |
| Ramachandran outliers | 138981 | 1240 (1.22-1.18) |
| Sidechain outliers | 138945 | 1239 (1.22-1.18) |
| RSRZ outliers | 127900 | 1200 (1.22-1.18) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---|
| 1 | S | 335 |  75% 21% |
| 1 | T | 335 |  74% 21% |
| 2 | L | 582 |  95% 5% |
| 2 | M | 582 |  96% |

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 28652 atoms, of which 13324 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogenase-1 small chain.

| Mol | Chain | Residues | Atoms | | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|------|-----|-----|----|---------|---------|-------|
| | | | Total | C | H | N | O | S | | | |
| 1 | S | 263 | 4110 | 1319 | 2034 | 359 | 378 | 20 | 148 | 7 | 0 |
| 1 | T | 263 | 4173 | 1336 | 2070 | 367 | 379 | 21 | 157 | 10 | 0 |

There are 16 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| S | 328 | ARG | - | expression tag | UNP P69739 |
| S | 329 | SER | - | expression tag | UNP P69739 |
| S | 330 | HIS | - | expression tag | UNP P69739 |
| S | 331 | HIS | - | expression tag | UNP P69739 |
| S | 332 | HIS | - | expression tag | UNP P69739 |
| S | 333 | HIS | - | expression tag | UNP P69739 |
| S | 334 | HIS | - | expression tag | UNP P69739 |
| S | 335 | HIS | - | expression tag | UNP P69739 |
| T | 328 | ARG | - | expression tag | UNP P69739 |
| T | 329 | SER | - | expression tag | UNP P69739 |
| T | 330 | HIS | - | expression tag | UNP P69739 |
| T | 331 | HIS | - | expression tag | UNP P69739 |
| T | 332 | HIS | - | expression tag | UNP P69739 |
| T | 333 | HIS | - | expression tag | UNP P69739 |
| T | 334 | HIS | - | expression tag | UNP P69739 |
| T | 335 | HIS | - | expression tag | UNP P69739 |


- Molecule 2 is a protein called Hydrogenase-1 large chain.

| Mol | Chain | Residues | Atoms | | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|------|-----|-----|----|---------|---------|-------|
| | | | Total | C | H | N | O | S | | | |
| 2 | L | 581 | 9248 | 2961 | 4586 | 817 | 856 | 28 | 375 | 21 | 0 |
| 2 | M | 581 | 9195 | 2948 | 4560 | 807 | 853 | 27 | 361 | 17 | 0 |

There are 2 discrepancies between the modelled and reference sequences:


| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|------------|
| L | 28 | GLN | GLU | conflict | UNP P0ACD8 |
| M | 28 | GLN | GLU | conflict | UNP P0ACD8 |

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|---------|---------|
| 3 | S | 1 | Total | Fe | S | 0 | 0 |
| | | | 8 | 4 | 4 | | |
| 3 | T | 1 | Total | Fe | S | 0 | 0 |
| | | | 8 | 4 | 4 | | |

- Molecule 4 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe₃S₄).




| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|---------|---------|
| 4 | S | 1 | Total | Fe | S | 0 | 0 |
| | | | 7 | 3 | 4 | | |
| 4 | T | 1 | Total | Fe | S | 0 | 0 |
| | | | 7 | 3 | 4 | | |

- Molecule 5 is FE4-S3 CLUSTER (three-letter code: SF3) (formula: Fe₄S₃).



| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|---------|---------|
| 5 | S | 1 | Total | Fe | S | 0 | 0 |
| | | | 7 | 4 | 3 | | |
| 5 | T | 1 | Total | Fe | S | 0 | 0 |
| | | | 7 | 4 | 3 | | |

- Molecule 6 is DODECYL-BETA-D-MALTOSE (three-letter code: LMT) (formula: $C_{24}H_{46}O_{11}$).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|----|----|---------|---------|
| | | | Total | C | H | O | | |
| 6 | S | 1 | 81 | 24 | 46 | 11 | 6 | 0 |
| 6 | T | 1 | 40 | 13 | 26 | 1 | 1 | 0 |

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| | | | Total | Cl | | |
| 7 | S | 2 | 2 | 2 | 0 | 0 |
| 7 | T | 2 | 2 | 2 | 0 | 0 |

- Molecule 8 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 8 | L | 1 | Total | O | S | 0 | 0 |
| | | | 5 | 4 | 1 | | |

- Molecule 9 is NI-FE REDUCED ACTIVE CENTER (three-letter code: EJ2) (formula: C₃HFeN₂NiO).



| Mol | Chain | Residues | Atoms | | | | | | ZeroOcc | AltConf | |
|-----|-------|----------|-------|---|----|---|---|----|---------|---------|---|
| 9 | L | 1 | Total | C | Fe | H | N | Ni | O | 0 | 0 |
| | | | 9 | 3 | 1 | 1 | 2 | 1 | 1 | | |
| 9 | M | 1 | Total | C | Fe | H | N | Ni | O | 0 | 0 |
| | | | 9 | 3 | 1 | 1 | 2 | 1 | 1 | | |

- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|------------|---------|---------|---------|
| 10 | L | 1 | Total 1 | Mg 1 | 0 | 0 |
| 10 | M | 1 | Total 1 | Mg 1 | 0 | 0 |

- Molecule 11 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|--------------|----------|---------|---------|
| 11 | S | 241 | Total 241 | O 241 | 0 | 0 |
| 11 | L | 590 | Total 590 | O 590 | 0 | 0 |
| 11 | T | 287 | Total 287 | O 287 | 0 | 0 |
| 11 | M | 614 | Total 614 | O 614 | 0 | 0 |

4 Data and refinement statistics i

| Property | Value | Source |
|---|---|------------------|
| Space group | P 21 21 21 | Depositor |
| Cell constants a, b, c, α , β , γ | 94.07Å 97.81Å 183.21Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 67.89 – 1.20 67.80 – 1.20 | Depositor EDS |
| % Data completeness (in resolution range) | 96.5 (67.89-1.20) 96.5 (67.80-1.20) | Depositor EDS |
| R_{merge} | 0.07 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 1.90 (at 1.20Å) | Xtrriage |
| Refinement program | REFMAC 5.8.0222 | Depositor |
| R, R_{free} | 0.111 , 0.133 0.111 , 0.133 | Depositor DCC |
| R_{free} test set | 25633 reflections (5.08%) | wwPDB-VP |
| Wilson B-factor (Å ²) | 10.2 | Xtrriage |
| Anisotropy | 0.637 | Xtrriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.41 , 46.3 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$ | Xtrriage |
| Estimated twinning fraction | 0.000 for k,h,-l | Xtrriage |
| F_o, F_c correlation | 0.99 | EDS |
| Total number of atoms | 28652 | wwPDB-VP |
| Average B, all atoms (Å ²) | 14.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF3, F3S, MG, EJ2, SO4, SF4, CL, LMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | S | 0.54 | 0/2143 | 0.91 | 6/2908 (0.2%) |
| 1 | T | 0.53 | 0/2176 | 0.82 | 4/2951 (0.1%) |
| 2 | L | 0.50 | 1/4844 (0.0%) | 0.76 | 6/6586 (0.1%) |
| 2 | M | 0.50 | 0/4802 | 0.76 | 5/6533 (0.1%) |
| All | All | 0.51 | 1/13965 (0.0%) | 0.79 | 21/18978 (0.1%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | S | 0 | 2 |
| 1 | T | 0 | 3 |
| 2 | L | 0 | 2 |
| 2 | M | 0 | 1 |
| All | All | 0 | 8 |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|--------|------|-------------|----------|
| 2 | L | 561 | GLU | CD-OE1 | 5.37 | 1.31 | 1.25 |

The worst 5 of 21 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|--------|------|-----------|--------|-------------|----------|
| 1 | S | 185[A] | ARG | NE-CZ-NH1 | 10.48 | 125.54 | 120.30 |
| 1 | S | 185[B] | ARG | NE-CZ-NH1 | 10.48 | 125.54 | 120.30 |
| 1 | S | 185[A] | ARG | NE-CZ-NH2 | -10.30 | 115.15 | 120.30 |
| 1 | S | 185[B] | ARG | NE-CZ-NH2 | -10.30 | 115.15 | 120.30 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|------|-------------|----------|
| 2 | L | 89 | TYR | CB-CG-CD1 | 8.40 | 126.04 | 121.00 |

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 2 | L | 169 | ARG | Sidechain |
| 2 | L | 74 | ARG | Sidechain |
| 1 | S | 26 | ARG | Sidechain |
| 1 | S | 260 | ARG | Sidechain |
| 1 | T | 26 | ARG | Sidechain |

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | S | 2076 | 2034 | 2027 | 15 | 0 |
| 1 | T | 2103 | 2070 | 2065 | 25 | 0 |
| 2 | L | 4662 | 4586 | 4590 | 21 | 0 |
| 2 | M | 4635 | 4560 | 4558 | 11 | 0 |
| 3 | S | 8 | 0 | 0 | 0 | 0 |
| 3 | T | 8 | 0 | 0 | 0 | 0 |
| 4 | S | 7 | 0 | 0 | 0 | 0 |
| 4 | T | 7 | 0 | 0 | 0 | 0 |
| 5 | S | 7 | 0 | 0 | 0 | 0 |
| 5 | T | 7 | 0 | 0 | 0 | 0 |
| 6 | S | 35 | 46 | 46 | 1 | 0 |
| 6 | T | 14 | 26 | 25 | 0 | 0 |
| 7 | S | 2 | 0 | 0 | 0 | 0 |
| 7 | T | 2 | 0 | 0 | 0 | 0 |
| 8 | L | 5 | 0 | 0 | 0 | 0 |
| 9 | L | 8 | 1 | 0 | 0 | 0 |
| 9 | M | 8 | 1 | 0 | 0 | 0 |
| 10 | L | 1 | 0 | 0 | 0 | 0 |
| 10 | M | 1 | 0 | 0 | 0 | 0 |
| 11 | L | 590 | 0 | 0 | 13 | 0 |
| 11 | M | 614 | 0 | 0 | 3 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 11 | S | 241 | 0 | 0 | 0 | 0 |
| 11 | T | 287 | 0 | 0 | 2 | 0 |
| All | All | 15328 | 13324 | 13311 | 63 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|-----------------|--------------------------|-------------------|
| 2:L:28:GLN:CD | 11:L:703:HOH:O | 1.73 | 1.24 |
| 1:T:93[B]:ARG:CZ | 1:T:97:GLU:OE1 | 1.98 | 1.12 |
| 2:L:28:GLN:CG | 11:L:703:HOH:O | 1.91 | 1.10 |
| 1:T:101:ARG:HH11 | 1:T:101:ARG:HG3 | 1.25 | 1.00 |
| 2:L:28:GLN:NE2 | 11:L:703:HOH:O | 1.91 | 0.99 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|---------|----------|-------------|-----|
| 1 | S | 268/335 (80%) | 256 (96%) | 12 (4%) | 0 | 100 | 100 |
| 1 | T | 271/335 (81%) | 259 (96%) | 12 (4%) | 0 | 100 | 100 |
| 2 | L | 600/582 (103%) | 585 (98%) | 15 (2%) | 0 | 100 | 100 |
| 2 | M | 596/582 (102%) | 582 (98%) | 14 (2%) | 0 | 100 | 100 |
| All | All | 1735/1834 (95%) | 1682 (97%) | 53 (3%) | 0 | 100 | 100 |

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|-------------|-----|
| 1 | S | 223/274 (81%) | 220 (99%) | 3 (1%) | 69 | 33 |
| 1 | T | 226/274 (82%) | 221 (98%) | 5 (2%) | 52 | 14 |
| 2 | L | 501/481 (104%) | 499 (100%) | 2 (0%) | 91 | 76 |
| 2 | M | 497/481 (103%) | 497 (100%) | 0 | 100 | 100 |
| All | All | 1447/1510 (96%) | 1437 (99%) | 10 (1%) | 86 | 59 |

5 of 10 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|--------|------|
| 1 | T | 174[B] | ARG |
| 1 | T | 191 | TYR |
| 1 | T | 242 | PRO |
| 2 | L | 169 | ARG |
| 2 | L | 524 | ASP |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | T | 13 | HIS |
| 1 | T | 244 | GLN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 6 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 5 | SF3 | T | 403 | 1 | 0,8,8 | - | - | - | | |
| 8 | SO4 | L | 601 | - | 4,4,4 | 0.11 | 0 | 6,6,6 | 0.32 | 0 |
| 3 | SF4 | T | 401 | 1 | 0,12,12 | - | - | - | | |
| 4 | F3S | S | 402 | 1 | 0,9,9 | - | - | - | | |
| 6 | LMT | S | 404 | - | 36,36,36 | 0.68 | 0 | 47,47,47 | 1.26 | 5 (10%) |
| 9 | EJ2 | M | 601 | 2 | 1,6,6 | 2.48 | 1 (100%) | - | | |
| 4 | F3S | T | 402 | 1 | 0,9,9 | - | - | - | | |
| 3 | SF4 | S | 401 | 1 | 0,12,12 | - | - | - | | |
| 6 | LMT | T | 404 | - | 13,13,36 | 0.40 | 0 | 12,12,47 | 0.45 | 0 |
| 5 | SF3 | S | 403 | 1 | 0,8,8 | - | - | - | | |
| 9 | EJ2 | L | 602 | 2 | 1,6,6 | 3.13 | 1 (100%) | - | | |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|------------|---------|
| 5 | SF3 | T | 403 | 1 | - | - | 0/2/2/2 |
| 3 | SF4 | T | 401 | 1 | - | - | 0/6/5/5 |
| 4 | F3S | S | 402 | 1 | - | - | 0/3/3/3 |
| 6 | LMT | S | 404 | - | - | 2/21/61/61 | 0/2/2/2 |
| 4 | F3S | T | 402 | 1 | - | - | 0/3/3/3 |
| 3 | SF4 | S | 401 | 1 | - | - | 0/6/5/5 |
| 6 | LMT | T | 404 | - | - | 2/11/11/61 | - |
| 5 | SF3 | S | 403 | 1 | - | - | 0/2/2/2 |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 9 | L | 602 | EJ2 | O3-C3 | -3.13 | 1.12 | 1.16 |
| 9 | M | 601 | EJ2 | O3-C3 | -2.48 | 1.12 | 1.16 |

All (5) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|-------|-------------|----------|
| 6 | S | 404 | LMT | C6-C5-C4 | 3.26 | 130.97 | 114.42 |
| 6 | S | 404 | LMT | O6B-C6B-C5B | -3.04 | 100.87 | 111.29 |
| 6 | S | 404 | LMT | O5B-C5B-C4B | 3.01 | 115.16 | 109.69 |
| 6 | S | 404 | LMT | C8-C7-C6 | 2.56 | 127.43 | 114.42 |
| 6 | S | 404 | LMT | C6'-C5'-C4' | 2.41 | 120.35 | 113.33 |

There are no chirality outliers.

All (4) torsion outliers are listed below:


| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-----------------|
| 6 | T | 404 | LMT | C2-C1-O1'-C1' |
| 6 | T | 404 | LMT | C1-C2-C3-C4 |
| 6 | S | 404 | LMT | C5-C6-C7-C8 |
| 6 | S | 404 | LMT | C4B-C5B-C6B-O6B |

There are no ring outliers.

1 monomer is involved in 1 short contact:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 6 | S | 404 | LMT | 1 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | | OWAB(Å ²) | Q<0.9 |
|-----|-------|-----------------|--------|---------|-----------|-----------------------|-------|
| 1 | S | 263/335 (78%) | -0.72 | 0 | 100 100 | 7, 11, 18, 28 | 0 |
| 1 | T | 263/335 (78%) | -0.71 | 0 | 100 100 | 7, 11, 20, 29 | 0 |
| 2 | L | 581/582 (99%) | -0.80 | 0 | 100 100 | 7, 12, 26, 49 | 0 |
| 2 | M | 581/582 (99%) | -0.80 | 1 (0%) | 95 94 | 7, 12, 22, 35 | 0 |
| All | All | 1688/1834 (92%) | -0.77 | 1 (0%) | 95 95 | 7, 12, 23, 49 | 0 |

All (1) RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 2 | M | 175 | GLU | 3.1 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 6 | LMT | S | 404 | 35/35 | 0.82 | 0.22 | 22,26,37,49 | 6 |


Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(\AA^2) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|-----------------------------|-------|
| 6 | LMT | T | 404 | 14/35 | 0.90 | 0.12 | 0,18,29,35 | 40 |
| 8 | SO4 | L | 601 | 5/5 | 0.98 | 0.08 | 14,14,18,19 | 5 |
| 7 | CL | T | 406 | 1/1 | 0.99 | 0.12 | 30,30,30,30 | 0 |
| 5 | SF3 | S | 403 | 7/7 | 1.00 | 0.07 | 7,7,7,8 | 0 |
| 5 | SF3 | T | 403 | 7/7 | 1.00 | 0.06 | 8,8,8,8 | 0 |
| 3 | SF4 | S | 401 | 8/8 | 1.00 | 0.07 | 7,7,8,8 | 0 |
| 3 | SF4 | T | 401 | 8/8 | 1.00 | 0.07 | 7,7,7,8 | 0 |
| 7 | CL | S | 405 | 1/1 | 1.00 | 0.04 | 13,13,13,13 | 0 |
| 7 | CL | S | 406 | 1/1 | 1.00 | 0.12 | 27,27,27,27 | 0 |
| 7 | CL | T | 405 | 1/1 | 1.00 | 0.04 | 15,15,15,15 | 0 |
| 4 | F3S | S | 402 | 7/7 | 1.00 | 0.07 | 7,8,8,8 | 0 |
| 4 | F3S | T | 402 | 7/7 | 1.00 | 0.07 | 8,8,8,8 | 0 |
| 9 | EJ2 | L | 602 | 8/8 | 1.00 | 0.06 | 8,9,10,11 | 0 |
| 9 | EJ2 | M | 601 | 8/8 | 1.00 | 0.07 | 8,8,9,11 | 0 |
| 10 | MG | L | 603 | 1/1 | 1.00 | 0.12 | 6,6,6,6 | 0 |
| 10 | MG | M | 602 | 1/1 | 1.00 | 0.12 | 6,6,6,6 | 0 |

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.