



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 18, 2023 – 09:32 AM EST

PDB ID : 1GAX
Title : CRYSTAL STRUCTURE OF THERMUS THERMOPHILUS VALYL-TRNA SYNTHETASE COMPLEXED WITH TRNA(VAL) AND VALYL-ADENYLATE ANALOGUE
Authors : Fukai, S.; Nureki, O.; Sekine, S.; Shimada, A.; Tao, J.; Vassylyev, D.G.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2000-06-23
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RNA backbone	3102	1007 (3.16-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	75	27% 39% 35%
1	D	75	25% 43% 31% .
2	A	862	37% 52% 9% .
2	B	862	36% 51% 12% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 17368 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called TRNA(VA).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	75	1603	714	293	521	75	0	0	0
1	D	75	1603	714	293	521	75	0	0	0


- Molecule 2 is a protein called VALYL-TRNA SYNTHETASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	862	6970	4449	1228	1266	27	0	0	0
2	B	862	6970	4449	1228	1266	27	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		

- Molecule 4 is N-[VALINYLYL]-N'-[ADENOSYL]-DIAMINOSUFONE (three-letter code: VAA) (formula: C₁₅H₂₄N₈O₆S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	Total	C	N	O	S	0	0
			30	15	8	6	1		
4	B	1	Total	C	N	O	S	0	0
			30	15	8	6	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	C	19	Total	O	0	0
			19	19		
5	D	11	Total	O	0	0
			11	11		
5	A	67	Total	O	0	0
			67	67		
5	B	61	Total	O	0	0
			61	61		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: TRNA(VAL)




- Molecule 1: TRNA(VAL)





- Molecule 2: VALYL-TRNA SYNTHETASE





● Molecule 2: VALYL-TRNA SYNTHETASE



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	411.81Å 411.81Å 81.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.90	Depositor
% Data completeness (in resolution range)	96.5 (30.00-2.90)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.245 , 0.272	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	17368	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VAA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.47	1/1791 (0.1%)	0.78	0/2789
1	D	0.46	1/1791 (0.1%)	0.78	1/2789 (0.0%)
2	A	0.44	0/7143	0.69	3/9678 (0.0%)
2	B	0.43	0/7143	0.68	2/9678 (0.0%)
All	All	0.44	2/17868 (0.0%)	0.71	6/24934 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	901	G	OP3-P	-7.02	1.52	1.61
1	D	901	G	OP3-P	-6.91	1.52	1.61

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	172	LEU	CA-CB-CG	7.55	132.66	115.30
1	D	936	A	N9-C1'-C2'	6.97	123.06	114.00
2	B	382	VAL	C-N-CD	5.74	140.45	128.40
2	B	409	GLY	N-CA-C	5.50	126.84	113.10
2	A	409	GLY	N-CA-C	5.13	125.93	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1603	0	816	67	0
1	D	1603	0	816	66	0
2	A	6970	0	6943	728	0
2	B	6970	0	6942	751	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	30	0	24	3	0
4	B	30	0	24	4	0
5	A	67	0	0	13	0
5	B	61	0	0	17	0
5	C	19	0	0	0	0
5	D	11	0	0	1	0
All	All	17368	0	15565	1571	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 48.

The worst 5 of 1571 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:777:LYS:HE3	2:B:777:LYS:H	1.06	1.18
2:B:201:ARG:HE	2:B:211:GLU:HB3	1.02	1.17
2:A:102:ARG:HH21	2:A:104:ASP:HA	1.04	1.16
2:A:382:VAL:HG11	2:A:516:LEU:HD12	1.19	1.16
2:A:28:ASN:HD21	2:A:30:LYS:HG2	1.05	1.13

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	860/862 (100%)	676 (79%)	139 (16%)	45 (5%)	2	6
2	B	860/862 (100%)	652 (76%)	156 (18%)	52 (6%)	1	4
All	All	1720/1724 (100%)	1328 (77%)	295 (17%)	97 (6%)	2	5

5 of 97 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	94	LEU
2	A	103	HIS
2	A	104	ASP
2	A	234	ARG
2	A	312	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	724/724 (100%)	641 (88%)	83 (12%)	5	17
2	B	724/724 (100%)	630 (87%)	94 (13%)	4	12
All	All	1448/1448 (100%)	1271 (88%)	177 (12%)	5	15

5 of 177 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	270	LYS
2	B	479	PHE
2	B	284	GLU
2	B	404	ARG
2	B	545	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	396	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	584	ASN
2	B	401	ASN
2	B	518	HIS
2	B	692	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	C	74/75 (98%)	26 (35%)	10 (13%)
1	D	74/75 (98%)	27 (36%)	7 (9%)
All	All	148/150 (98%)	53 (35%)	17 (11%)

5 of 53 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	C	903	G
1	C	907	C
1	C	908	U
1	C	909	A
1	C	910	G

5 of 17 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	D	947	C
1	D	959	U
1	C	947	C
1	C	957	A
1	C	959	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	VAA	B	1990	-	27,32,32	3.05	9 (33%)	30,48,48	1.69	5 (16%)
4	VAA	A	990	-	27,32,32	3.18	10 (37%)	30,48,48	1.87	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	VAA	B	1990	-	-	1/19/39/39	0/3/3/3
4	VAA	A	990	-	-	5/19/39/39	0/3/3/3

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	990	VAA	C5'-N5'	-9.33	1.31	1.47
4	B	1990	VAA	C5'-N5'	-8.38	1.33	1.47
4	B	1990	VAA	S-N5'	7.13	1.69	1.61
4	A	990	VAA	S-N5'	6.44	1.69	1.61
4	A	990	VAA	O1S-S	6.27	1.52	1.43

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	990	VAA	O2S-S-O1S	-6.27	111.01	120.40
4	B	1990	VAA	O2S-S-O1S	-5.46	112.23	120.40
4	A	990	VAA	C-N3S-S	-3.83	117.87	124.23
4	B	1990	VAA	C-N3S-S	-3.05	119.18	124.23
4	B	1990	VAA	O1S-S-N5'	-2.84	101.73	106.73

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	990	VAA	C5'-N5'-S-O1S
4	A	990	VAA	C3'-C4'-C5'-N5'
4	A	990	VAA	O4'-C4'-C5'-N5'
4	A	990	VAA	C5'-N5'-S-O2S
4	A	990	VAA	C5'-N5'-S-N3S


There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1990	VAA	4	0
4	A	990	VAA	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.