

# wwPDB X-ray Structure Validation Summary Report (i)

#### Sep 7, 2023 – 09:25 AM EDT

PDB ID : 4GAE

Title: Crystal structure of plasmodium dxr in complex with a pyridine-containing

inhibitor

Authors : Diao, J.; Xue, J.; Cai, G.; Deng, L.; Song, Y.

Deposited on : 2012-07-25

Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	426	71%	25%	
1	В	426	9%	21%	



# 2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 6931 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 1-deoxy-D-xylulose 5-phosphate reductoisomerase, apicoplast.

$\mathbf{Mol}$	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	A	416	Total 3324	C 2132	N 549	O 623	S 20	0	0	0
1	В	410	Total 3276	C 2103	N 537	O 616	S 20	0	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	63	MET	-	expression tag	UNP O96693
A	64	ARG	-	expression tag	UNP O96693
A	65	GLY	-	expression tag	UNP O96693
A	66	SER	-	expression tag	UNP O96693
A	67	HIS	-	expression tag	UNP O96693
A	68	HIS	-	expression tag	UNP O96693
A	69	HIS	-	expression tag	UNP O96693
A	70	HIS	-	expression tag	UNP O96693
A	71	HIS	-	expression tag	UNP O96693
A	72	HIS	-	expression tag	UNP O96693
A	73	GLY	-	expression tag	UNP O96693
A	74	SER	-	expression tag	UNP O96693
A	247	ILE	LEU	engineered mutation	UNP O96693
В	63	MET	-	expression tag	UNP O96693
В	64	ARG	-	expression tag	UNP O96693
В	65	GLY	-	expression tag	UNP O96693
В	66	SER	-	expression tag	UNP O96693
В	67	HIS	-	expression tag	UNP O96693
В	68	HIS	-	expression tag	UNP O96693
В	69	HIS	-	expression tag	UNP O96693
В	70	HIS	-	expression tag	UNP O96693
В	71	HIS	-	expression tag	UNP O96693
В	72	HIS	-	expression tag	UNP O96693
В	73	GLY	-	expression tag	UNP O96693
В	74	SER	-	expression tag	UNP O96693


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	247	ILE	LEU	engineered mutation	UNP O96693


• Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).



Mol	Chain	Residues		$\mathbf{At}$	oms			ZeroOcc	AltConf
2	Λ	1	Total	С	N	О	Р	0	0
	A	1	48	21	7	17	3		U
9	D	1	Total	С	N	О	Р	0	0
	Б	1	48	21	7	17	3		


• Molecule 3 is [(1R)-3-[acetyl(hydroxy)amino]-1-(pyridin-4-yl)propyl]phosphonic acid (three-letter code: FOB) (formula:  $C_{10}H_{15}N_2O_5P$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
9	٨	1	Total	С	N	О	Р	0	1	
3	A	1	18	10	2	5	1		1	
2	D	1	Total	С	N	О	Р	0	1	
3	Б	1	18	10	2	5	1	U	1	

• Molecule 4 is [(1S)-3-[acetyl(hydroxy)amino]-1-(pyridin-4-yl)propyl]phosphonic acid (three-letter code: FOQ) (formula:  $C_{10}H_{15}N_2O_5P$ ).



$\mathbf{Mol}$	Chain	Residues		Ato	$\mathbf{ms}$			ZeroOcc	AltConf
4	A	1	Total 18	C 10	N 2	O 5	P 1	0	1

Continued on next page...



Continued from previous page...

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
1	D	1	Total	С	N	О	Р	0	1
4	Б	1	18	10	2	5	1	U	1


• Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mn 1 1	0	0
5	В	3	Total Mn 3 3	0	0

• Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0

• Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0
7	В	1	Total C O 4 2 2	0	0

• Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 4 2 2	0	0
8	В	1	Total C O 4 2 2	0	0
8	В	1	Total C O 4 2 2	0	0

#### • Molecule 9 is water.


$\mathbf{Mol}$	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	67	Total O 67 67	0	0
9	В	71	Total O 71 71	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase, apicoplast



• Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase, apicoplast





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	51.36Å 77.28Å 109.41Å	Donositon
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $91.50^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	44.64 - 2.30	Depositor
Resolution (A)	44.64 - 2.30	EDS
% Data completeness	89.0 (44.64-2.30)	Depositor
(in resolution range)	89.1 (44.64-2.30)	EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	5.53  (at  2.29Å)	Xtriage
Refinement program	CNS 1.3	Depositor
D.D.	0.230 , $0.273$	Depositor
$R, R_{free}$	0.210 , $0.258$	DCC
$R_{free}$ test set	1698 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.3	Xtriage
Anisotropy	0.170	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.31, 30.2	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	0.037 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	6931	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 11.67% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, EDO, MN, FOQ, FOB, NDP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles	
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.35	0/3389	0.60	0/4575
1	В	0.38	0/3339	0.61	0/4509
All	All	0.37	0/6728	0.60	0/9084

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3324	0	3369	88	0
1	В	3276	0	3322	67	0
2	A	48	0	26	2	0
2	В	48	0	26	0	0
3	A	18	0	12	1	0
3	В	18	0	12	0	0
4	A	18	0	12	0	0
4	В	18	0	12	0	0
5	A	1	0	0	0	0
5	В	3	0	0	0	0
6	A	1	0	0	1	0

Continued on next page...



Continued	tmom	mmonia	maaa
Canadana		THETHIS	THEFT

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	4	0	3	0	0
7	В	4	0	3	0	0
8	A	4	0	6	0	0
8	В	8	0	12	0	0
9	A	67	0	0	0	0
9	В	71	0	0	3	0
All	All	6931	0	6815	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:B:295:LYS:HD2	1:B:358:PRO:HG2	1.36	1.04
1:B:269:SER:HB2	1:B:312:LYS:HE3	1.50	0.92
1:A:156:PRO:HG2	1:A:158:ILE:HD11	1.65	0.79
1:B:295:LYS:CD	1:B:358:PRO:HG2	2.11	0.78
1:B:283:LYS:NZ	1:B:283:LYS:HB2	1.98	0.77

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	414/426 (97%)	385 (93%)	24 (6%)	5 (1%)	13	14
1	В	408/426 (96%)	386 (95%)	18 (4%)	4 (1%)	15	17
All	All	822/852 (96%)	771 (94%)	42 (5%)	9 (1%)	14	15

5 of 9 Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	В	294	PRO
1	A	342	SER
1	A	453	SER
1	A	105	ASN
1	В	342	SER

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	382/391 (98%)	371 (97%)	11 (3%)	42 58
1	В	377/391 (96%)	368 (98%)	9 (2%)	49 66
All	All	759/782~(97%)	739 (97%)	20 (3%)	46 63

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	165	MET
1	В	264	LYS
1	В	292	LYS
1	В	284	ASN
1	A	213	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	255	ASN
1	В	420	ASN
1	В	433	ASN
1	В	385	GLN
1	A	459	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 5 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	T	Chain	Das	T :1-	Во	ond leng	$_{ m ths}$	Bond angles		
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	FOQ	В	503[B]	5	16,18,18	2.28	7 (43%)	17,25,25	1.64	3 (17%)
3	FOB	A	502[A]	5	16,18,18	2.27	5 (31%)	17,25,25	2.63	5 (29%)
8	EDO	В	508	-	3,3,3	0.52	0	2,2,2	0.31	0
2	NDP	A	501	-	45,52,52	1.54	5 (11%)	53,80,80	1.39	4 (7%)
4	FOQ	A	503[B]	5	16,18,18	2.21	7 (43%)	17,25,25	1.79	5 (29%)
8	EDO	В	509	-	3,3,3	0.57	0	2,2,2	0.27	0
2	NDP	В	501	-	45,52,52	1.61	5 (11%)	53,80,80	1.44	5 (9%)
8	EDO	A	507	-	3,3,3	0.52	0	2,2,2	0.32	0
7	ACT	В	507	-	3,3,3	1.04	0	3,3,3	0.82	0
3	FOB	В	502[A]	5	16,18,18	2.25	6 (37%)	17,25,25	2.60	3 (17%)
7	ACT	A	506	-	3,3,3	1.04	0	3,3,3	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FOQ	В	503[B]	5	-	3/19/19/19	0/1/1/1
3	FOB	A	502[A]	5	-	4/19/19/19	0/1/1/1

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	EDO	В	508	-	-	1/1/1/1	-
2	NDP	A	501	-	-	4/30/77/77	0/5/5/5
4	FOQ	A	503[B]	5	-	4/19/19/19	0/1/1/1
8	EDO	В	509	-	-	0/1/1/1	-
2	NDP	В	501	-	-	4/30/77/77	0/5/5/5
8	EDO	A	507	-	-	0/1/1/1	-
3	FOB	В	502[A]	5	-	8/19/19/19	0/1/1/1

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
3	A	502[A]	FOB	PA1-C4	-6.38	1.73	1.82
2	В	501	NDP	O7N-C7N	6.01	1.38	1.24
4	В	503[B]	FOQ	PA1-C4	-5.87	1.73	1.82
3	В	502[A]	FOB	PA1-C4	-5.71	1.74	1.82
4	A	503[B]	FOQ	PA1-C4	-5.65	1.74	1.82

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\mathbf{Ideal}(^o)$
3	В	502[A]	FOB	C3-C2-N1	-9.37	98.19	111.36
3	A	502[A]	FOB	C3-C2-N1	-9.03	98.67	111.36
2	В	501	NDP	N3A-C2A-N1A	-6.60	118.37	128.68
2	A	501	NDP	N3A-C2A-N1A	-6.59	118.38	128.68
4	A	503[B]	FOQ	C3-C2-N1	-4.28	105.35	111.36

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	502[A]	FOB	N1-C2-C3-C4
8	В	508	EDO	O1-C1-C2-O2
2	A	501	NDP	O4D-C1D-N1N-C6N
3	A	502[A]	FOB	N1-C2-C3-C4
4	A	503[B]	FOQ	C3-C2-N1-C1


There are no ring outliers.

2 monomers are involved in 3 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502[A]	FOB	1	0
2	A	501	NDP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























# 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	$416/426 \ (97\%)$	0.69	57 (13%) 3 4	21, 36, 71, 104	1 (0%)
1	В	410/426 (96%)	0.44	37 (9%) 9 12	18, 30, 54, 98	0
All	All	826/852 (96%)	0.57	94 (11%) 5 7	18, 33, 65, 104	1 (0%)

The worst 5 of 94 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	294	PRO	16.9
1	В	291	LEU	11.2
1	A	292	LYS	10.4
1	A	295	LYS	9.4
1	A	132	TYR	9.4

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathrm{A}}^2)$	Q<0.9
7	ACT	A	506	4/4	0.74	0.32	93,93,94,94	0
8	EDO	A	507	4/4	0.80	0.35	78,79,80,80	0
8	EDO	В	508	4/4	0.81	0.16	64,65,65,65	0
8	EDO	В	509	4/4	0.81	0.37	78,79,79,79	0
7	ACT	В	507	4/4	0.90	0.28	65,67,67,67	0
4	FOQ	A	503[B]	18/18	0.90	0.19	26,33,36,37	18
3	FOB	A	502[A]	18/18	0.91	0.20	28,35,38,39	18
2	NDP	A	501	48/48	0.92	0.12	30,38,42,42	0
5	MN	В	506	1/1	0.93	0.04	49,49,49,49	0
3	FOB	В	502[A]	18/18	0.94	0.16	26,30,36,36	18
4	FOQ	В	503[B]	18/18	0.94	0.15	20,24,31,31	18
2	NDP	В	501	48/48	0.95	0.10	22,28,30,32	0
5	MN	В	505	1/1	0.98	0.05	40,40,40,40	0
6	CL	A	505	1/1	0.99	0.06	45,45,45,45	0
5	MN	В	504	1/1	0.99	0.04	24,24,24,24	0
5	MN	A	504	1/1	1.00	0.14	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















# Electron density around FOB B 502 (A): 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)



# Electron density around FOQ B 503 (B): $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





# 6.5 Other polymers (i)

There are no such residues in this entry.

