



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 10:28 PM BST

PDB ID : 6GFE
Title : High-resolution Structure of a therapeutic full-length anti-NPRA Antibody with exceptional Conformational Diversity
Authors : Hoerer, S.
Deposited on : 2018-04-30
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	450	<p>10% (red), 90% (green), 6% (yellow), 5% (grey)</p>
1	K	450	<p>4% (red), 91% (green), 5% (yellow), 0% (grey)</p>
2	L	215	<p>0% (red), 97% (green), 0% (yellow), 0% (grey)</p>
2	M	215	<p>6% (red), 98% (green), 0% (yellow), 0% (grey)</p>
3	A	7	<p>0% (red), 71% (green), 29% (yellow), 0% (grey)</p>
3	B	7	<p>0% (red), 43% (green), 57% (yellow), 0% (grey)</p>

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 11641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Immunoglobulin gamma-4 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	429	Total 3405	C 2148	N 574	O 666	S 17	6	7	0
1	K	426	Total 3400	C 2143	N 573	O 665	S 19	6	10	0

- Molecule 2 is a protein called Immunoglobulin gamma-4 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	215	Total 1703	C 1056	N 293	O 347	S 7	0	9	0
2	M	215	Total 1662	C 1035	N 285	O 335	S 7	6	3	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	A	7	Total 85	C 48	N 3	O 34	0	0	0
3	B	7	Total 85	C 48	N 3	O 34	0	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total Ca 1 1	0	0
4	K	1	Total Ca 1 1	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	K	1	Total C O 4 2 2	0	0
5	L	1	Total C O 4 2 2	0	0
5	L	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	372	Total O 372 372	0	1
6	K	493	Total O 494 494	0	2
6	L	249	Total O 249 249	0	0
6	M	172	Total O 172 172	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Immunoglobulin gamma-4 heavy chain




- Molecule 1: Immunoglobulin gamma-4 heavy chain



- Molecule 2: Immunoglobulin gamma-4 light chain



- Molecule 2: Immunoglobulin gamma-4 light chain



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyra


nose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain A:  71% 29%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  43% 57%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.78Å 160.51Å 87.30Å 90.00° 110.72° 90.00°	Depositor
Resolution (Å)	40.00 – 1.80 39.57 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (40.00-1.80) 99.5 (39.57-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.79Å)	Xtrriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.181 , 0.200 0.189 , 0.210	Depositor DCC
R_{free} test set	7796 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtrriage
Anisotropy	0.297	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11641	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CA, FUC, ACT, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.41	0/3496	0.61	0/4762
1	K	0.42	0/3488	0.61	0/4754
2	L	0.41	0/1738	0.61	0/2362
2	M	0.40	0/1701	0.62	0/2311
All	All	0.41	0/10423	0.61	0/14189

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	3405	0	3303	7	0
1	K	3400	0	3285	5	0
2	L	1703	0	1648	2	0
2	M	1662	0	1612	3	0
3	A	85	0	73	0	0
3	B	85	0	73	0	0
4	H	1	0	0	0	0
4	K	1	0	0	0	0
5	K	4	0	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	8	0	6	0	0
6	H	372	0	0	0	0
6	K	494	0	0	0	0
6	L	249	0	0	0	0
6	M	172	0	0	0	0
All	All	11641	0	10003	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:346:PRO:HB3	1:H:372:PHE:HB3	1.78	0.66
1:H:278:TYR:HB2	1:H:320:LYS:HB3	1.85	0.59
1:K:127:PRO:HD3	1:K:213:LYS:HE2	1.85	0.58
1:H:3:GLU:HG3	1:H:25:SER:HB2	1.87	0.56
1:H:314:LEU:HD23	1:H:338:LYS:HE3	1.89	0.54
2:M:145:LYS:HB3	2:M:197:THR:HB	1.89	0.53
2:M:198[A]:HIS:CD2	2:M:200:GLY:H	2.26	0.53
2:L:145:LYS:HB3	2:L:197:THR:HB	1.90	0.53
1:K:346:PRO:HB3	1:K:372:PHE:HB3	1.94	0.48
1:H:35:SER:HB2	1:H:95:MET:HB3	1.97	0.47
1:K:23:ALA:HA	1:K:77:GLN:HG2	1.97	0.47
1:H:290:LYS:HB2	1:H:303:VAL:HG13	2.00	0.44
2:M:141:PRO:O	2:M:198[A]:HIS:HE1	2.00	0.44
2:L:28:VAL:HG13	2:L:92:SER:HB2	1.99	0.44
1:K:11:LEU:HB2	1:K:151:PRO:HG3	2.01	0.43
1:H:263:VAL:HG12	1:H:302:VAL:HB	2.02	0.41
1:K:343:PRO:HA	1:K:373:TYR:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	430/450 (96%)	423 (98%)	4 (1%)	3 (1%)	22	10
1	K	430/450 (96%)	425 (99%)	4 (1%)	1 (0%)	47	33
2	L	222/215 (103%)	217 (98%)	5 (2%)	0	100	100
2	M	216/215 (100%)	211 (98%)	5 (2%)	0	100	100
All	All	1298/1330 (98%)	1276 (98%)	18 (1%)	4 (0%)	41	27

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	97	VAL
1	H	272	GLU
1	K	97	VAL
1	H	271	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	398/408 (98%)	389 (98%)	9 (2%)	50	37
1	K	398/408 (98%)	394 (99%)	4 (1%)	76	71
2	L	195/186 (105%)	193 (99%)	2 (1%)	76	71
2	M	189/186 (102%)	189 (100%)	0	100	100
All	All	1180/1188 (99%)	1165 (99%)	15 (1%)	71	62

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	35(B)	SER
1	H	132	SER
1	H	200[A]	CYS
1	H	200[B]	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	201	ASN
1	H	285	HIS
1	H	326	LYS
1	H	328	LEU
1	H	399	ASP
1	K	132	SER
1	K	201	ASN
1	K	384	ASN
1	K	399	ASP
2	L	45	ARG
2	L	91	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	93	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1	1,3	14,14,15	0.31	0	17,19,21	0.62	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	2	3	14,14,15	0.32	0	17,19,21	0.73	1 (5%)
3	BMA	A	3	3	11,11,12	0.31	0	15,15,17	0.60	0
3	MAN	A	4	3	11,11,12	0.31	0	15,15,17	0.83	1 (6%)
3	NAG	A	5	3	14,14,15	0.30	0	17,19,21	0.44	0
3	MAN	A	6	3	11,11,12	0.25	0	15,15,17	0.67	0
3	FUC	A	7	3	10,10,11	0.53	0	14,14,16	0.68	0
3	NAG	B	1	1,3	14,14,15	0.24	0	17,19,21	0.67	1 (5%)
3	NAG	B	2	3	14,14,15	0.29	0	17,19,21	0.58	0
3	BMA	B	3	3	11,11,12	0.29	0	15,15,17	0.60	0
3	MAN	B	4	3	11,11,12	0.32	0	15,15,17	0.92	1 (6%)
3	NAG	B	5	3	14,14,15	0.28	0	17,19,21	0.43	0
3	MAN	B	6	3	11,11,12	0.39	0	15,15,17	0.79	1 (6%)
3	FUC	B	7	3	10,10,11	0.41	0	14,14,16	0.77	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	2	3	-	0/6/23/26	0/1/1/1
3	BMA	A	3	3	-	0/2/19/22	0/1/1/1
3	MAN	A	4	3	-	0/2/19/22	0/1/1/1
3	NAG	A	5	3	-	0/6/23/26	0/1/1/1
3	MAN	A	6	3	-	0/2/19/22	0/1/1/1
3	FUC	A	7	3	-	-	0/1/1/1
3	NAG	B	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	2	3	-	0/6/23/26	0/1/1/1
3	BMA	B	3	3	-	0/2/19/22	0/1/1/1
3	MAN	B	4	3	-	0/2/19/22	0/1/1/1
3	NAG	B	5	3	-	0/6/23/26	0/1/1/1
3	MAN	B	6	3	-	0/2/19/22	0/1/1/1
3	FUC	B	7	3	-	-	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	4	MAN	C1-O5-C5	3.09	116.38	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	4	MAN	C1-O5-C5	2.99	116.24	112.19
3	B	7	FUC	C1-O5-C5	2.30	118.00	112.78
3	B	6	MAN	C1-O5-C5	2.20	115.17	112.19
3	B	1	NAG	C1-O5-C5	2.11	115.06	112.19
3	A	2	NAG	O5-C1-C2	-2.09	107.99	111.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACT	L	302	-	1,3,3	4.30	1 (100%)	0,3,3	0.00	-
5	ACT	L	301	-	1,3,3	2.10	1 (100%)	0,3,3	0.00	-
5	ACT	K	509	-	1,3,3	3.96	1 (100%)	0,3,3	0.00	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	L	302	ACT	CH3-C	4.30	1.54	1.48
5	K	509	ACT	CH3-C	3.96	1.53	1.48
5	L	301	ACT	CH3-C	2.10	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	429/450 (95%)	0.31	47 (10%) 5 4	16, 34, 87, 103	0
1	K	426/450 (94%)	-0.32	18 (4%) 36 30	17, 27, 54, 102	2 (0%)
2	L	215/215 (100%)	-0.39	2 (0%) 84 82	17, 27, 51, 71	0
2	M	215/215 (100%)	-0.04	13 (6%) 21 17	18, 32, 70, 79	0
All	All	1285/1330 (96%)	-0.07	80 (6%) 20 16	16, 29, 68, 103	2 (0%)

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	328	LEU	9.6
1	H	330	SER	7.8
1	H	329	PRO	6.6
1	H	327	GLY	6.4
1	H	277	TRP	6.1
1	H	332	ILE	6.1
2	L	214	CYS	5.9
1	K	444	SER	5.7
1	H	323	VAL	5.6
1	H	136	SER	5.4
2	L	212	GLY	5.4
1	H	275	PHE	5.1
1	K	217	SER	5.0
1	H	273	VAL	4.9
1	K	296	PHE	4.8
1	H	284	VAL	4.8
1	K	195	THR	4.7
1	H	326	LYS	4.7
1	K	329	PRO	4.7
1	H	269	GLU	4.5
1	H	274	GLN	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	276	ASN	4.4
1	K	194	GLY	4.4
1	K	328	LEU	4.3
2	M	154	LEU	4.1
1	H	285	HIS	4.1
1	H	444	SER	3.9
1	H	271	PRO	3.8
1	H	282	VAL	3.8
1	H	331	SER	3.6
1	H	253	ILE	3.5
1	H	325	ASN	3.5
2	M	214	CYS	3.5
1	H	289	THR	3.3
1	H	321	CYS	3.2
1	K	192	SER	3.1
1	H	218	LYS	3.1
1	K	327	GLY	3.1
1	H	278	TYR	3.1
2	M	127	SER	3.1
1	H	303	VAL	3.1
1	K	138	SER	3.0
1	H	272	GLU	2.9
1	K	191	SER	2.9
2	M	212	GLY	2.8
1	H	324	SER	2.7
1	H	270	ASP	2.7
2	M	150	VAL	2.7
2	M	188	LYS	2.6
1	H	268	GLN	2.6
1	H	137	GLU	2.5
1	H	256	THR	2.5
2	M	184	ALA	2.5
2	M	190	LYS	2.5
2	M	126	LYS	2.4
1	H	308	VAL	2.4
1	H	322	LYS	2.4
1	H	259	VAL	2.4
2	M	152	ASN	2.4
1	K	1	GLN	2.4
1	H	333	GLU	2.3
1	K	189	PRO	2.3
1	K	132	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	K	326	LYS	2.3
1	H	291	PRO	2.3
1	K	193	LEU	2.3
1	H	263	VAL	2.2
1	H	288	LYS	2.2
2	M	125	LEU	2.2
1	K	385	GLY	2.2
1	H	280	ASP	2.2
1	H	340	LYS	2.1
1	H	54	HIS	2.1
1	H	283	GLU	2.1
1	H	319	TYR	2.0
1	K	384	ASN	2.0
2	M	210	ASN	2.0
1	H	252	MET	2.0
2	M	189	HIS	2.0
1	H	336	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MAN	A	6	11/12	0.80	0.14	40,45,49,51	0
3	FUC	B	7	10/11	0.80	0.24	62,68,69,70	0
3	MAN	B	6	11/12	0.90	0.10	40,43,50,53	0
3	MAN	B	4	11/12	0.92	0.08	38,40,45,46	0
3	NAG	B	5	14/15	0.92	0.07	39,42,44,44	0
3	NAG	A	5	14/15	0.92	0.10	35,38,40,42	0
3	NAG	B	1	14/15	0.93	0.08	31,37,46,55	0
3	BMA	B	3	11/12	0.93	0.07	28,33,37,40	0
3	FUC	A	7	10/11	0.95	0.06	23,24,25,27	0
3	NAG	A	1	14/15	0.95	0.07	26,32,40,40	0
3	MAN	A	4	11/12	0.95	0.06	29,30,32,33	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	A	2	14/15	0.95	0.07	25,31,38,42	0
3	NAG	B	2	14/15	0.96	0.06	27,31,34,36	0
3	BMA	A	3	11/12	0.96	0.06	25,28,31,34	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ACT	K	509	4/4	0.76	0.19	39,40,43,48	0
5	ACT	L	301	4/4	0.89	0.13	44,49,50,51	0
5	ACT	L	302	4/4	0.90	0.16	40,46,47,47	0
4	CA	H	508	1/1	0.92	0.10	75,75,75,75	0
4	CA	K	508	1/1	0.99	0.04	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.