



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 05:22 AM BST

PDB ID : 1GLM  
Title : REFINED CRYSTAL STRUCTURES OF GLUCOAMYLASE FROM AS-  
PERGILLUS AWAMORI VAR. X100  
Authors : Aleshin, A.E.; Hoffman, C.; Firsov, L.M.; Honzatko, R.B.  
Deposited on : 1994-04-25  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.




Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	470	 88% 12%
2	B	5	 100%
3	C	8	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	B	3	X	-	-	-
3	MAN	C	3	X	-	-	-

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4404 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUCOAMYLASE-471.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	470	3578	2242	585	743	8	0	5	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	58	LEU	ILE	conflict	UNP P69327
A	60	ILE	LEU	conflict	UNP P69327
A	117	THR	ALA	conflict	UNP P69327

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	5	61	34	2	25	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	8	94	52	2	40	0	0	0

- Molecule 4 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0
4	A	1	11	6	5	0	0

- Molecule 5 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	561	Total 561	O 561	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: GLUCOAMYLASE-471



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.80Å 104.40Å 48.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.40	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.40)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.122 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4404	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.85	0/3692	1.35	35/5051 (0.7%)

There are no bond length outliers.

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	428	ARG	NE-CZ-NH1	14.66	127.63	120.30
1	A	273	ARG	NE-CZ-NH1	11.37	125.98	120.30
1	A	68	ARG	NE-CZ-NH2	-11.26	114.67	120.30
1	A	273	ARG	NE-CZ-NH2	-11.02	114.79	120.30
1	A	428	ARG	NE-CZ-NH2	-9.53	115.53	120.30
1	A	97[A]	ASP	CB-CG-OD1	-8.32	110.81	118.30
1	A	97[B]	ASP	CB-CG-OD1	-8.32	110.81	118.30
1	A	194	ARG	CD-NE-CZ	7.80	134.53	123.60
1	A	341	LEU	CA-CB-CG	7.72	133.05	115.30
1	A	289	TYR	CB-CG-CD1	7.50	125.50	121.00
1	A	194	ARG	NE-CZ-NH2	-7.24	116.68	120.30
1	A	305	ARG	NE-CZ-NH2	-7.21	116.69	120.30
1	A	297	ASP	CB-CG-OD1	-6.92	112.07	118.30
1	A	429	ARG	NE-CZ-NH1	6.68	123.64	120.30
1	A	160	ARG	NE-CZ-NH1	6.54	123.57	120.30
1	A	289	TYR	CB-CG-CD2	-6.43	117.14	121.00
1	A	198	GLU	CA-CB-CG	6.32	127.31	113.40
1	A	293	ASP	CB-CG-OD1	6.25	123.93	118.30
1	A	414	ASP	CB-CG-OD2	6.12	123.80	118.30
1	A	48	TYR	CB-CG-CD1	6.11	124.67	121.00
1	A	454	ALA	N-CA-CB	6.05	118.56	110.10
1	A	329	TYR	CB-CG-CD1	5.94	124.57	121.00
1	A	241	ARG	NE-CZ-NH2	-5.87	117.37	120.30
1	A	329	TYR	CB-CG-CD2	-5.83	117.50	121.00

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	413	ARG	NE-CZ-NH1	5.67	123.14	120.30
1	A	241	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	A	305	ARG	NE-CZ-NH1	5.59	123.09	120.30
1	A	413	ARG	NE-CZ-NH2	-5.45	117.58	120.30
1	A	122	ARG	NE-CZ-NH2	5.39	123.00	120.30
1	A	429	ARG	CD-NE-CZ	5.27	130.97	123.60
1	A	406	ASP	CB-CG-OD1	5.22	123.00	118.30
1	A	160	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	264	ASP	CB-CG-OD1	5.17	122.95	118.30
1	A	68	ARG	NE-CZ-NH1	5.15	122.88	120.30
1	A	194	ARG	NE-CZ-NH1	5.12	122.86	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3578	0	3317	19	0
2	B	61	0	52	0	0
3	C	94	0	79	0	0
4	A	110	0	100	1	0
5	A	561	0	0	3	0
All	All	4404	0	3548	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:371:SER:HB2	5:A:1103:HOH:O	1.98	0.62
1:A:217:ALA:HB3	1:A:218:PRO:HD3	1.81	0.62
1:A:462:THR:HB	4:A:494(A):MAN:H62	1.83	0.60
1:A:21:ASN:ND2	1:A:28:TRP:HE3	2.04	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ASN:HD21	1:A:28:TRP:HE3	1.54	0.54
1:A:137:GLY:HA3	5:A:1048:HOH:O	2.07	0.54
1:A:3:LEU:HD11	1:A:385:VAL:HG12	1.89	0.54
1:A:312:TYR:O	1:A:313:ASN:HB2	2.09	0.53
1:A:60:ILE:O	1:A:64:VAL:HG23	2.12	0.50
1:A:86:ALA:HA	1:A:158:LEU:HD21	1.96	0.47
1:A:316:PRO:HD2	1:A:402:PHE:O	2.14	0.47
1:A:60:ILE:HG13	1:A:81:TYR:CD1	2.50	0.46
1:A:325:ALA:HB2	1:A:384:PHE:HB2	1.98	0.46
1:A:106:GLU:HA	1:A:107:PRO:HD3	1.84	0.45
1:A:403:ASP:HB3	1:A:406:ASP:OD1	2.18	0.44
1:A:341:LEU:HD11	1:A:373:ILE:HG21	2.02	0.42
1:A:433:VAL:HB	1:A:434:PRO:HD2	2.02	0.42
1:A:268:GLN:HA	5:A:781:HOH:O	2.21	0.41
1:A:128:PRO:HG2	1:A:165:TYR:CD1	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	473/470 (101%)	453 (96%)	20 (4%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	390/385 (101%)	381 (98%)	9 (2%)	50 70

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55[A]	ASP
1	A	55[B]	ASP
1	A	247	ASN
1	A	310	SER
1	A	341	LEU
1	A	356	SER
1	A	394	SER
1	A	399	SER
1	A	465	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	45	ASN
1	A	168	GLN
1	A	182	ASN
1	A	426	ASN
1	A	427	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	1.15	1 (7%)	17,19,21	1.33	2 (11%)
2	NAG	B	2	2	14,14,15	0.78	1 (7%)	17,19,21	1.46	3 (17%)
2	MAN	B	3	2	11,11,12	0.49	0	15,15,17	1.48	1 (6%)
2	MAN	B	4	2	11,11,12	0.74	0	15,15,17	1.61	2 (13%)
2	MAN	B	5	2	11,11,12	1.05	1 (9%)	15,15,17	1.86	4 (26%)
3	NAG	C	1	1,3	14,14,15	1.03	1 (7%)	17,19,21	1.37	2 (11%)
3	NAG	C	2	3	14,14,15	0.89	1 (7%)	17,19,21	0.98	1 (5%)
3	MAN	C	3	3	11,11,12	0.70	0	15,15,17	1.54	3 (20%)
3	MAN	C	4	3	11,11,12	0.41	0	15,15,17	1.44	1 (6%)
3	MAN	C	5	3	11,11,12	0.57	0	15,15,17	1.29	2 (13%)
3	MAN	C	6	3	11,11,12	0.55	0	15,15,17	0.96	1 (6%)
3	MAN	C	7	3	11,11,12	0.75	0	15,15,17	1.21	2 (13%)
3	MAN	C	8	3	11,11,12	0.51	0	15,15,17	1.16	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	MAN	B	3	2	1/1/4/5	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	2/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	MAN	C	3	3	1/1/4/5	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
3	MAN	C	6	3	-	1/2/19/22	0/1/1/1
3	MAN	C	7	3	-	0/2/19/22	0/1/1/1
3	MAN	C	8	3	-	0/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	C8-C7	3.69	1.58	1.50
3	C	1	NAG	C8-C7	3.21	1.57	1.50
3	C	2	NAG	C8-C7	2.80	1.56	1.50
2	B	2	NAG	C8-C7	2.31	1.55	1.50
2	B	5	MAN	O5-C5	2.21	1.47	1.43

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	MAN	O5-C1-C2	-4.83	103.32	110.77
3	C	4	MAN	C1-O5-C5	4.79	118.68	112.19
2	B	3	MAN	C1-O5-C5	4.70	118.56	112.19
3	C	3	MAN	C1-O5-C5	3.53	116.97	112.19
2	B	5	MAN	C6-C5-C4	3.50	121.20	113.00
3	C	5	MAN	C1-O5-C5	3.40	116.80	112.19
2	B	1	NAG	C1-O5-C5	3.40	116.79	112.19
3	C	6	MAN	C1-O5-C5	3.20	116.52	112.19
3	C	1	NAG	C2-N2-C7	3.17	127.41	122.90
2	B	5	MAN	C3-C4-C5	-3.16	104.60	110.24
3	C	3	MAN	O5-C1-C2	-3.11	105.97	110.77
3	C	5	MAN	O5-C1-C2	-3.02	106.12	110.77
2	B	5	MAN	O5-C5-C4	-2.99	103.55	110.83
2	B	2	NAG	C1-O5-C5	2.84	116.04	112.19
3	C	7	MAN	O6-C6-C5	2.56	120.06	111.29
2	B	5	MAN	C1-C2-C3	2.53	112.77	109.67
3	C	8	MAN	O5-C1-C2	2.48	114.59	110.77
3	C	2	NAG	O4-C4-C5	-2.41	103.31	109.30
3	C	3	MAN	O2-C2-C1	2.38	114.02	109.15
3	C	1	NAG	C1-O5-C5	2.25	115.25	112.19
2	B	1	NAG	O4-C4-C5	-2.17	103.92	109.30
2	B	4	MAN	C6-C5-C4	2.15	118.05	113.00
3	C	7	MAN	C6-C5-C4	2.07	117.86	113.00
2	B	2	NAG	C2-N2-C7	-2.07	119.96	122.90
2	B	2	NAG	O5-C1-C2	-2.03	108.09	111.29

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	C	3	MAN	C1
2	B	3	MAN	C1


All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	5	MAN	O5-C5-C6-O6
2	B	5	MAN	C4-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6
3	C	4	MAN	C4-C5-C6-O6
3	C	6	MAN	O5-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MAN	A	487(A)	1	11,11,12	0.93	0	15,15,17	1.54	4 (26%)
4	MAN	A	493(A)	1	11,11,12	0.60	0	15,15,17	1.42	1 (6%)
4	MAN	A	494(A)	1	11,11,12	1.07	1 (9%)	15,15,17	1.00	1 (6%)
4	MAN	A	488(A)	1	11,11,12	0.90	1 (9%)	15,15,17	1.46	2 (13%)
4	MAN	A	492(A)	1	11,11,12	0.56	0	15,15,17	1.12	2 (13%)
4	MAN	A	485(A)	1	11,11,12	0.75	0	15,15,17	0.78	0
4	MAN	A	491(A)	1	11,11,12	0.54	0	15,15,17	1.16	2 (13%)
4	MAN	A	489(A)	1	11,11,12	0.59	0	15,15,17	0.76	0
4	MAN	A	486(A)	1	11,11,12	0.87	1 (9%)	15,15,17	1.31	3 (20%)
4	MAN	A	490(A)	1	11,11,12	0.90	1 (9%)	15,15,17	1.17	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	A	487(A)	1	-	2/2/19/22	0/1/1/1
4	MAN	A	493(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	494(A)	1	-	2/2/19/22	0/1/1/1
4	MAN	A	488(A)	1	-	2/2/19/22	0/1/1/1
4	MAN	A	492(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	485(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	491(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	489(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	486(A)	1	-	2/2/19/22	0/1/1/1
4	MAN	A	490(A)	1	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	494(A)	MAN	O5-C1	3.05	1.48	1.43
4	A	490(A)	MAN	O5-C1	2.31	1.47	1.43
4	A	486(A)	MAN	O5-C5	2.28	1.48	1.43
4	A	488(A)	MAN	O5-C5	2.01	1.47	1.43

All (16) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	493(A)	MAN	C1-O5-C5	4.59	118.41	112.19
4	A	494(A)	MAN	C1-O5-C5	3.04	116.31	112.19
4	A	492(A)	MAN	C1-O5-C5	2.99	116.25	112.19
4	A	491(A)	MAN	C1-O5-C5	2.92	116.15	112.19
4	A	487(A)	MAN	O5-C5-C4	-2.78	104.08	110.83
4	A	488(A)	MAN	C6-C5-C4	2.69	119.32	113.00
4	A	487(A)	MAN	C1-O5-C5	2.60	115.71	112.19
4	A	486(A)	MAN	O5-C5-C4	-2.55	104.63	110.83
4	A	486(A)	MAN	C3-C4-C5	-2.45	105.88	110.24
4	A	492(A)	MAN	O3-C3-C4	-2.44	104.71	110.35
4	A	486(A)	MAN	C6-C5-C4	2.44	118.71	113.00
4	A	490(A)	MAN	C1-O5-C5	2.34	115.37	112.19
4	A	488(A)	MAN	O5-C5-C6	2.31	110.82	107.20
4	A	487(A)	MAN	C3-C4-C5	-2.22	106.28	110.24
4	A	487(A)	MAN	O5-C1-C2	-2.21	107.36	110.77
4	A	491(A)	MAN	O3-C3-C4	-2.07	105.56	110.35

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	487(A)	MAN	O5-C5-C6-O6
4	A	486(A)	MAN	C4-C5-C6-O6
4	A	488(A)	MAN	C4-C5-C6-O6
4	A	487(A)	MAN	C4-C5-C6-O6
4	A	494(A)	MAN	O5-C5-C6-O6
4	A	494(A)	MAN	C4-C5-C6-O6
4	A	486(A)	MAN	O5-C5-C6-O6
4	A	488(A)	MAN	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	494(A)	MAN	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.