



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 21, 2025 – 03:03 pm GMT

PDB ID : 9GO1
Title : C1C2 Channelrhodopsin - SMX Dark structure
Authors : Mulder, M.; Weinert, T.; Skopintsev, P.; Bruenle, S.; Broser, M.; Hegemann, P.; Standfuss, J.
Deposited on : 2024-09-04
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	 4% 71% 17% 12%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Archaeal-type opsin 1, Archaeal-type opsin 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	294	2301	1509	367	410	15	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	349	SER	-	expression tag	UNP Q8RUT8
A	350	SER	-	expression tag	UNP Q8RUT8
A	351	GLU	-	expression tag	UNP Q8RUT8
A	352	ASP	-	expression tag	UNP Q8RUT8
A	353	LEU	-	expression tag	UNP Q8RUT8
A	354	TYR	-	expression tag	UNP Q8RUT8
A	355	PHE	-	expression tag	UNP Q8RUT8
A	356	GLN	-	expression tag	UNP Q8RUT8

- Molecule 2 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: C₂₁H₄₀O₄).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	C O	0	0
			25	21 4		
2	A	1	Total	C O	0	0
			16	14 2		
2	A	1	Total	C O	0	0
			14	10 4		
2	A	1	Total	C O	0	0
			16	12 4		
2	A	1	Total	C O	0	0
			18	14 4		
2	A	1	Total	C O	0	0
			10	8 2		
2	A	1	Total	C O	0	0
			10	8 2		
2	A	1	Total	C	0	0
			9	9		
2	A	1	Total	C	0	0
			8	8		

- Molecule 3 is RETINAL (three-letter code: RET) (formula: C₂₀H₂₈O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C	0	0
			20	20		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	13	Total	O	0	0
			13	13		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	61.40Å 141.70Å 94.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.85 – 2.59 70.85 – 2.59	Depositor EDS
% Data completeness (in resolution range)	66.9 (70.85-2.59) 66.9 (70.85-2.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.83 (at 2.58Å)	Xtrriage
Refinement program	PHENIX 1.20_4459	Depositor
R, R_{free}	0.211 , 0.270 0.211 , 0.268	Depositor DCC
R_{free} test set	12310 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	57.8	Xtrriage
Anisotropy	0.072	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2460	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RET, OLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2361	0.57	0/3215

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2301	0	2239	33	0
2	A	126	0	172	3	0
3	A	20	0	27	4	0
4	A	13	0	0	0	0
All	All	2460	0	2438	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:VAL:HG21	2:A:405:OLC:H2	1.79	0.65
1:A:244:LYS:HA	1:A:248:ARG:HH11	1.61	0.64
1:A:305:TYR:O	1:A:308:VAL:HG22	2.01	0.60
1:A:99:PHE:HB2	1:A:133:PHE:CD1	2.36	0.60
1:A:136:GLU:O	1:A:140:GLU:HB2	2.03	0.58
1:A:171:LEU:HD13	1:A:191:LEU:HG	1.86	0.56
1:A:324:THR:HA	1:A:348:VAL:O	2.06	0.56
1:A:254:MET:HE1	1:A:303:GLY:HA2	1.89	0.55
1:A:267:ILE:O	1:A:271:LEU:HD13	2.07	0.55
1:A:139:HIS:HB3	1:A:142:ASP:HB2	1.90	0.54
1:A:325:LYS:HE3	1:A:349:SER:HB3	1.92	0.52
1:A:304:HIS:O	1:A:308:VAL:HG13	2.09	0.52
1:A:53:GLN:HE21	1:A:55:SER:H	1.59	0.51
1:A:244:LYS:HA	1:A:248:ARG:NH1	2.25	0.50
1:A:351:GLU:N	1:A:351:GLU:OE2	2.44	0.49
1:A:119:GLY:O	1:A:123:ILE:HD13	2.12	0.49
1:A:283:GLY:O	1:A:286:VAL:HG22	2.12	0.48
1:A:246:ARG:HD2	1:A:309:LEU:HD21	1.97	0.47
1:A:266:PRO:HA	3:A:410:RET:H172	1.98	0.46
1:A:109:TYR:CE2	1:A:122:GLU:HB3	2.50	0.46
1:A:224:GLY:O	1:A:228:PHE:HD2	2.00	0.45
1:A:163:TRP:CD1	3:A:410:RET:H14	2.51	0.45
3:A:410:RET:H8	3:A:410:RET:H171	2.00	0.44
1:A:252:THR:HG21	2:A:409:OLC:H7	1.99	0.44
3:A:410:RET:H181	3:A:410:RET:H7	1.79	0.43
1:A:239:TYR:CZ	1:A:248:ARG:HD3	2.54	0.43
1:A:215:ILE:O	1:A:219:MET:HG3	2.19	0.43
1:A:134:ILE:HG13	1:A:135:ILE:N	2.33	0.42
2:A:403:OLC:H3	2:A:403:OLC:H6	1.75	0.42
1:A:157:TRP:CE2	1:A:161:ALA:HB2	2.54	0.42
1:A:240:HIS:HA	1:A:248:ARG:NH2	2.35	0.42
1:A:240:HIS:HA	1:A:248:ARG:HH21	1.85	0.41
1:A:106:LEU:HD12	1:A:130:MET:HE1	2.03	0.41
1:A:239:TYR:HD2	1:A:240:HIS:ND1	2.19	0.41
1:A:325:LYS:HE3	1:A:349:SER:CB	2.50	0.41
1:A:175:SER:OG	1:A:188:THR:HG23	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	288/333 (86%)	281 (98%)	7 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	240/282 (85%)	231 (96%)	9 (4%)	28 54

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	102	SER
1	A	106	LEU
1	A	144	PRO
1	A	162	GLU
1	A	173	HIS
1	A	321	ARG
1	A	322	LYS
1	A	325	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	GLN
1	A	85	ASN
1	A	95	GLN
1	A	182	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OLC	A	404	-	15,15,24	1.19	1 (6%)	16,16,25	1.08	1 (6%)
2	OLC	A	408	-	8,8,24	0.91	0	7,7,25	1.62	1 (14%)
2	OLC	A	407	-	9,9,24	1.45	1 (11%)	9,9,25	1.20	0
2	OLC	A	406	-	9,9,24	1.47	1 (11%)	9,9,25	1.20	0
3	RET	A	410	1	20,20,21	0.74	1 (5%)	27,27,28	0.59	1 (3%)
2	OLC	A	401	-	24,24,24	1.05	1 (4%)	25,25,25	1.24	2 (8%)
2	OLC	A	405	-	17,17,24	1.20	1 (5%)	18,18,25	1.18	2 (11%)
2	OLC	A	403	-	13,13,24	1.21	1 (7%)	14,14,25	1.07	1 (7%)
2	OLC	A	402	-	15,15,24	1.27	1 (6%)	15,15,25	1.47	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OLC	A	409	-	7,7,24	0.72	0	6,6,25	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OLC	A	404	-	-	9/15/15/24	-
2	OLC	A	408	-	-	3/6/6/24	-
2	OLC	A	407	-	-	4/7/7/24	-
2	OLC	A	406	-	-	5/7/7/24	-
3	RET	A	410	1	-	0/13/30/31	0/1/1/1
2	OLC	A	401	-	-	13/24/24/24	-
2	OLC	A	405	-	-	10/17/17/24	-
2	OLC	A	403	-	-	9/13/13/24	-
2	OLC	A	402	-	-	7/13/13/24	-
2	OLC	A	409	-	-	3/5/5/24	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	OLC	O20-C1	3.35	1.42	1.30
2	A	406	OLC	O20-C1	3.34	1.42	1.30
2	A	407	OLC	O20-C1	3.30	1.41	1.30
2	A	404	OLC	O20-C1	3.11	1.42	1.33
2	A	401	OLC	O20-C1	3.03	1.42	1.33
2	A	405	OLC	O20-C1	2.99	1.42	1.33
2	A	403	OLC	O20-C1	2.95	1.42	1.33
3	A	410	RET	C14-C13	2.88	1.36	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	408	OLC	C8-C9-C10	3.81	152.75	126.84
2	A	402	OLC	C8-C9-C10	3.70	153.13	124.73
2	A	401	OLC	C8-C9-C10	3.69	153.03	124.73
2	A	405	OLC	C8-C9-C10	2.72	152.66	131.07
2	A	404	OLC	O20-C1-C2	2.71	120.42	111.91
2	A	403	OLC	O20-C1-C2	2.64	120.18	111.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	OLC	O20-C1-C2	2.56	119.95	111.91
2	A	405	OLC	O20-C1-C2	2.55	119.90	111.91
3	A	410	RET	C8-C7-C6	2.03	132.89	127.20

There are no chirality outliers.

All (63) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	404	OLC	C21-C22-C24-O25
2	A	404	OLC	O20-C21-C22-C24
2	A	401	OLC	O19-C1-O20-C21
2	A	401	OLC	C2-C1-O20-C21
2	A	404	OLC	O20-C21-C22-O23
2	A	403	OLC	O23-C22-C24-O25
2	A	401	OLC	C1-C2-C3-C4
2	A	407	OLC	C1-C2-C3-C4
2	A	401	OLC	C11-C12-C13-C14
2	A	402	OLC	C5-C6-C7-C8
2	A	402	OLC	C3-C4-C5-C6
2	A	401	OLC	C14-C15-C16-C17
2	A	403	OLC	C21-C22-C24-O25
2	A	404	OLC	C3-C4-C5-C6
2	A	409	OLC	C4-C5-C6-C7
2	A	401	OLC	C13-C14-C15-C16
2	A	401	OLC	C3-C4-C5-C6
2	A	405	OLC	C2-C3-C4-C5
2	A	406	OLC	C3-C4-C5-C6
2	A	407	OLC	C4-C5-C6-C7
2	A	404	OLC	O23-C22-C24-O25
2	A	401	OLC	C6-C7-C8-C9
2	A	405	OLC	C6-C7-C8-C9
2	A	405	OLC	C1-C2-C3-C4
2	A	403	OLC	C1-C2-C3-C4
2	A	406	OLC	C1-C2-C3-C4
2	A	407	OLC	C3-C4-C5-C6
2	A	401	OLC	C2-C3-C4-C5
2	A	401	OLC	C10-C11-C12-C13
2	A	402	OLC	C6-C7-C8-C9
2	A	408	OLC	C6-C7-C8-C9
2	A	403	OLC	C4-C5-C6-C7
2	A	405	OLC	C7-C8-C9-C10
2	A	405	OLC	C4-C5-C6-C7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	408	OLC	C3-C4-C5-C6
2	A	405	OLC	C5-C6-C7-C8
2	A	403	OLC	O20-C21-C22-O23
2	A	404	OLC	C6-C7-C8-C9
2	A	409	OLC	C3-C4-C5-C6
2	A	403	OLC	O20-C21-C22-C24
2	A	403	OLC	C2-C1-O20-C21
2	A	406	OLC	C4-C5-C6-C7
2	A	403	OLC	O19-C1-O20-C21
2	A	401	OLC	C21-C22-C24-O25
2	A	402	OLC	C4-C5-C6-C7
2	A	402	OLC	C2-C3-C4-C5
2	A	403	OLC	O20-C1-C2-C3
2	A	401	OLC	O23-C22-C24-O25
2	A	409	OLC	C5-C6-C7-C8
2	A	401	OLC	C7-C8-C9-C10
2	A	408	OLC	C7-C8-C9-C10
2	A	404	OLC	C5-C6-C7-C8
2	A	406	OLC	O20-C1-C2-C3
2	A	402	OLC	C9-C10-C11-C12
2	A	406	OLC	O19-C1-C2-C3
2	A	407	OLC	C5-C6-C7-C8
2	A	405	OLC	O20-C1-C2-C3
2	A	405	OLC	O23-C22-C24-O25
2	A	404	OLC	C2-C1-O20-C21
2	A	405	OLC	C3-C4-C5-C6
2	A	404	OLC	O19-C1-O20-C21
2	A	405	OLC	O19-C1-C2-C3
2	A	402	OLC	C7-C8-C9-C10


There are no ring outliers.




4 monomers are involved in 7 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	410	RET	4	0
2	A	405	OLC	1	0
2	A	403	OLC	1	0
2	A	409	OLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/333 (88%)	0.10	12 (4%) 42 36	18, 51, 115, 144	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	THR	3.4
1	A	327	ASN	3.3
1	A	118	CYS	3.3
1	A	348	VAL	3.0
1	A	333	ILE	2.9
1	A	211	TYR	2.8
1	A	256	TRP	2.6
1	A	111	TYR	2.5
1	A	343	ALA	2.5
1	A	276	PHE	2.4
1	A	89	LEU	2.4
1	A	58	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OLC	A	403	14/25	0.50	0.20	65,106,123,133	0
2	OLC	A	402	16/25	0.73	0.20	64,86,90,90	0
2	OLC	A	409	8/25	0.74	0.20	66,69,75,84	0
2	OLC	A	405	18/25	0.78	0.16	36,77,107,110	0
2	OLC	A	406	10/25	0.82	0.13	38,65,102,104	0
2	OLC	A	407	10/25	0.83	0.15	50,70,86,93	0
2	OLC	A	401	25/25	0.84	0.17	45,72,89,94	0
2	OLC	A	404	16/25	0.84	0.14	29,51,85,92	0
2	OLC	A	408	9/25	0.88	0.11	40,50,57,60	0
3	RET	A	410	20/21	0.92	0.08	33,46,57,57	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around OLC A 409:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 405:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 407:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 404:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 408:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around RET A 410:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.