



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 10:50 AM EDT

PDB ID : 4GU8  
Title : Crystal Structure of Burkholderia oklahomensis agglutinin (BOA)  
Authors : Whitley, M.J.; Furey, W.; Gronenborn, A.M.  
Deposited on : 2012-08-29  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	 3% 94% 6%
1	B	279	 3% 97% .
1	C	279	 2% 95% ..
1	D	279	 6% 97% .
1	E	279	 4% 94% 5% .

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	279	 4% 97%
1	G	279	 5% 95%
1	H	279	 5% 96%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	301	-	-	-	X
2	GOL	A	303	-	-	-	X
2	GOL	D	301	-	-	-	X
2	GOL	D	302	-	-	-	X
3	DTT	E	301	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 17085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Burkholderia oklahomensis agglutinin (BOA).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	279	2046	1270	365	401	10	0	0	0
1	B	279	2046	1270	365	401	10	0	0	0
1	C	275	2016	1253	361	393	9	0	0	0
1	D	279	2050	1273	366	401	10	0	1	0
1	E	275	2016	1253	361	393	9	0	0	0
1	F	276	2029	1261	363	395	10	0	1	0
1	G	275	2016	1253	361	393	9	0	0	0
1	H	276	2029	1261	363	395	10	0	1	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0

- Molecule 3 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (three-letter code: DTT) (formula:  $C_4H_{10}O_2S_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	E	1	Total	C	O	S	0	0
			8	4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	131	Total	O	0	0
			131	131		
4	B	109	Total	O	0	0
			109	109		
4	C	118	Total	O	0	0
			118	118		
4	D	117	Total	O	0	0
			117	117		
4	E	87	Total	O	0	0
			87	87		
4	F	98	Total	O	0	0
			98	98		
4	G	68	Total	O	0	0
			68	68		
4	H	71	Total	O	0	0
			71	71		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)




- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)




- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)



- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)




- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)




- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)

Chain F:  4% 97%



- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)

Chain G:  5% 95%



- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)

Chain H:  5% 96%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.72Å 130.58Å 149.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.41 – 2.40 18.41 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (18.41-2.40) 97.9 (18.41-2.40)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.03 (at 2.40Å)	Xtrriage
Refinement program	BUSTER-TNT, BUSTER 2.10.0	Depositor
R, $R_{free}$	0.194 , 0.229 0.189 , 0.227	Depositor DCC
$R_{free}$ test set	4601 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.0	Xtrriage
Anisotropy	0.742	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 42.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	17085	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DTT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/2100	0.66	0/2858
1	B	0.52	0/2100	0.67	0/2858
1	C	0.52	0/2070	0.66	0/2819
1	D	0.51	0/2107	0.64	0/2869
1	E	0.53	0/2070	0.67	0/2819
1	F	0.50	0/2086	0.64	0/2840
1	G	0.48	0/2070	0.64	0/2819
1	H	0.49	0/2086	0.65	0/2840
All	All	0.51	0/16689	0.65	0/22722

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2046	0	1910	12	0
1	B	2046	0	1910	7	0
1	C	2016	0	1880	7	0
1	D	2050	0	1916	6	0
1	E	2016	0	1880	13	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2029	0	1898	4	0
1	G	2016	0	1880	5	0
1	H	2029	0	1898	4	0
2	A	18	0	24	3	0
2	D	12	0	16	4	0
3	E	8	0	10	9	0
4	A	131	0	0	0	0
4	B	109	0	0	0	0
4	C	118	0	0	1	0
4	D	117	0	0	0	0
4	E	87	0	0	0	0
4	F	98	0	0	0	0
4	G	68	0	0	0	0
4	H	71	0	0	0	0
All	All	17085	0	15222	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:ALA:HB1	2:A:301:GOL:H2	1.33	1.05
1:A:198:GLY:HA3	2:D:302:GOL:H12	1.48	0.95
1:A:249:GLY:HA2	2:D:301:GOL:H12	1.48	0.93
1:E:209:GLY:HA3	3:E:301:DTT:H2	1.69	0.75
1:D:182:THR:HG23	2:D:302:GOL:H2	1.75	0.68
1:A:149:GLU:HG3	2:A:303:GOL:H2	1.77	0.67
1:E:210:ASN:ND2	3:E:301:DTT:H3	2.13	0.64
1:E:210:ASN:OD1	3:E:301:DTT:H12	1.97	0.64
1:E:210:ASN:H	3:E:301:DTT:H41	1.63	0.62
1:E:238:GLY:N	3:E:301:DTT:S1	2.69	0.60
1:G:275:PRO:O	1:G:276:GLN:HB2	2.03	0.59
1:C:17:GLN:NE2	1:C:20:GLY:O	2.36	0.58
1:E:210:ASN:CG	3:E:301:DTT:H12	2.23	0.57
1:A:40:ASN:O	1:A:62:VAL:HG23	2.04	0.57
1:E:104:VAL:HG22	3:E:301:DTT:C2	2.36	0.56
1:E:236:ARG:O	3:E:301:DTT:H11	2.06	0.55
1:A:221:THR:HB	1:D:221:THR:HB	1.90	0.53
1:C:152:TRP:HB3	1:C:267:ILE:HG22	1.90	0.53
1:B:93:HIS:HB3	1:C:258:SER:HB2	1.90	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:46:ILE:HG21	1:E:69:PHE:CD1	2.44	0.52
1:B:152:TRP:HB3	1:B:267:ILE:HG22	1.93	0.51
1:G:152:TRP:HB3	1:G:267:ILE:HG22	1.94	0.50
1:H:152:TRP:HB3	1:H:267:ILE:HG22	1.94	0.49
1:D:152:TRP:HB3	1:D:267:ILE:HG22	1.93	0.49
1:A:152:TRP:HB3	1:A:267:ILE:HG22	1.95	0.49
1:H:71:ALA:HB1	1:H:79:TYR:HB3	1.95	0.48
1:B:71:ALA:HB1	1:B:79:TYR:HB3	1.95	0.48
1:A:42:VAL:HG13	1:A:62:VAL:HG22	1.97	0.47
1:A:71:ALA:HB1	1:A:79:TYR:HB3	1.97	0.47
1:B:53:ARG:HH11	1:B:53:ARG:HG2	1.80	0.46
1:F:71:ALA:HB1	1:F:79:TYR:HB3	1.97	0.46
1:B:84:GLN:HB2	1:B:92:TRP:CD2	2.52	0.45
1:D:71:ALA:HB1	1:D:79:TYR:HB3	1.98	0.45
1:C:71:ALA:HB1	1:C:79:TYR:HB3	1.99	0.45
1:G:258:SER:HB2	1:H:93:HIS:HB3	1.98	0.45
1:C:226:HIS:HD2	4:C:325:HOH:O	1.99	0.45
1:G:71:ALA:HB1	1:G:79:TYR:HB3	1.98	0.45
1:E:50:ASP:HB2	1:H:196:GLY:HA3	1.99	0.44
1:A:3:LYS:O	1:A:9:ASN:HB3	2.18	0.44
1:A:266:PRO:HG3	1:C:53:ARG:HH22	1.83	0.44
1:E:71:ALA:HB1	1:E:79:TYR:HB3	2.00	0.43
1:B:84:GLN:HB2	1:B:92:TRP:CE3	2.53	0.43
1:E:104:VAL:HG22	3:E:301:DTT:H2	2.00	0.43
1:C:17:GLN:HG2	1:C:25:TRP:CZ2	2.54	0.43
1:F:246:ILE:HD12	1:F:255:PHE:CD2	2.54	0.43
1:E:14:VAL:HG22	1:E:138:SER:HB3	2.01	0.42
1:B:-2:SER:N	1:F:92:TRP:O	2.52	0.42
1:D:258:SER:OG	2:D:301:GOL:H11	2.19	0.42
1:D:93:HIS:HB3	1:F:258:SER:HB2	2.02	0.41
1:G:201:GLY:O	1:G:216:ASN:HA	2.20	0.41
1:A:103:ARG:HD3	2:A:302:GOL:H31	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/279 (99%)	271 (98%)	6 (2%)	0	100	100
1	B	277/279 (99%)	273 (99%)	4 (1%)	0	100	100
1	C	273/279 (98%)	268 (98%)	5 (2%)	0	100	100
1	D	278/279 (100%)	273 (98%)	5 (2%)	0	100	100
1	E	273/279 (98%)	269 (98%)	4 (2%)	0	100	100
1	F	275/279 (99%)	269 (98%)	6 (2%)	0	100	100
1	G	273/279 (98%)	267 (98%)	6 (2%)	0	100	100
1	H	275/279 (99%)	269 (98%)	5 (2%)	1 (0%)	34	48
All	All	2201/2232 (99%)	2159 (98%)	41 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	155	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/206 (100%)	206 (100%)	0	100	100
1	B	206/206 (100%)	206 (100%)	0	100	100
1	C	202/206 (98%)	201 (100%)	1 (0%)	88	95
1	D	207/206 (100%)	207 (100%)	0	100	100
1	E	202/206 (98%)	200 (99%)	2 (1%)	76	88
1	F	204/206 (99%)	204 (100%)	0	100	100
1	G	202/206 (98%)	201 (100%)	1 (0%)	88	95
1	H	204/206 (99%)	204 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1633/1648 (99%)	1629 (100%)	4 (0%)	93 97

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	17	GLN
1	E	76	SER
1	E	178	SER
1	G	56	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	210	ASN
1	A	240	ASN
1	B	226	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	DTT	E	301	-	7,7,7	1.19	0	4,8,8	1.93	2 (50%)
2	GOL	A	301	-	5,5,5	0.70	0	5,5,5	1.03	0
2	GOL	D	302	-	5,5,5	0.68	0	5,5,5	0.77	0
2	GOL	D	301	-	5,5,5	0.40	0	5,5,5	0.61	0
2	GOL	A	302	-	5,5,5	0.44	0	5,5,5	1.10	0
2	GOL	A	303	-	5,5,5	0.69	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DTT	E	301	-	-	5/8/8/8	-
2	GOL	A	301	-	-	3/4/4/4	-
2	GOL	D	302	-	-	2/4/4/4	-
2	GOL	D	301	-	-	2/4/4/4	-
2	GOL	A	302	-	-	4/4/4/4	-
2	GOL	A	303	-	-	3/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	301	DTT	O2-C2-C3	-2.39	104.56	109.57
3	E	301	DTT	O3-C3-C2	-2.10	105.16	109.57

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	302	GOL	O1-C1-C2-C3
2	A	303	GOL	C1-C2-C3-O3
2	A	303	GOL	O2-C2-C3-O3
2	D	301	GOL	C1-C2-C3-O3
2	D	301	GOL	O2-C2-C3-O3
2	D	302	GOL	C1-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	E	301	DTT	C1-C2-C3-O3
3	E	301	DTT	C1-C2-C3-C4
3	E	301	DTT	O2-C2-C3-O3
3	E	301	DTT	O2-C2-C3-C4
2	A	301	GOL	C1-C2-C3-O3
2	A	302	GOL	C1-C2-C3-O3
2	A	301	GOL	O1-C1-C2-O2
2	A	302	GOL	O1-C1-C2-O2
2	A	302	GOL	O2-C2-C3-O3
2	D	302	GOL	O2-C2-C3-O3
2	A	303	GOL	O1-C1-C2-O2
2	A	301	GOL	O1-C1-C2-C3
3	E	301	DTT	S1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	301	DTT	9	0
2	A	301	GOL	1	0
2	D	302	GOL	2	0
2	D	301	GOL	2	0
2	A	302	GOL	1	0
2	A	303	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	279/279 (100%)	-0.35	8 (2%) 51 50	31, 42, 71, 88	0
1	B	279/279 (100%)	-0.26	8 (2%) 51 50	35, 49, 67, 94	0
1	C	275/279 (98%)	-0.26	6 (2%) 62 60	35, 47, 73, 102	0
1	D	279/279 (100%)	-0.11	16 (5%) 23 22	37, 53, 81, 95	0
1	E	275/279 (98%)	0.01	12 (4%) 34 33	33, 55, 87, 102	0
1	F	276/279 (98%)	-0.13	11 (3%) 38 37	39, 54, 82, 112	0
1	G	275/279 (98%)	0.10	13 (4%) 31 30	44, 62, 92, 112	0
1	H	276/279 (98%)	0.01	14 (5%) 28 26	43, 64, 85, 99	0
All	All	2214/2232 (99%)	-0.13	88 (3%) 38 37	31, 54, 82, 112	0

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	19	GLY	5.6
1	E	18	TRP	5.1
1	A	154	GLY	4.9
1	D	155	SER	4.8
1	G	153	GLY	4.7
1	D	19	GLY	4.4
1	E	116	GLY	4.2
1	H	155	SER	4.1
1	F	222	SER	4.0
1	F	155	SER	3.9
1	F	156	SER	3.9
1	D	89	SER	3.9
1	H	19	GLY	3.8
1	B	155	SER	3.8
1	B	156	SER	3.8
1	E	153	GLY	3.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	F	1	MET	3.7
1	C	155	SER	3.7
1	F	221	THR	3.5
1	G	218	TRP	3.5
1	C	153	GLY	3.5
1	D	130	GLY	3.4
1	E	20	GLY	3.3
1	A	-2	SER	3.3
1	H	105	ASN	3.3
1	D	90	ALA	3.2
1	B	-1	VAL	3.1
1	H	156	SER	3.1
1	B	157	ALA	3.1
1	G	196	GLY	3.1
1	G	222	SER	3.1
1	E	91	PRO	3.0
1	F	153	GLY	3.0
1	G	154	GLY	3.0
1	A	155	SER	3.0
1	B	154	GLY	2.8
1	D	-1	VAL	2.8
1	A	0	ASP	2.8
1	H	196	GLY	2.8
1	G	207	VAL	2.7
1	C	21	SER	2.7
1	D	-2	SER	2.7
1	H	18	TRP	2.6
1	B	-2	SER	2.6
1	F	63	GLY	2.6
1	E	130	GLY	2.6
1	G	225	TRP	2.6
1	D	22	SER	2.6
1	G	224	PRO	2.5
1	D	86	GLY	2.5
1	H	152	TRP	2.5
1	G	276	GLN	2.5
1	A	156	SER	2.5
1	D	156	SER	2.5
1	C	63	GLY	2.5
1	E	25	TRP	2.5
1	B	222	SER	2.5
1	E	23	ALA	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	H	221	THR	2.4
1	C	154	GLY	2.4
1	F	220	GLY	2.4
1	H	88	SER	2.4
1	D	154	GLY	2.4
1	A	157	ALA	2.4
1	A	-1	VAL	2.4
1	D	223	ALA	2.4
1	H	130	GLY	2.3
1	D	63	GLY	2.3
1	F	218	TRP	2.3
1	G	38	GLY	2.3
1	G	195	ALA	2.3
1	C	41	VAL	2.2
1	H	23	ALA	2.2
1	G	155	SER	2.2
1	A	19	GLY	2.2
1	E	21	SER	2.2
1	B	153	GLY	2.2
1	E	89	SER	2.1
1	D	0	ASP	2.1
1	G	105	ASN	2.1
1	H	25	TRP	2.1
1	E	93	HIS	2.1
1	D	88	SER	2.1
1	H	276	GLN	2.1
1	F	152	TRP	2.0
1	F	224	PRO	2.0
1	D	85	TRP	2.0
1	H	85	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	301	6/6	0.66	0.47	46,50,55,56	6
2	GOL	D	302	6/6	0.68	0.62	38,41,43,45	6
2	GOL	D	301	6/6	0.73	0.40	57,58,59,60	6
2	GOL	A	303	6/6	0.79	0.44	36,46,47,48	6
3	DTT	E	301	8/8	0.81	0.46	45,49,51,55	8
2	GOL	A	302	6/6	0.91	0.16	62,67,68,71	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.