



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 05:55 pm BST

PDB ID : 6GUE
Title : CDK2/CyclinA in complex with AZD5438
Authors : Wood, D.J.; Korolchuk, S.; Tatum, N.J.; Wang, L.Z.; Endicott, J.A.; Noble, M.E.M.; Martin, M.P.
Deposited on : 2018-06-19
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	
2	B	268	
2	D	268	
3	C	302	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 9499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	294	2363	1536	402	416	1	8	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P24941
A	-2	PRO	-	expression tag	UNP P24941
A	-1	GLY	-	expression tag	UNP P24941
A	0	SER	-	expression tag	UNP P24941
A	73	ASN	GLU	conflict	UNP P24941

- Molecule 2 is a protein called Cyclin-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	262	2112	1367	344	391	10	0	0	0
2	D	262	2112	1367	344	391	10	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	433	HIS	-	expression tag	UNP P30274
B	434	HIS	-	expression tag	UNP P30274
B	435	HIS	-	expression tag	UNP P30274
B	436	HIS	-	expression tag	UNP P30274
B	437	HIS	-	expression tag	UNP P30274
B	438	HIS	-	expression tag	UNP P30274
D	433	HIS	-	expression tag	UNP P30274
D	434	HIS	-	expression tag	UNP P30274
D	435	HIS	-	expression tag	UNP P30274

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	436	HIS	-	expression tag	UNP P30274
D	437	HIS	-	expression tag	UNP P30274
D	438	HIS	-	expression tag	UNP P30274


- Molecule 3 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
3	C	289	2321	1502	393	417	1	8	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	expression tag	UNP P24941
C	-2	PRO	-	expression tag	UNP P24941
C	-1	GLY	-	expression tag	UNP P24941
C	0	SER	-	expression tag	UNP P24941

- Molecule 4 is 4-(2-methyl-3-propan-2-yl-imidazol-4-yl)- {N}-(4-methylsulfonylphenyl)pyrimidin-2-amine (three-letter code: FB8) (formula: C₁₈H₂₁N₅O₂S) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	26	18	5	2	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	C	1	26	18	5	2	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	179	Total 179	O 179	0	0
5	B	141	Total 141	O 141	0	0
5	C	114	Total 114	O 114	0	0
5	D	105	Total 105	O 105	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclin-dependent kinase 2




- Molecule 2: Cyclin-A2



- Molecule 2: Cyclin-A2



- Molecule 3: Cyclin-dependent kinase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.27Å 133.22Å 148.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	99.26 – 1.99 74.07 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.5 (99.26-1.99) 98.5 (74.07-1.99)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.98Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.197 , 0.234 0.204 , 0.240	Depositor DCC
R_{free} test set	4909 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtrriage
Anisotropy	0.127	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9499	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, FB8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	0/2412	0.92	0/3271
2	B	0.82	0/2162	0.82	0/2940
2	D	0.76	0/2162	0.83	0/2940
3	C	0.82	0/2364	0.87	0/3201
All	All	0.84	0/9100	0.86	0/12352

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2363	0	2412	25	0
2	B	2112	0	2130	11	0
2	D	2112	0	2130	10	0
3	C	2321	0	2362	53	0
4	A	26	0	0	0	0
4	C	26	0	0	0	0
5	A	179	0	0	2	0
5	B	141	0	0	6	0
5	C	114	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	105	0	0	2	0
All	All	9499	0	9034	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:121:HIS:O	3:C:122:ARG:HG3	1.49	1.13
3:C:237:LYS:HB3	3:C:238:PRO:HD2	1.30	1.10
2:D:312:ASN:HB3	5:D:552:HOH:O	1.50	1.09
3:C:237:LYS:HB3	3:C:238:PRO:CD	1.87	1.04
1:A:121:HIS:O	1:A:122:ARG:HG3	1.56	1.02
2:B:327:CYS:SG	5:B:621:HOH:O	2.22	0.98
3:C:237:LYS:CB	3:C:238:PRO:CD	2.43	0.97
2:B:334:MET:SD	5:B:622:HOH:O	2.26	0.92
5:A:481:HOH:O	2:B:171:SER:HB2	1.71	0.90
3:C:223:ASP:O	3:C:230:VAL:HG12	1.73	0.87
2:D:346:PRO:HB2	2:D:349:LYS:HE3	1.54	0.87
2:B:334:MET:CG	5:B:622:HOH:O	2.23	0.85
2:B:334:MET:HG3	5:B:622:HOH:O	1.77	0.84
3:C:71:HIS:CD2	2:D:296:HIS:NE2	2.48	0.81
3:C:237:LYS:HG3	3:C:238:PRO:HD3	1.62	0.80
3:C:237:LYS:CG	3:C:238:PRO:HD3	2.12	0.79
3:C:223:ASP:OD1	3:C:224:GLU:HG2	1.87	0.75
2:D:241:ARG:HH22	2:D:305:ASP:HB3	1.52	0.75
2:D:346:PRO:HB2	2:D:349:LYS:CE	2.16	0.74
1:A:128:LEU:O	1:A:192:ILE:HD11	1.88	0.73
3:C:237:LYS:CB	3:C:238:PRO:HD2	2.09	0.73
3:C:223:ASP:O	3:C:230:VAL:CG1	2.37	0.72
1:A:121:HIS:C	1:A:122:ARG:HG3	2.12	0.70
1:A:175:LEU:CD1	1:A:233:MET:HE1	2.22	0.70
3:C:222:PRO:O	3:C:224:GLU:N	2.26	0.69
3:C:84:HIS:HB3	5:C:497:HOH:O	1.91	0.69
3:C:128:LEU:O	3:C:192:ILE:HD11	1.93	0.69
3:C:237:LYS:CG	3:C:238:PRO:CD	2.72	0.68
1:A:177:CYS:SG	1:A:233:MET:CE	2.82	0.68
3:C:230:VAL:HG22	3:C:236:TYR:HE1	1.58	0.67
2:D:241:ARG:NH2	2:D:305:ASP:HB3	2.09	0.67
3:C:121:HIS:C	3:C:122:ARG:HG3	2.15	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:223:ASP:CG	3:C:224:GLU:N	2.49	0.66
3:C:237:LYS:CB	3:C:238:PRO:HD3	2.26	0.65
1:A:223:ASP:H	1:A:226:VAL:HG12	1.62	0.63
3:C:71:HIS:HD2	2:D:296:HIS:NE2	1.96	0.63
2:B:183:HIS:HB2	2:B:317:GLN:HE22	1.63	0.63
3:C:15:TYR:HD2	3:C:35:ILE:HG12	1.64	0.62
1:A:246:GLN:O	1:A:247:ASP:C	2.37	0.62
3:C:198:THR:O	3:C:199:ARG:HB2	2.00	0.62
3:C:15:TYR:CD2	3:C:35:ILE:HG12	2.35	0.61
3:C:230:VAL:HG22	3:C:236:TYR:CE1	2.36	0.60
1:A:175:LEU:HD12	1:A:233:MET:HE1	1.83	0.58
1:A:246:GLN:O	1:A:247:ASP:O	2.22	0.58
3:C:38:ASP:HB3	3:C:41:THR:OG1	2.03	0.57
3:C:209:ILE:HD13	3:C:237:LYS:HE3	1.86	0.57
1:A:175:LEU:HB2	1:A:233:MET:CE	2.35	0.57
1:A:71:HIS:CD2	2:B:296:HIS:CE1	2.93	0.57
1:A:177:CYS:SG	1:A:233:MET:HE3	2.47	0.55
3:C:22:ARG:HG2	3:C:29:VAL:HG22	1.89	0.55
1:A:33:LYS:NZ	1:A:51:GLU:OE1	2.40	0.54
3:C:212:LEU:HD22	3:C:216:PHE:CZ	2.43	0.54
1:A:198:THR:O	1:A:199:ARG:HB2	2.06	0.53
1:A:256:ASP:O	1:A:260:ARG:HG3	2.08	0.53
1:A:95:ALA:O	1:A:199:ARG:NH1	2.41	0.53
3:C:22:ARG:CZ	3:C:22:ARG:HB2	2.39	0.52
2:D:429:THR:HB	5:D:575:HOH:O	2.10	0.52
1:A:218:THR:O	1:A:246:GLN:HG3	2.09	0.51
3:C:15:TYR:HE2	3:C:35:ILE:HD11	1.75	0.51
3:C:238:PRO:HB3	5:C:499:HOH:O	2.10	0.51
1:A:177:CYS:SG	1:A:233:MET:HE2	2.51	0.50
3:C:245:ARG:O	3:C:245:ARG:HG3	2.12	0.50
3:C:247:ASP:OD1	3:C:249:SER:N	2.45	0.49
3:C:94:SER:O	3:C:199:ARG:CD	2.60	0.49
3:C:22:ARG:HB2	3:C:22:ARG:NH1	2.28	0.48
3:C:94:SER:O	3:C:199:ARG:HD3	2.13	0.48
3:C:223:ASP:OD2	3:C:224:GLU:N	2.47	0.47
3:C:247:ASP:CG	3:C:249:SER:H	2.18	0.47
2:D:345:ASP:HA	2:D:346:PRO:HA	1.46	0.47
3:C:14:THR:HG21	3:C:148:LEU:CD1	2.45	0.46
1:A:0:SER:N	5:A:409:HOH:O	2.46	0.45
3:C:62:ASN:C	3:C:142:LYS:HB3	2.37	0.45
3:C:162:GLU:HG2	3:C:169:ARG:CZ	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:22:ARG:CB	3:C:22:ARG:CZ	2.95	0.44
3:C:206:ASP:HA	5:C:418:HOH:O	2.16	0.44
2:B:233:HIS:HD2	5:B:603:HOH:O	2.00	0.44
3:C:237:LYS:HB2	3:C:240:PHE:CE1	2.52	0.44
1:A:175:LEU:HB2	1:A:233:MET:HE1	2.00	0.43
1:A:97:THR:OG1	1:A:98:GLY:O	2.31	0.43
1:A:71:HIS:NE2	2:B:296:HIS:CE1	2.87	0.43
1:A:25:LEU:HD21	2:D:293:ARG:HB3	2.00	0.43
3:C:197:VAL:HG11	3:C:252:VAL:CG1	2.49	0.43
3:C:165:THR:HG22	5:C:460:HOH:O	2.19	0.43
1:A:163:VAL:O	1:A:164:VAL:HB	2.19	0.42
2:B:323:GLN:HA	2:B:323:GLN:NE2	2.34	0.42
3:C:175:LEU:HD13	3:C:233:MET:CE	2.50	0.42
2:B:183:HIS:HD2	5:B:512:HOH:O	2.02	0.42
1:A:175:LEU:HD13	1:A:233:MET:HE1	1.97	0.41
3:C:128:LEU:O	3:C:192:ILE:CD1	2.66	0.41
3:C:162:GLU:CG	3:C:162:GLU:O	2.69	0.40
3:C:94:SER:O	3:C:199:ARG:HD2	2.21	0.40
3:C:198:THR:O	3:C:199:ARG:CB	2.67	0.40
3:C:237:LYS:CG	3:C:238:PRO:HD2	2.48	0.40
3:C:236:TYR:CE2	3:C:240:PHE:HB2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/302 (96%)	279 (96%)	8 (3%)	2 (1%)	22	16
2	B	260/268 (97%)	258 (99%)	2 (1%)	0	100	100
2	D	260/268 (97%)	257 (99%)	3 (1%)	0	100	100
3	C	282/302 (93%)	270 (96%)	8 (3%)	4 (1%)	11	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1091/1140 (96%)	1064 (98%)	21 (2%)	6 (0%)	29 23

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	247	ASP
3	C	223	ASP
1	A	164	VAL
3	C	164	VAL
3	C	238	PRO
3	C	237	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	258/264 (98%)	242 (94%)	16 (6%)	18 13
2	B	235/241 (98%)	232 (99%)	3 (1%)	69 74
2	D	235/241 (98%)	230 (98%)	5 (2%)	53 57
3	C	255/264 (97%)	237 (93%)	18 (7%)	14 10
All	All	983/1010 (97%)	941 (96%)	42 (4%)	29 26

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLU
1	A	51	GLU
1	A	74	ASN
1	A	83	LEU
1	A	89	LYS
1	A	96	LEU
1	A	97	THR
1	A	122	ARG
1	A	150	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	161	HIS
1	A	206	ASP
1	A	237	LYS
1	A	246	GLN
1	A	247	ASP
1	A	248	PHE
1	A	255	LEU
2	B	196	LYS
2	B	328	LYS
2	B	391	LEU
3	C	22	ARG
3	C	41	THR
3	C	74	ASN
3	C	122	ARG
3	C	138	GLU
3	C	142	LYS
3	C	150	ARG
3	C	161	HIS
3	C	199	ARG
3	C	206	ASP
3	C	217	ARG
3	C	223	ASP
3	C	236	TYR
3	C	237	LYS
3	C	241	PRO
3	C	247	ASP
3	C	248	PHE
3	C	295	HIS
2	D	201	LYS
2	D	202	LYS
2	D	293	ARG
2	D	334	MET
2	D	391	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	161	HIS
1	A	295	HIS
2	B	183	HIS
2	B	233	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	296	HIS
2	B	317	GLN
2	B	323	GLN
2	B	370	GLN
2	B	378	GLN
2	B	395	HIS
2	B	431	ASN
3	C	59	ASN
3	C	71	HIS
2	D	233	HIS
2	D	254	GLN
2	D	370	GLN
2	D	395	HIS
2	D	431	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TPO	C	160	3	8,10,11	1.72	2 (25%)	10,14,16	1.17	1 (10%)
1	TPO	A	160	1	8,10,11	0.86	0	10,14,16	1.90	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TPO	C	160	3	-	0/9/11/13	-
1	TPO	A	160	1	-	1/9/11/13	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	160	TPO	P-O3P	-2.94	1.43	1.54
3	C	160	TPO	P-O2P	-2.77	1.44	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	TPO	P-OG1-CB	-5.42	106.82	123.21
3	C	160	TPO	P-OG1-CB	-2.35	116.10	123.21

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	160	TPO	CB-OG1-P-O3P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FB8	A	301	-	27,28,28	0.68	1 (3%)	35,41,41	1.25	6 (17%)
4	FB8	C	301	-	27,28,28	0.70	1 (3%)	35,41,41	1.33	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FB8	A	301	-	-	6/18/18/18	0/3/3/3
4	FB8	C	301	-	-	4/18/18/18	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	FB8	C4-N5	2.42	1.41	1.36
4	C	301	FB8	C4-N5	2.26	1.41	1.36

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	301	FB8	O2-S1-C12	-4.51	104.57	108.25
4	A	301	FB8	C2-N5-C16	3.13	128.01	125.50
4	C	301	FB8	C13-S1-C12	2.91	108.02	104.58
4	A	301	FB8	C13-S1-C12	2.77	107.86	104.58
4	C	301	FB8	N1-C2-N5	-2.68	107.76	112.00
4	A	301	FB8	O1-S1-C12	-2.50	106.21	108.25
4	A	301	FB8	N3-C8-N4	2.42	125.14	116.92
4	C	301	FB8	O1-S1-C12	-2.36	106.32	108.25
4	A	301	FB8	N3-C8-N2	-2.35	108.99	116.28
4	A	301	FB8	N1-C2-N5	-2.30	108.36	112.00

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	301	FB8	C14-C12-S1-C13
4	A	301	FB8	C11-C12-S1-C13
4	A	301	FB8	C11-C12-S1-O2
4	C	301	FB8	C14-C12-S1-O2

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	301	FB8	C14-C12-S1-O2
4	C	301	FB8	C11-C12-S1-O2
4	C	301	FB8	C14-C12-S1-C13
4	C	301	FB8	C11-C12-S1-C13
4	A	301	FB8	C17-C16-N5-C4
4	A	301	FB8	C18-C16-N5-C4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	293/302 (97%)	0.45	11 (3%) 40 39	15, 25, 64, 90	0
2	B	262/268 (97%)	0.13	5 (1%) 66 65	17, 31, 53, 79	0
2	D	262/268 (97%)	0.23	7 (2%) 54 53	21, 39, 63, 93	0
3	C	288/302 (95%)	1.10	38 (13%) 3 2	24, 37, 148, 216	0
All	All	1105/1140 (96%)	0.49	61 (5%) 25 24	15, 33, 70, 216	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	236	TYR	21.0
3	C	221	THR	17.3
3	C	223	ASP	15.3
3	C	232	SER	14.5
3	C	233	MET	12.4
3	C	234	PRO	11.4
3	C	222	PRO	11.2
3	C	224	GLU	11.1
3	C	246	GLN	10.8
3	C	245	ARG	10.6
3	C	235	ASP	10.0
3	C	231	THR	8.7
3	C	240	PHE	8.6
3	C	247	ASP	7.9
3	C	238	PRO	7.6
3	C	237	LYS	6.9
3	C	295	HIS	6.4
3	C	239	SER	6.3
3	C	251	VAL	5.5
3	C	296	LEU	5.1
1	A	96	LEU	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	249	SER	5.1
3	C	248	PHE	4.9
3	C	162	GLU	4.5
2	B	432	VAL	4.3
1	A	73	ASN	4.1
3	C	241	PRO	4.0
3	C	253	PRO	3.9
1	A	240	PHE	3.7
2	D	432	VAL	3.6
1	A	14	THR	3.6
2	D	346	PRO	3.4
3	C	230	VAL	3.3
3	C	287	GLN	3.2
3	C	161	HIS	3.2
3	C	250	LYS	3.2
3	C	13	GLY	3.2
3	C	220	GLY	3.1
3	C	293	VAL	3.0
3	C	14	THR	2.7
3	C	269	TYR	2.7
1	A	2	GLU	2.7
2	B	284	ASP	2.6
1	A	298	LEU	2.6
2	D	424	LEU	2.6
2	B	280	TYR	2.5
1	A	15	TYR	2.5
2	D	323	GLN	2.4
2	D	428	GLU	2.4
1	A	246	GLN	2.4
3	C	22	ARG	2.4
2	B	201	LYS	2.4
2	D	284	ASP	2.3
2	D	364	LEU	2.2
1	A	71	HIS	2.2
1	A	161	HIS	2.2
3	C	15	TYR	2.2
2	B	283	ASP	2.1
3	C	252	VAL	2.1
3	C	288	ASP	2.0
1	A	157	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	160	11/12	0.93	0.17	18,21,24,24	0
3	TPO	C	160	11/12	0.98	0.08	23,29,34,35	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FB8	C	301	26/26	0.96	0.11	26,31,39,41	0
4	FB8	A	301	26/26	0.98	0.11	20,26,35,36	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FB8 C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FB8 A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.